

MEASURES AND DIMENSIONS OF JULIA SETS OF SEMI-HYPERBOLIC RATIONAL SEMIGROUPS

HIROKI SUMI AND MARIUSZ URBAŃSKI

ABSTRACT. We consider the dynamics of semi-hyperbolic semigroups generated by finitely many rational maps on the Riemann sphere. Assuming that the nice open set condition holds it is proved that there exists a geometric measure on the Julia set with exponent h equal to the Hausdorff dimension of the Julia set. Both h -dimensional Hausdorff and packing measures are finite and positive on the Julia set and are mutually equivalent with Radon-Nikodym derivatives uniformly separated from zero and infinity. All three fractal dimensions, Hausdorff, packing and box counting are equal. It is also proved that for the canonically associated skew-product map there exists a unique h -conformal measure. Furthermore, it is shown that this conformal measure admits a unique Borel probability absolutely continuous invariant (under the skew-product map) measure. In fact these two measures are equivalent, and the invariant measure is metrically exact, hence ergodic.

Mathematics Subject Classification (2001). Primary 37F35; Secondary 37F15.

1. INTRODUCTION

In this paper, we frequently use the notation from [36]. A “rational semigroup” G is a semigroup generated by a family of non-constant rational maps $g : \hat{\mathbb{C}} \rightarrow \hat{\mathbb{C}}$, where $\hat{\mathbb{C}}$ denotes the Riemann sphere, with the semigroup operation being functional composition. For a rational semigroup G , we set

$$F(G) := \{z \in \hat{\mathbb{C}} \mid G \text{ is normal in a neighborhood of } z\}$$

and

$$J(G) := \hat{\mathbb{C}} \setminus F(G).$$

Date: November 12, 2008.

Key words and phrases. Complex dynamical systems, rational semigroups, semi-hyperbolic semigroups, Julia set, Hausdorff dimension, conformal measure, skew product.

Research of the second author supported in part by the NSF Grant DMS 0700831.

Hiroki Sumi

Department of Mathematics, Graduate School of Science, Osaka University, 1-1 Machikaneyama, Toyonaka, Osaka, 560-0043, Japan

E-mail: sumi@math.sci.osaka-u.ac.jp

Web: <http://www.math.sci.osaka-u.ac.jp/~sumi/>

Mariusz Urbański

Department of Mathematics, University of North Texas, Denton, TX 76203-1430, USA

E-mail: urbanski@unt.edu

Web: <http://www.math.unt.edu/~urbanski/>

$F(G)$ is called the Fatou set of G and $J(G)$ is called the Julia set of G . If G is generated by a family $\{f_i\}_i$, then we write $G = \langle f_1, f_2, \dots \rangle$.

The work on the dynamics of rational semigroups was initiated by Hinkkanen and Martin ([14]), who were interested in the role of the dynamics of polynomial semigroups while studying various one-complex-dimensional moduli spaces for discrete groups, and by F. Ren's group ([50]), who studied such semigroups from the perspective of random complex dynamics. The theory of the dynamics of rational semigroups on $\hat{\mathbb{C}}$ has developed in many directions since the 1990s ([14, 50, 15, 28, 29, 30, 31, 34, 35, 36, 37, 38, 39, 40, 41, 45, 42, 32, 43]).

Since the Julia set $J(G)$ of a rational semigroup G generated by finitely many elements f_1, \dots, f_u has **backward self-similarity**, i.e.,

$$(1.1) \quad J(G) = f_1^{-1}(J(G)) \cup \dots \cup f_u^{-1}(J(G))$$

(see [36]), it can be viewed as a significant generalization and extension of both, the theory of iteration of rational maps (see [23]), and conformal iterated function systems (see [22]). For example, the Sierpiński gasket can be regarded as the Julia set of a rational semigroup. The theory of the dynamics of rational semigroups borrows and develops tools from both of these theories. It has also developed its own unique methods, notably the skew product approach (see [36, 37, 38, 39, 42, 45], and [46]). We remark that by (1.1), the analysis of the Julia sets of rational semigroups somewhat resembles “backward iterated functions systems”, however since each map f_j is not in general injective (critical points), some qualitatively different extra effort in the cases of semigroups is needed.

The theory of the dynamics of rational semigroups is intimately related to that of the random dynamics of rational maps. For the study of random complex dynamics, the reader may consult [13, 4, 5, 3, 2, 16, 24]. We remark that the complex dynamical systems can be used to describe some mathematical models. For example, the behavior of the population of a certain species can be described as the dynamical system of a polynomial $f(z) = az(1-z)$ such that f preserves the unit interval and the postcritical set in the plane is bounded (cf. [10]). From this point of view, it is very important to consider the random dynamics of polynomials. For the random dynamics of polynomials on the unit interval, see [33].

The deep relation between these fields (rational semigroups, random complex dynamics, and (backward) IFS) is explained in detail in the subsequent papers ([40, 41, 42, 43, 44]) of the first author.

In this paper, we investigate the Hausdorff, packing, and box dimension of the Julia sets of semi-hyperbolic rational semigroups $G = \langle f_1, \dots, f_u \rangle$ satisfying the nice open set condition. We will show that these dimensions coincide, that $0 < H^h(J(G)), P^h(J(G)) < \infty$, where h is the Hausdorff dimension of $J(G)$ and H^h (resp. P^h) denotes the h -dimensional Hausdorff (resp. packing) measure, that h is equal to the critical exponent of the Poincaré series of the semigroup G , that there exists a unique h -conformal measure \tilde{m}_h on the Julia set $J(\tilde{f})$ of the “skew product map” \tilde{f} , that there exists a unique Borel probability measure $\tilde{\mu}_h$ on $J(\tilde{f})$ which is absolutely continuous with respect to \tilde{m}_h , and that $\tilde{\mu}_h$ is metrically exact and equivalent with \tilde{m}_h . The precise statements of these results are given in Theorem 1.11. In order to prove these results, we develop and combine the idea of usual iteration of non-recurrent critical point maps ([47]), conformal iterated function systems ([22]), and the

dynamics of expanding rational semigroups ([38]). However, as we mentioned before, since the generators may have critical points in the Julia set, we need some careful treatment on the critical points in the Julia set and some observation on the overlapping of the backward images of the Julia set under the elements of the semigroup.

Our approach develops the methods from [38], [47], and [48]. In order to prove that a conformal measure exists, is atomless, and, ultimately, geometric, we expand the concepts of estimability of measures, which originally appeared in [47], we introduce a partial order in the set of critical points, and a stratification of invariant subsets of the Julia set. As an entirely new tool to all [38], [47], and [48], we introduce the concept of essential families of inverse branches. This concept, supported by the notion of nice open set, is extremely useful in the realm of semi-hyperbolic rational semigroups, as it would also (without nice open set) substantially simplified considerations in the expanding case.

In the second part of the paper, devoted to proving the existence and uniqueness of an invariant (with respect to the canonical skew-product) probability measure equivalent with the h -conformal measure, the most challenging task is to prove the uniqueness of the latter. We do it by bringing up and elaborating the tool of Vitali relations due to Federer (see [12]), the tool which has not come up in [47], [48] nor [38]. We rely here heavily on deep results from [12]. The second tool, already employed in [48] and subsequent papers of the second author, is the Marco Martens method of producing σ -finite invariant measures absolutely continuous with respect to a given quasi invariant measure. We apply and develop this method, proving in particular its validity for abstract measure spaces and not only for σ -compact measure spaces. This is possible because of our use of Banach limits rather than weak convergence of measures.

We remark that as illustrated in [41, 40, 44], estimating the Hausdorff dimension of the Julia sets of rational semigroups plays an important role when we investigate random complex dynamics and its associated Markov process on $\hat{\mathbb{C}}$. For example, when we consider the random dynamics of a compact family Γ of polynomials of degree greater than or equal to two, then the function $T_\infty : \hat{\mathbb{C}} \rightarrow [0, 1]$ of probability of tending to $\infty \in \hat{\mathbb{C}}$ varies only inside the Julia set of rational semigroup generated by Γ , and under some condition, this $T_\infty : \hat{\mathbb{C}} \rightarrow [0, 1]$ is continuous on $\hat{\mathbb{C}}$ and varies precisely on $J(G)$. If the Hausdorff dimension of the Julia set is strictly less than two, then it means that $T_\infty : \hat{\mathbb{C}} \rightarrow [0, 1]$ is a complex version of devil's staircase (Cantor function) ([40, 41, 44]).

In order to present the precise statements of the main result, we give some basic notations. For each meromorphic function φ , we denote by $|\varphi'(z)|_s$ the norm of the derivative with respect to the spherical metric. Moreover, we denote by $CV(\varphi)$ the set of critical values of φ .

Given a set $A \subset \mathbb{C}$ and $r > 0$, the symbol $B(A, r)$ denotes the Euclidean open r -neighborhood of the set A . Moreover, $\text{diam}(A)$ denotes the diameter of A with respect to the Euclidean distance. Moreover, given a subset A of $\hat{\mathbb{C}}$, $B_s(A, r)$ denotes the spherical open r -neighborhood of the set B . Moreover, $\text{diam}_s(A)$ denotes the diameter of A with respect to the spherical distance.

Let $u \in \mathbb{N}$. In this paper, an element of $(\text{Rat})^u$ is called a multi-map.

Let $f = (f_1, \dots, f_u) \in (\text{Rat})^u$ be a multi-map and let $G = \langle f_1, \dots, f_u \rangle$ be the rational semigroup generated by $\{f_1, \dots, f_u\}$. Then, we use the following notation. Let $\Sigma_u := \{1, \dots, u\}^{\mathbb{N}}$ be the space of one-sided sequences of u -symbols endowed with the product topology. This is a compact metric space. Let $\tilde{f} : \Sigma_u \times \hat{\mathbb{C}} \rightarrow \Sigma_u \times \hat{\mathbb{C}}$ be the skew product map associated with $f = (f_1, \dots, f_u)$ given by the formula

$$\tilde{f}(\omega, z) = (\sigma(\omega), f_{\omega_1}(z)),$$

where $(\omega, z) \in \Sigma_u \times \hat{\mathbb{C}}$, $\omega = (\omega_1, \omega_2, \dots)$, and $\sigma : \Sigma_u \rightarrow \Sigma_u$ denotes the shift map. We denote by $p_1 : \Sigma_u \times \hat{\mathbb{C}} \rightarrow \Sigma_u$ the projection onto Σ_u and $p_2 : \Sigma_u \times \hat{\mathbb{C}} \rightarrow \hat{\mathbb{C}}$ the projection onto $\hat{\mathbb{C}}$. That is,

$$p_1(\omega, z) = \omega \quad \text{and} \quad p_2(\omega, z) = z.$$

Under the canonical identification $p_1^{-1}\{\omega\} \cong \hat{\mathbb{C}}$, each fiber $p_1^{-1}\{\omega\}$ is a Riemann surface which is isomorphic to $\hat{\mathbb{C}}$. Let $\Sigma_u^* := \bigcup_{n \in \mathbb{N}} \{1, \dots, u\}^n$. For each $\omega = (\omega_1, \dots, \omega_n) \in \Sigma_u^*$, let $f_\omega := f_{\omega_n} \circ \dots \circ f_{\omega_1}$. Moreover, let $\tau \in \Sigma_u^*$, $x \in \hat{\mathbb{C}}$, and $n \in \mathbb{N}$. Suppose that $z = f_\tau(x)$ is not a critical value of f_τ . Then we denote by $f_{\tau,x}^{-1}$ the inverse branch of f_τ mapping z to x . Furthermore, we denote by $\tilde{f}_{\tau,x}^{-|\tau|}$ the inverse branch of $\tilde{f}^{|\tau|}$ such that $\tilde{f}_{\tau,x}^{-|\tau|}(\omega, y) = (\tau\omega, f_{\tau,x}^{-1}(y))$.

Let $\text{Crit}(\tilde{f}) := \bigcup_{\omega \in \Sigma_u} \{v \in p_1^{-1}\{\omega\} \mid v \text{ is a critical point of } \tilde{f}|_{p_1^{-1}\{\omega\}} \rightarrow p_1^{-1}\{\sigma(\omega)\}\}$ ($\subset \Sigma_u \times \hat{\mathbb{C}}$) be the set of critical points of \tilde{f} . For each $n \in \mathbb{N}$ and $(\omega, z) \in \Sigma_u \times \hat{\mathbb{C}}$, we set $(\tilde{f}^n)'(\omega, z) := (f_{\omega_n} \circ \dots \circ f_{\omega_1})'(z)$.

For each $\omega \in \Sigma_u$ we define

$$J_\omega := \{z \in \hat{\mathbb{C}} \mid \{f_{\omega_n} \circ \dots \circ f_{\omega_1}\}_{n \in \mathbb{N}} \text{ is not normal in any neighborhood of } z\}$$

and we then set

$$J(\tilde{f}) := \overline{\bigcup_{\omega \in \Sigma_u} \{\omega\} \times J_\omega},$$

where the closure is taken in the product space $\Sigma_u \times \hat{\mathbb{C}}$. By definition, $J(\tilde{f})$ is compact. Furthermore, by Proposition 3.2 in [36], $J(\tilde{f})$ is completely invariant under \tilde{f} , \tilde{f} is an open map on $J(\tilde{f})$, $(\tilde{f}, J(\tilde{f}))$ is topologically exact under a mild condition, and $J(\tilde{f})$ is equal to the closure of the set of repelling periodic points of \tilde{f} provided that $\sharp J(G) \geq 3$, where we say that a periodic point (ω, z) of \tilde{f} with period n is repelling if $|(\tilde{f}^n)'(\omega, z)| > 1$. Furthermore,

$$p_2(J(\tilde{f})) = J(G).$$

Definition 1.1. Let G be a rational semigroup and let F be a subset of $\hat{\mathbb{C}}$. We set $G(F) = \bigcup_{g \in G} g(F)$ and $G^{-1}(F) = \bigcup_{g \in G} g^{-1}(F)$. Moreover, we set $G^* = G \cup \{Id\}$, where Id denotes the identity map on $\hat{\mathbb{C}}$. Furthermore, let $E(G) := \{z \in \hat{\mathbb{C}} \mid \sharp \bigcup_{g \in G} g^{-1}(\{z\}) < \infty\}$.

Proposition 1.2 (Proposition 3.2(f) in [36]). *(topological exactness)* Let $G = \langle f_1, \dots, f_u \rangle$ be a finitely generated rational semigroup. Suppose $\sharp J(G) \geq 3$ and $E(G) \subset F(G)$. Then, the action of the semigroup G on the Julia set $J(G)$ is topologically exact, meaning that for every non-empty open set $U \subset J(G)$ there exist $g_1, g_2, \dots, g_n \in G$ such that

$$g_1(U) \cup g_2(U) \cup \dots \cup g_n(U) \supset J(G).$$

Definition 1.3. A rational semigroup G is called *semi-hyperbolic* if and only if there exists an $N \in \mathbb{N}$ and a $\delta > 0$ such that for each $x \in J(G)$ and $g \in G$,

$$\deg(g : V \rightarrow B_s(x, \delta)) \leq N$$

for each connected component V of $g^{-1}(B_s(x, \delta))$.

Definition 1.4. Let $f = (f_1, \dots, f_u) \in (\text{Rat})^u$ be a multi-map and let $G = \langle f_1, \dots, f_u \rangle$. Moreover, let U be a non-empty open subset of $\hat{\mathbb{C}}$. We say that G (or f) satisfies the *open set condition* with U if (f, U) satisfies the following two properties:

- (osc1) $f_1^{-1}(U) \cup f_2^{-1}(U) \cup \dots \cup f_u^{-1}(U) \subset U$,
- (osc2) $f_i^{-1}(U) \cap f_j^{-1}(U) = \emptyset$ whenever $i \neq j$.

Moreover, we say that G (or f) satisfies the *nice open set condition* with U if (f, U) satisfies the above (osc1), (osc2), and the following (osc3).

- (osc3) $\exists(\alpha \in (0, 1)) \forall(0 < r \leq 1) \forall(x \in \bar{U}) \quad l_2(U \cap B_s(x, r)) \geq \alpha l_2(B_s(x, r))$, where l_2 denotes the 2-dimensional Lebesgue measure on $\hat{\mathbb{C}}$.

Remark 1.5. Condition (osc3) is not needed if our semigroup G is expanding (see [38] or note that our proofs would use only (osc1) and (osc2) under this assumption). Condition (osc3) is satisfied in the theory of conformal infinite iterated function systems (see [21], comp. [22]), where it follows from the open set condition and the cone condition. Moreover, condition (osc3) holds for example if the boundary of U is smooth enough; piecewise smooth with no exterior cusps suffices. Furthermore, (osc3) holds if U is a John domain (see [6]).

Definition 1.6 ([38]). Let G be a countable rational semigroup. For any $t \geq 0$ and $z \in \hat{\mathbb{C}}$, we set $S_G(z, t) := \sum_{g \in G} \sum_{g(y)=z} |g'(y)|_s^{-t}$, counting multiplicities. We also set $S_G(z) := \inf\{t \geq 0 : S_G(z, t) < \infty\}$ (if no t exists with $S_G(z, t) < \infty$, then we set $S_G(z) := \infty$). Furthermore, we set $s_0(G) := \inf\{S_G(z) : z \in \hat{\mathbb{C}}\}$. This $s_0(G)$ is called the **critical exponent of the Poincaré series** of G .

Definition 1.7 ([38]). Let $f = (f_1, \dots, f_u) \in (\text{Rat})^u$, $t \geq 0$, and $z \in \hat{\mathbb{C}}$. We put $T_f(z, t) := \sum_{\omega \in \Sigma_u^*} \sum_{f_\omega(y)=z} |f'_\omega(y)|_s^{-t}$, counting multiplicities. Moreover, we set $T_f(z) := \inf\{t \geq 0 : T_f(z, t) < \infty\}$ (if no t exists with $T_f(z, t) < \infty$, then we set $T_f(z) = \infty$). Furthermore, we set $t_0(f) := \inf\{T_f(z) : z \in \hat{\mathbb{C}}\}$. This $t_0(f)$ is called the **critical exponent of the Poincaré series** of $f = (f_1, \dots, f_u) \in (\text{Rat})^u$.

Remark 1.8. Let $f = (f_1, \dots, f_u) \in (\text{Rat})^u$, $t \geq 0$, $z \in \hat{\mathbb{C}}$ and let $G = \langle f_1, \dots, f_u \rangle$. Then, $S_G(t, z) \leq T_f(t, z)$, $S_G(z) \leq T_f(z)$, and $s_0(G) \leq t_0(f)$. Note that for almost every $f \in (\text{Rat})^u$ with respect to the Lebesgue measure, $G = \langle f_1, \dots, f_u \rangle$ is a free semigroup and so we have $S_G(t, z) = T_f(t, z)$, $S_G(z) = T_f(z)$, and $s_0(G) = t_0(f)$.

Definition 1.9. Let $\varphi : J(\tilde{f}) \rightarrow \mathbb{R}$ be a function. Let ν be a Borel probability measure on $J(\tilde{f})$. We say that ν is a φ -conformal measure for the map $\tilde{f} : J(\tilde{f}) \rightarrow J(\tilde{f})$ if for each Borel subset A of $J(\tilde{f})$ such that $\tilde{f}|_A : A \rightarrow J(\tilde{f})$ is injective, we have $\nu(\tilde{f}(A)) = \int_A \varphi d\nu$. A $|\tilde{f}'|_s^t$ -conformal measure ν is sometimes called a t -conformal measure. When $J(G) \subset \mathbb{C}$, a $|\tilde{f}'|^t$ -conformal measure is also sometimes called a t -conformal measure.

Definition 1.10. Let $G = \langle f_1, \dots, f_u \rangle$ and let $t \geq 0$. For all $z \in \hat{\mathbb{C}} \setminus \overline{G^*(\bigcup_{j=1}^u CV(f_j))}$, we set $P_z(t) := \limsup_{n \rightarrow \infty} \frac{1}{n} \log \sum_{|\omega|=n} \sum_{x \in f_\omega^{-1}(z)} |f'_\omega(x)|_s^{-t}$.

The main result of this paper is the following.

Theorem 1.11 (see Lemma 7.2, Lemma 7.3, Theorem 7.16, Corollary 7.18 and Theorem 8.4). Let $f = (f_1, \dots, f_u) \in (\text{Rat})^u$ be a multi-map. Let $G = \langle f_1, \dots, f_u \rangle$. Suppose that there exists an element g of G such that $\deg(g) \geq 2$, that each element of $\text{Aut}(\hat{\mathbb{C}}) \cap G$ (if this is not empty) is loxodromic, that G is semi-hyperbolic, and that G satisfies the nice open set condition. Then, we have the following.

- (a) $J(G) \cap \overline{G^*(\bigcup_{j=1}^u CV(f_j))}$ is nowhere dense in $J(G)$ and, for each $t \geq 0$, the function $z \mapsto P_z(t)$ is constant throughout a neighborhood of $J(G) \setminus \overline{G^*(\bigcup_{j=1}^u CV(f_j))}$ in $\hat{\mathbb{C}}$. We denote by $P(t)$ the constant.
- (b) The function $t \mapsto P(t)$ has a unique zero. This zero is denoted by $h = h(f)$.
- (c) There exists a unique $|\tilde{f}'|_s^h$ -conformal measure \tilde{m}_h for the map $\tilde{f} : J(\tilde{f}) \rightarrow J(\tilde{f})$.
- (d) Let $m_h := \tilde{m}_h \circ p_2^{-1}$. Then there exists a constant $C \geq 1$ such that

$$C^{-1} \leq \frac{m_h(B_s(z, r))}{r^h} \leq C$$

for all $z \in J(G)$ and all $r \in (0, 1]$.

- (e) $h(f) = \text{HD}(J(G)) = \text{PD}(J(G)) = \text{BD}(J(G))$, where HD, PD, BD denotes the Hausdorff dimension, packing dimension, and box dimension, respectively, with respect to the spherical distance in $\hat{\mathbb{C}}$. Moreover, for each $z \in J(G) \setminus \overline{G^*(\bigcup_{j=1}^u CV(f_j))}$, we have $h(f) = T_f(z) = t_0(f) = S_G(z) = s_0(G)$.
- (f) Let H^h and P^h be the h -dimensional Hausdorff dimension and h -dimensional packing measure respectively. Then, all the measures H^h , P^h , and m_h are mutually equivalent with Radon-Nikodym derivatives uniformly separated away from zero and infinity.
- (g) $0 < H^h(J(G)), P^h(J(G)) < \infty$.
- (h) There exists a unique Borel probability \tilde{f} -invariant measure $\tilde{\mu}_h$ on $J(\tilde{f})$ which is absolutely continuous with respect to \tilde{m}_h . The measure $\tilde{\mu}_h$ is metrically exact and equivalent with \tilde{m}_h .

The proof of Theorem 1.11 will be given in the following Sections 2–8. In Section 9, we give some examples of semi-hyperbolic rational semigroups satisfying the nice open set condition.

2. PRELIMINARIES

2.1. Distortion and Measures. All the points (numbers) appearing in this paper are complex unless it is clear from the context that they are real. In particular x and y are always assumed to be complex numbers and not the real and imaginary parts of a complex number.

Theorem 2.1. (Koebe's $\frac{1}{4}$ -Theorem) If $z \in \mathbb{C}$, $r > 0$ and $H : B(z, r) \rightarrow \mathbb{C}$ is an arbitrary univalent analytic function, then $H(B(z, r)) \supset B(H(z), 4^{-1}|H'(z)|r)$.

Theorem 2.2. (*Koebe's Distortion Theorem, I*) *There exists a function $k : [0, 1) \rightarrow [1, \infty)$ such that for any $z \in \mathbb{C}$, $r > 0$, $t \in [0, 1)$ and any univalent analytic function $H : B(z, r) \rightarrow \mathbb{C}$ we have that*

$$\sup\{|H'(w)| : w \in B(z, tr)\} \leq k(t) \inf\{|H'(w)| : w \in B(z, tr)\}.$$

We put $K = k(1/2)$.

The following is a straightforward consequence of these two distortion theorems.

Lemma 2.3. *Suppose that $D \subset \mathbb{C}$ is an open set, $z \in D$ and $H : D \rightarrow \mathbb{C}$ is an analytic map which has an analytic inverse H_z^{-1} defined on $B(H(z), 2R)$ for some $R > 0$. Then for every $0 \leq r \leq R$*

$$B(z, K^{-1}r|H'(z)|^{-1}) \subset H_z^{-1}(B(H(z), r)) \subset B(z, Kr|H'(z)|^{-1}).$$

We also use the following more geometric versions of Koebe's Distortion Theorems involving moduli of annuli.

Theorem 2.4. (*Koebe's Distortion Theorem, II*) *There exists a function $w : (0, +\infty) \rightarrow [1, \infty)$ such that for any two open topological disks $Q_1 \subset Q_2 \subset \mathbb{C}$ with $\text{Mod}(Q_2 \setminus Q_1) \geq t$ and any univalent analytic function $H : Q_2 \rightarrow \mathbb{C}$ we have*

$$\sup\{|H'(\xi)| : \xi \in Q_1\} \leq w(t) \inf\{|H'(\xi)| : \xi \in Q_1\}.$$

Definition 2.5. *If $f : D \rightarrow \mathbb{C}$ is an analytic map, $z \in \mathbb{C}$, and $r > 0$, then by*

$$\text{Comp}(z, H, r)$$

we denote the connected component of $H^{-1}(B(H(z), r))$ that contains z .

Given an analytic function H defined throughout a region $D \subset \mathbb{C}$, we put

$$\text{Crit}(H) = \{z \in D : H'(z) = 0\}.$$

Suppose now that c is a critical point of an analytic map $H : D \rightarrow \mathbb{C}$. Then there exists $R = R(H, c) > 0$ and $A = A(H, c) \geq 1$ such that

$$A^{-1}|z - c|^q \leq |H(z) - H(c)| \leq A|z - c|^q$$

and

$$A^{-1}|z - c|^{q-1} \leq |H'(z)| \leq A|z - c|^{q-1}$$

for every $z \in \text{Comp}(c, H, R)$, and that

$$H(\text{Comp}(c, H, R)) = B(H(c), R),$$

where $q = q(H, c)$ is the order of H at the critical point c . In particular

$$\text{Comp}(c, H, R) \subset B(c, (AR)^{1/q}).$$

Moreover, by taking $R > 0$ sufficiently small, we can ensure that the above two inequalities hold for every $z \in B(c, (AR)^{1/q})$ and the ball $B(c, (AR)^{1/q})$ can be expressed as a union of q closed topological disks with smooth boundaries and mutually disjoint interiors such that the map H restricted to each of these interiors, is injective.

In the sequel we require the following technical lemma proven in [47] as Lemma 2.11.

Lemma 2.6. *Let $H : D \rightarrow \mathbb{C}$ be an analytic function. Suppose that an analytic map $Q \circ H : D \rightarrow \mathbb{C}$, a radius $R > 0$ and a point $z \in D$ are such that*

$$\text{Comp}(H(z), Q, 2R) \cap \text{Crit}(Q) = \emptyset \text{ and } \text{Comp}(z, Q \circ H, R) \cap \text{Crit}(H) \neq \emptyset. \quad (\text{a})$$

If c belongs to the last intersection, $A = A(H, c)$, and q is the order of H at c , and

$$\text{diam}(\text{Comp}(z, Q \circ H, R)) \leq (AR(H, c))^{1/q}, \quad (\text{b})$$

then

$$|z - c| \leq KA^2|(Q \circ H)'(z)|^{-1}R.$$

Proof. In view of Lemma 2.3

$$\text{Comp}(H(z), Q, R) \subset B(H(z), KR|Q'(H(z))|^{-1}).$$

So, since $H(c) \in \text{Comp}(H(z), Q, R)$, we get

$$H(c) \in B(H(z), KR|Q'(H(z))|^{-1}).$$

Thus, using this and (b) we obtain

$$\begin{aligned} A^{-1}|z - c|^q &\leq |H(z) - H(c)| \\ &\leq KR|Q'(H(z))|^{-1} \\ &= KR|(Q \circ H)'(z)|^{-1}|H'(z)| \\ &\leq KR|(Q \circ H)'(z)|^{-1}A|z - c|^{q-1}. \end{aligned}$$

So, $|z - c| \leq KA^2|(Q \circ H)'(z)|^{-1}R$. □

Developing the appropriate concepts from [47] we now shall define the notions of estimabilities (upper, lower and strongly lower) of measures, and we shall prove some of its properties and consequences.

Definition 2.7. *Suppose m is a Borel finite measure on Borel set $X \subset \mathbb{R}^n$.*

- (1) *(Upper Estimability) The measure m is said to be upper t -estimable at a point $x \in X$ if there exist $L > 0$ and $R > 0$ such that*

$$m(B(x, r)) \leq Lr^t$$

for all $0 \leq r \leq R$. The number L is referred to as the upper estimability constant of the measure m at x and the number R is referred to as the upper estimability radius of the measure m at x . If there exists an $L > 0$ and an $R > 0$ such that the measure

m is upper t -estimable at each point of X with the upper estimability constant L and the upper estimability radius R , the measure m is said to be uniformly upper t -estimable.

- (2) (Lower Estimability) The measure m is said to be lower t -estimable at a point $x \in X$ if there exists an $L > 0$ and an $R > 0$ such that

$$m(B(x, r)) \geq Lr^t$$

for all $0 \leq r \leq R$. The number L is referred to as the upper estimability constant of the measure m at x and the number R is referred to as the upper estimability radius of the measure m at x . If there exists an $L > 0$ and an $R > 0$ such that the measure m is lower t -estimable at each point of X with the lower estimability constant L and the lower estimability radius R , then the measure m is said to be uniformly lower t -estimable.

- (3) (Strongly Lower Estimability) The measure m is said to be strongly lower t -estimable at a point $x \in X$ if there exists an $L > 0$, a $\lambda \in (0, \infty)$, and an $R > 0$ such that

$$m(B(y, \lambda r)) \geq Lr^t$$

for every $y \in B(x, R)$ for all $0 \leq r \leq R$. The number L is referred to as the lower estimability constant of the measure m at x , the number R is referred to as the lower estimability radius of the measure m at x , and λ is referred to as the lower estimability size of the measure m at x . If there exists an $L > 0$, a λ , and an $R > 0$ such that the measure m is strongly lower t -estimable at each point of X with the lower estimability constant L , the lower estimability radius R , and the lower estimability size λ , then the measure m is said to be uniformly lower t -estimable.

Suppose U and V are open subsets of \mathbb{C} , z is a point of U , and $H : U \rightarrow V$ is an analytic map. Fix $t \geq 0$. A pair (m_1, m_2) of finite Borel measures respectively on U and V is called upper t -conformal for H if and only if

$$m_2(H(A)) \geq \int_A |H'|^t dm_1$$

for all Borel sets $A \subset U$ such that the restriction $H|_A$ is injective. The pair (m_1, m_2) is called t -conformal if the above inequality sign can be replaced by equality. We will need the following lemmas.

Lemma 2.8. *Suppose U and V are open subsets of \mathbb{C} and $H : U \rightarrow V$ is an analytic map which has an analytic inverse H_z^{-1} defined on $B(H(z), 2R)$ for some $R > 0$. Suppose (m_1, m_2) is a t -conformal pair of measures for H . Suppose m_2 is strongly lower t -estimable at $H(z)$ with estimability constant L , estimability radius $0 < r_0 \leq R/2$, and the lower estimability size $\lambda \leq 1$. Then the measure m_1 is strongly lower t -estimable at z with lower estimability constant L , lower estimability radius $K^{-1}|H'(z)|^{-1}r_0$, and lower estimability size $K^2\lambda$.*

Proof. Let $0 \leq r \leq r_0$. Consider $x \in B(z, K^{-1}r|H'(z)|^{-1})$. Then by Lemma 2.3 $H(x) \in B(H(z), r)$ and therefore $m_2(B(H(x), \lambda r)) \geq Lr^t$. Since

$$B(H(x), \lambda r) \subset B(H(z), 2r) \subset B(H(z), R)$$

we have

$$H_z^{-1}(B(H(x), \lambda r)) \subset B(x, K\lambda r|H'(z)|^{-1}) = B(x, K^2\lambda(K^{-1}|H'(z)|^{-1}r)).$$

Thus

$$m_1(B(x, K^2\lambda(K^{-1}|H'(z)|^{-1}r))) \geq K^{-t}|H'(z)|^{-t}Lr^t = L(K^{-1}|H'(z)|^{-1}r)^t.$$

The proof is finished. \square

Lemma 2.9. *Suppose U and V are open subsets of \mathbb{C} and $H : U \rightarrow V$ is an analytic map. Let $c \in U$ be a critical point of H of order q . Suppose (m_1, m_2) is a t -conformal pair of measures for H . If m_2 is lower t -estimable at $H(c)$ with estimability constant L and estimability radius $0 < T \leq R(H, c)$, then the measure m_1 is lower t -estimable at c with estimability constant $A^{-2t}L$ and estimability radius $(A(c)T)^{1/q}$.*

Proof. Put $A = A(c)$. Let $0 < r \leq T$. Notice that $B(H(c), r) = H(\text{Comp}(c, H, r))$. If $x \in \text{Comp}(c, H, r)$, then $A^{-1}|x - c|^q \leq |H(x) - H(c)| < r$ which implies that $x \in B(c, (Ar)^{1/q})$. Thus $B(H(c), r) \subset H(B(c, (Ar)^{1/q}))$ and therefore

$$\begin{aligned} Lr^t &\leq m_2(B(H(c), r)) \\ &\leq m_2(H(B(c, (Ar)^{1/q}))) \\ &\leq \int_{B(c, (Ar)^{1/q})} |H'(z)|^t dm_1(z) \\ &\leq \int_{B(c, (Ar)^{1/q})} A^t(|z - c|^{q-1})^t dm_1(z) \\ &\leq A^t(Ar)^{\frac{q-1}{q}t} m_1(B(c, (Ar)^{1/q})). \end{aligned}$$

So, $m_1(B(c, (Ar)^{1/q})) \geq A^{-2t}L((Ar)^{1/q})^t$. \square

Lemma 2.10. *Suppose U and V are open subsets of \mathbb{C} and $H : U \rightarrow V$ is an analytic map. Let $c \in U$ be a critical point of H of order q . Suppose (m_1, m_2) is an upper t -conformal pair of measures for H such that $m_1(\{c\}) = 0$. If m_2 is upper t -estimable at $H(c)$ with estimability constant L and estimability radius $0 < T < R(H, c)$, then the measure m_1 is upper t -estimable at c with estimability constant $q(2A(c)^2)^t(2^{t/q} - 1)^{-1}L$ and estimability radius $(A(c)^{-1}T)^{1/q}$.*

Proof. Put $A = A(c)$. Take any $0 < s \leq T$. then $H(B(c, (A^{-1}s)^{1/q})) \subset B(H(c), s)$. Set $R(c, a, b) = \{z : a \leq |z - c| < b\}$ and abbreviate $R(c, 2^{-1/q}(A^{-1}s)^{1/q}, (A^{-1}s)^{1/q})$ to $R(c)$.

Using our assumptions and the fact that the map H is q -to-1 on $B(c, (A^{-1}s)^{1/q})$, we obtain

$$\begin{aligned}
Ls^t &\geq m_2(B(H(c), s)) \\
&\geq m_2(H(B(c, (A^{-1}s)^{1/q})) \\
&\geq q^{-1} \int_{B(c, (A^{-1}s)^{1/q})} |H'(z)|^t dm_1(z) \\
&\geq q^{-1} \int_{R(c)} |H'(z)|^t dm_1(z) \\
&\geq q^{-1} A^{-t} (2^{-1} A^{-1} s)^{\frac{q-1}{q}t} m_1(R(c)).
\end{aligned}$$

So, $m_1(R(c, 2^{-1/q}(A^{-1}s)^{1/q}, (A^{-1}s)^{1/q})) \leq q2^{t(1-\frac{1}{q})} A^{2t} L((A^{-1}s)^{1/q})^t$ and therefore for any $0 < r \leq T$,

$$\begin{aligned}
m_1(B(c, (A^{-1}r)^{1/q})) &= m_1\left(\bigcup_{n=0}^{\infty} R(c, 2^{-\frac{n+1}{q}}(A^{-1}r)^{1/q}, 2^{-\frac{n}{q}}(A^{-1}r)^{1/q})\right) \\
&= \sum_{n=0}^{\infty} m_1(R(c, 2^{-\frac{1}{q}}(A^{-1}2^{-n}r)^{1/q}, (A^{-1}2^{-n}r)^{1/q})) \\
&\leq q(2^{1-\frac{1}{q}}A^2)^t L \sum_{n=0}^{\infty} (A^{-1}2^{-n}r)^{t/q} \\
&= q(2^{1-\frac{1}{q}}A^2)^t \frac{L}{1-2^{-\frac{t}{q}}} ((A^{-1}r)^{1/q})^t \\
&= q(2A^2)^t (2^{t/q} - 1)^{-1} L((A^{-1}r)^{1/q})^t.
\end{aligned}$$

The proof is finished. \square

Lemma 2.11. *Suppose U and V are open subsets of \mathbb{C} and $H : U \rightarrow V$ is an analytic map. Let $c \in U$ be a critical point of H of order q . Suppose (m_1, m_2) is a t -conformal pair of measures for H . If m_2 is strongly lower t -estimable at $H(c)$ with estimability constant L , estimability radius $0 < T < R(H, c)/3$, and the lower estimability size λ . Then the measure m_1 is strongly lower t -estimable at c with lower estimability constant $\tilde{L} = L \min\{K^{-t}, (A(c)^2\lambda)^{\frac{1-q}{q}t}\}$, lower estimability radius $(A^{-1}T)^{1/q}$, and lower estimability size $\tilde{\lambda} = (2^{q+1}KA^2\lambda)^{1/q}$.*

Proof. As in the proof of the previous lemma put $A = A(c)$. Let $0 < r \leq T$ and let $x \in B(c, (A^{-1}r)^{1/q})$. If $\tilde{\lambda}(A^{-1}r)^{1/q} \geq 2|x - c|$, then

$$\begin{aligned}
B(x, \tilde{\lambda}(A^{-1}r)^{1/q}) &\supset B(c, \tilde{\lambda}(A^{-1}r)^{1/q}/2) \\
&= B(c, (2K)^{1/q}(A\lambda r)^{1/q}) \\
&\supset B(c, (A\lambda r)^{1/q}).
\end{aligned}$$

It follows from the assumptions that m_2 is lower t -estimable at $H(c)$ with lower estimability constant $\lambda^{-t}L$ and lower estimability radius λT . Therefore, in view of Lemma 2.9 the

critical point c is lower t -estimable with lower estimability constant $A^{-2t}\lambda^{-t}L$ and lower estimability radius $(A\lambda T)^{1/q}$. Thus

$$(2.1) \quad \begin{aligned} m_1(B(x, \tilde{\lambda}(A^{-1}r)^{1/q})) &\geq A^{-2t}\lambda^{-t}L(A\lambda r)^{t/q} \\ &= (A^2\lambda)^{\frac{1-q}{q}t}L((A^{-1}r)^{1/q})^t. \end{aligned}$$

So, suppose that

$$(2.2) \quad \tilde{\lambda}(A^{-1}r)^{1/q} < 2|x - c|.$$

Since c is a critical point we have

$$|H'(x)| \geq A^{-1}|x - c|^{q-1} \geq A^{-1}\tilde{\lambda}^{q-1}(A^{-1}r)^{\frac{q-1}{q}}2^{1-q},$$

which means that

$$(2.3) \quad \begin{aligned} \tilde{\lambda}(A^{-1}r)^{1/q} &\geq A^{-1}\tilde{\lambda}^q A^{-1}r 2^{1-q} |H'(x)|^{-1} \\ &= 4K\lambda r |H'(x)|^{-1} \geq K\lambda r |H'(x)|^{-1}. \end{aligned}$$

In view of (2.2)

$$|H(x) - H(c)| \geq A^{-1}|x - c|^q \geq A^{-1}2^{-q}\tilde{\lambda}^q A^{-1}r = 2K\lambda r \geq 2\lambda r$$

which implies that

$$(2.4) \quad H(c) \notin B(H(x), 2\lambda r).$$

Since $|H(x) - H(c)| \leq A|x - c|^q \leq R(H, c)/3$, we have $B(H(x), 2\lambda r) \subset B(H(c), R(H, c))$. So, (2.4) implies the existence of a holomorphic inverse branch $H_x^{-1} : B(H(x), 2\lambda r) \rightarrow \mathbb{C}$ of H which sends $H(x)$ to x . Since, by the assumptions, the measure m_2 is lower t -estimable at $H(x)$ with lower estimability constant $\lambda^{-t}L$ and lower estimability radius λr , it follows from the proof of Lemma 2.8 that the measure m_1 is lower t -estimable at x with lower estimability constant $K^{-2t}\lambda^{-t}L$ and lower estimability radius $K\lambda r |H'(x)|^{-1}$. Thus, using (2.3), we get

$$\begin{aligned} m_2(B(x, \tilde{\lambda}(A^{-1}r)^{1/q})) &\geq m_2(B(x, K^{-1}\lambda r |H'(x)|^{-1})) \\ &\geq K^{-2t}\lambda^{-t}L(K\lambda r |H'(x)|^{-1})^t \\ &\geq K^{-t}Lr^t A^{-t}|x - c|^{(1-q)t} \\ &\geq K^{-t}L(A^{-1}r)^t (A^{-1}r)^{\frac{1-q}{q}t} \\ &= K^{-t}L((A^{-1}r)^{1/q})^t. \end{aligned}$$

In view of this and (2.1) the proof is completed. \square

By writing $A \preceq B$ we mean that there exists a positive constant C such that $A \leq CB$ for all A and B under consideration. Then $A \succeq B$ means that $B \preceq A$, and $A \asymp B$ says that $A \preceq B$ and $B \preceq A$.

2.2. Open Set Condition and Essential Families. In this section, starting with the open set condition, we develop the machinery of essential families of inverse branches. We first prove the following two lemmas.

Lemma 2.12. *Let $G = \langle f_1, \dots, f_u \rangle$ be a rational semigroup satisfying the nice open set condition with U . Let $j \in \{1, \dots, u\}$ and let $c \in f_j^{-1}(\bar{U})$ be a critical point of f_j . Then there exist constants $\zeta_{j,c} > 0$, $\xi_{j,c} > 0$, and $T_{j,c} > 0$ such that for each $x \in B_s(c, \zeta_{j,c}) \cap f_j^{-1}(\bar{U})$ and for each $0 < r < T_{j,c}$, $l_2(f_j^{-1}(U) \cap B_s(x, r)) \geq \xi_{j,c} r^2$.*

Proof. By conjugating G by an element of $\text{Aut}(\hat{\mathbb{C}})$, we may assume that $\infty \notin f_j^{-1}(\{f_j(c)\})$. Let W be an open neighborhood of $f_j(c)$ in \mathbb{C} such that $f_j^{-1}(W) \subset \mathbb{C}$. Let $m_1 := l_{2,e}|_{f_j^{-1}(U \cap W)}$ and $m_2 := l_{2,e}|_{U \cap W}$, where $l_{2,e}$ denotes the Euclidian measure on \mathbb{C} . Then (m_1, m_2) is a 2-conformal pair for f_j . By the nice open set condition, there exist constants $C > 0$ and $0 < R < \infty$ such that for each $y \in \bar{U} \cap W$ and for each $0 < r < R$, $m_2(B(y, r)) \geq Cr^2$. By using the method of the proof of Lemma 2.11, it is easy to see that there exist constants $\zeta'_{j,c} > 0$, $\xi'_{j,c} > 0$ and $T'_{j,c} > 0$ such that for each $x \in B(c, \zeta'_{j,c}) \cap f_j^{-1}(\bar{U})$ and for each $0 < r < T'_{j,c}$, $m_2(B(x, r)) \geq \xi'_{j,c} r^2$. Thus, the statement of our lemma holds. We are done. \square

Combining Lemma 2.12 and Koebe's Distortion Theorem, we immediately obtain the following lemma.

Lemma 2.13. *Let $G = \langle f_1, \dots, f_u \rangle$ be a rational semigroup satisfying the nice open set condition with U . Then, there exist constants $\xi > 0$ and $T > 0$ such that for each $j = 1, \dots, u$ and for each $x \in f_j^{-1}(\bar{U})$, $l_2(f_j^{-1}(U) \cap B_s(x, r)) \geq \xi r^2$.*

Let Σ_u^* be the family of finite words over the alphabet $\{1, 2, \dots, u\}$. For every $\tau \in \Sigma_u^*$, we denote by $|\tau|$ the n such that $\tau \in \{1, \dots, u\}^n$. For every $\tau \in \Sigma_u$ we set $|\tau| = \infty$. Moreover, for every $\tau = (\tau_1, \tau_2, \dots) \in \Sigma_u^* \cup \Sigma_u$ and $n \in \mathbb{N}$ with $n \leq |\tau|$, we set $\tau|_n := (\tau_1, \tau_2, \dots, \tau_n) \in \Sigma_u^*$. For every $\tau \in \Sigma_u^*$, we set $\hat{\tau} = \tau|_{|\tau|-1}$, $\tau_* := \tau|_{|\tau|}$, and $[\tau] := \{\omega \in \Sigma_u \mid \omega|_{|\tau|} = \tau\}$. Furthermore, for every $\omega \in \Sigma_u^* \cup \Sigma_u$ and $a, b \in \mathbb{N}$ with $a < b \leq |\omega|$, we set $\omega_a^b := (\omega_a, \dots, \omega_b) \in \Sigma_u^*$. For every $\omega, \tau \in \Sigma_u^*$, we say that ω and τ are comparable if either (1) $|\tau| \leq |\omega|$ and $\omega \in [\tau]$, or (2) $|\omega| \leq |\tau|$ and $\tau \in [\omega]$. We say that ω, τ are incomparable if they are not comparable.

For every family $\mathcal{F} \subset \Sigma_u^*$ let

$$\hat{\mathcal{F}} = \{\hat{\tau} : \tau \in \mathcal{F}\} \quad \text{and} \quad \mathcal{F}_* = \{\tau_* : \tau \in \mathcal{F}\}.$$

Definition 2.14. *Let $G = \langle f_1, \dots, f_u \rangle$ be a rational semigroup satisfying the nice open set condition. Suppose that $J(G) \subset \mathbb{C}$. Fix a number $M > 0$, a number $a > 0$, and V , an open subset of Σ_u . Suppose $x \in J(G)$ and $r \in (0, 1]$. A family $\mathcal{F} \subset \Sigma_u^*$ is called (M, a, V) -essential for the pair (x, r) provided that the following conditions are satisfied.*

(ess0) *For every $\tau \in \mathcal{F}$, $f_\tau(x) \in J(G)$.*

(ess1) For every $\tau \in \mathcal{F}$ there exists a number R_τ with $0 < R_\tau < a$ and an $f_{\hat{\tau},x}^{-1} : B(f_{\hat{\tau}}(x), 2R_\tau) \rightarrow \mathbb{C}$, an analytic inverse branch of $f_{\hat{\tau}}^{-1}$ sending $f_{\hat{\tau}}(x)$ to x , such that

$$M^{-1}R_\tau \leq |f'_{\hat{\tau}}(x)|r \leq \frac{1}{4}R_\tau.$$

(ess2) The family \mathcal{F} consists of mutually incomparable words.

(ess3) $\bigcup_{\tau \in \mathcal{F}} [\tau] = V$.

If $V = \Sigma_u$, the family \mathcal{F} is simply called (M, a) -essential for the pair (x, r) .

We shall prove the following.

Proposition 2.15. *Let $G = \langle f_1, \dots, f_u \rangle$ be a rational semigroup satisfying the nice open set condition with U . Suppose that $J(G) \subset \mathbb{C}$. Then, for every number $M > 0$ and for every $a > 0$ there exists an integer $\#_{(M,a)} \geq 1$ with the following properties. If V is an open subset of Σ_u , $x \in J(G)$, $r \in (0, 1]$, and $\mathcal{F} \subset \Sigma_u^*$ is an (M, a, V) -essential family for (x, r) , then we have the following.*

(a)

$$B(x, r) \subset f_{\hat{\tau},x}^{-1}(B(f_{\hat{\tau}}(x), R_\tau)) \subset \bigcup_{\gamma \in \mathcal{F}} f_{\hat{\gamma},x}^{-1}(B(f_{\hat{\gamma}}(x), R_\gamma)) \subset B(x, KMr)$$

for all $\tau \in \mathcal{F}$.

(b)

$$J(\tilde{f}) \cap (V \times B(x, r)) \subset \bigcup_{\tau \in \mathcal{F}} \tilde{f}_{\hat{\tau},x}^{-|\hat{\tau}|} (p_2^{-1}(B(f_{\hat{\tau}}(x), R_\tau))) = \bigcup_{\tau \in \mathcal{F}} [\tau] \times f_{\hat{\tau},x}^{-1}(B(f_{\hat{\tau}}(x), R_\tau)).$$

(c) $\#\mathcal{F} \leq \#_{(M,a)}$.

Proof. Item (a) follows immediately from Theorem 2.1 ($\frac{1}{4}$ -Koebe's Distortion Theorem), and Theorem 2.2. The equality part in item (b) is obvious. In order to prove the inclusion take $(\omega, z) \in J(\tilde{f}) \cap (V \times B(x, r))$. By item (ess3) of Definition 2.14 there exists $\tau \in \mathcal{F}$ such that $\omega \in [\tau]$. But then, by the first in item (a), $(\omega, z) \in [\tau] \times f_{\hat{\tau},x}^{-1}(B(f_{\hat{\tau}}(x), R_\tau))$ and item (b) is entirely proved. Let us deal with item (c). By item (osc2) of Definition 1.4,

$$\{f_{\hat{\tau},x}^{-1}((f_{\tau^*}|_{B(f_{\hat{\tau}}(x), R_\tau)})^{-1}(U))\}_{\tau \in \mathcal{F}}$$

is a family of mutually disjoint sets. Hence, using also (a), we get

$$(2.5) \quad \sum_{\tau \in \mathcal{F}} l_2(f_{\hat{\tau},x}^{-1}((f_{\tau^*}|_{B(f_{\hat{\tau}}(x), R_\tau)})^{-1}(U))) \leq l_2(B(x, KMr)) = C\pi(KM)^2r^2,$$

where $C > 0$ is a constant independent of r, M , and a . Let $L_a := \xi \min\{(T/a)^2, 1\}$, where ξ and T come from Lemma 2.13. By Lemma 2.13, we obtain that for each $j = 1, \dots, u$, for each $y \in \overline{f_j^{-1}(U)}$, and for each $0 < b \leq a$,

$$(2.6) \quad l_2(B(y, b) \cap f_j^{-1}(U)) \geq L_a b^2.$$

It follows from Theorem 2.2, (2.6), and (ess1) that for all $\tau \in \mathcal{F}$, we have

$$\begin{aligned} l_2(f_{\hat{\tau},x}^{-1}((f_{\tau_*}|_{B(f_{\hat{\tau}}(x),R_\tau)})^{-1}(U))) &\geq K^{-2}|f'_{\hat{\tau}}(x)|^{-2}l_2((f_{\tau_*}|_{B(f_{\hat{\tau}}(x),R_\tau)})^{-1}(U)) \\ &\geq K^{-2}|f'_\tau(x)|^{-2}l_2(B(f_{\hat{\tau}}(x),R_\tau) \cap f_{\tau_*}^{-1}(U)) \\ &\geq K^{-2}|f_{\hat{\tau}}(x)|^{-2}L_a R_\tau^2 \\ &\geq K^{-2}16L_a r^2. \end{aligned}$$

Combining this with (2.5) we get that $\#\mathcal{F} \leq (16L_a)^{-1}CK^4\pi M^2$. We are done. \square

3. BASIC PROPERTIES OF SEMI-HYPERBOLIC RATIONAL SEMIGROUPS

In this section define semi-hyperbolic rational semigroups and we collect and prove their dynamical properties which will be needed in the sequel.

Definition 3.1. *A rational semigroup G is called semi-hyperbolic if and only if there exists an $N \in \mathbb{N}$ and a $\delta > 0$ such that for each $x \in J(G)$ and $g \in G$,*

$$\deg(g : V \rightarrow B_s(x, \delta)) \leq N$$

for each connected component V of $g^{-1}(B_s(x, \delta))$.

The crucial tool, which makes all further considerations possible, is given by the following semigroup version of Mane's Theorem proved in [37].

Theorem 3.2. *Let $G = \langle f_1, \dots, f_u \rangle$ be a finitely generated rational semigroup. Assume that there exists an element of G with the degree at least two, that each element of $\text{Aut}(\hat{\mathbb{C}}) \cap G$ (if this is not empty) is loxodromic, and that $F(G) \neq \emptyset$. Then, G is semi-hyperbolic if and only if all of the following conditions are satisfied.*

- (a) *For each $z \in J(G)$ there exists a neighborhood U of z in $\hat{\mathbb{C}}$ such that for any sequence $\{g_n\}_{n=1}^\infty$ in G , any domain V in $\hat{\mathbb{C}}$ and any point $\zeta \in U$, the sequence $\{g_n\}_{n=1}^\infty$ does not converge to ζ locally uniformly on V .*
- (b) *For each $j = 1, \dots, u$, each $c \in \text{Crit}(f_j) \cap J(G)$ satisfies $\text{dist}(c, G^*(f_j(c))) > 0$.*

The first author proved in [37] the following.

Theorem 3.3. *Let $G = \langle f_1, \dots, f_u \rangle$ be a semi-hyperbolic finitely generated rational semigroup. Assume that there exists an element of G with the degree at least two, that each element of $\text{Aut}(\hat{\mathbb{C}}) \cap G$ (if this is not empty) is loxodromic, and that $F(G) \neq \emptyset$. Then there exist $R > 0$, $C > 0$, and $\alpha > 0$ such that if $x \in J(G)$, $\omega \in \Sigma_u^*$ and V is a connected component of $f_\omega^{-1}(B_s(x, R))$, then V is simply connected and $\text{diam}_s(V) \leq Ce^{-\alpha|\omega|}$.*

Throughout the rest of the paper, we assume the following:

Assumption (*):

- Let $f = (f_1, \dots, f_u) \in (\text{Rat})^u$ be a multi-map and let $G = \langle f_1, \dots, f_u \rangle$.
- There exists an element g of G such that $\deg(g) \geq 2$.

- Each element of $\text{Aut}(\hat{\mathbb{C}}) \cap G$ (if this is not empty) is loxodromic.
- G is semi-hyperbolic.
- G satisfies the nice open set condition.

In order to prove the main results (Theorem 1.11 etc.), in virtue of [47] and [48], we may assume that $u \geq 2$. If $u \geq 2$, then the open set condition implies that $F(G) \neq \emptyset$. Hence, conjugating G by some element of $\text{Aut}(\hat{\mathbb{C}})$ if necessary, we may assume that $J(G) \subset \mathbb{C}$. Thus, throughout the rest of the paper, in addition to the above assumption, we also assume that

- $u \geq 2$ and $J(G) \subset \mathbb{C}$.

Note that in Theorem 1.11, we work with the spherical distance. However, throughout the rest of the paper, we will work with the Euclidian distance. If we want to get the results on the spherical distance (and this would include the case $u = 1$), then we have only to consider some minor modifications in our argument.

We now give further notation. A pair $(c, j) \in \hat{\mathbb{C}} \times \{1, 2, \dots, u\}$ is called critical if $f'_i(c) = 0$. The set of all critical pairs of f will be denoted by $\text{CP}(f)$. Let $\text{Crit}(f)$ be the union of $\bigcup_{j=1}^u \text{Crit}(f_j)$. For every $c \in \text{Crit}(f)$ put

$$c_+ = \{f_j(c) : (c, j) \in \text{CP}(f)\}.$$

The set c_+ is called the set of critical values of c . For any subset A of $\text{Crit}(f)$ put

$$A_+ = \{c_+ : c \in A\}.$$

For each $(c, j) \in \text{CP}(f)$ let $q(c, j)$ be the local order of f_j at c . For any set $F \subset \hat{\mathbb{C}}$, set

$$\omega_G(F) = \bigcap_{n=0}^{\infty} \overline{\bigcup_{|\omega| \geq n} \phi_{\omega}(F)}.$$

The latter is called the ω -limit set of F with respect to the semigroup G . Similarly, for every set $B \subset \Sigma_u \times \hat{\mathbb{C}}$,

$$\omega(B) = \bigcap_{N=0}^{\infty} \overline{\bigcup_{n \geq N} \tilde{f}^n(B)},$$

and this set is called the ω -limit set of F with respect to the skew product map $\tilde{f} : \Sigma_u \times \hat{\mathbb{C}} \rightarrow \Sigma_u \times \hat{\mathbb{C}}$.

Given $\omega \in \Sigma_u^*$, $j \in \{1, 2, \dots, u\}$, $z \in \tilde{f}_{\omega}^{-1}(J(\tilde{f}))$ and $r > 0$, we say that a critical pair (c, j) sticks to $\text{Comp}(z, f_{\omega}, r)$ if $c \in \text{Comp}(z, f_{\omega}, r)$ and $j = \omega_1$. We then write

$$(c, j) \sim \text{Comp}(z, f_{\omega}, r).$$

Set

$$A = A_f := \max\{A(f_j, c) : (c, j) \in \text{CP}(f)\} \quad \text{and} \quad R_f := \min\{R(f_j, c) : (c, j) \in \text{CP}(f)\}.$$

For A, B , any two subsets of a metric space put

$$\text{dist}(A, B) = \inf\{\text{dist}(a, b) : a \in A, b \in B\}$$

and

$$\text{Dist}(A, B) = \sup\{\text{dist}(a, b) : a \in A, b \in B\}.$$

Fix a positive β smaller than the following four positive numbers (a)–(d).

(a) $\min\{\text{dist}(c, G^*(c_+)) : c \in \text{Crit}(f) \cap J(G)\}$, (b) R_f , (c) $\min\{|c' - c| : c, c' \in \text{Crit}(f) \cap J(G), c \neq c'\}$,

and

$$(d) \text{dist}\left(\bigcup_{j=1}^u CV(f_j) \cap F(G), J(G)\right),$$

where, (a) is positive because of semi-hyperbolicity (Theorem 3.2). It immediately follows from Theorem 3.3 that there exists $\gamma \in (0, 1/4)$ so small that if $g \in G^*$ and $g(x) \in J(G)$, then

$$(3.1) \quad \text{Comp}(x, g, 2\gamma) \subset \mathbb{C} \quad \text{and} \quad \text{diam}(\text{Comp}(x, g, 2\gamma)) < \beta.$$

We shall prove the following.

Lemma 3.4. *Fix $\eta \in (0, \beta)$, an integer $n \geq 0$ and $(\omega, z) \in J(\tilde{f})$. Suppose that for every $0 \leq k \leq n - 1$,*

$$\text{diam}(\text{Comp}(f_{\omega|_k}(z), f_{\omega|_{k+1}}^n, \eta)) \leq \beta.$$

Then each connected component $\text{Comp}(f_{\omega|_k}(z), f_{\omega|_{k+1}}^n, \eta)$ is sticked to by at most one critical pair (c, j) of f ; and if a critical pair (c, j) sticks to a component $\text{Comp}(f_{\omega|_k}(z), f_{\omega|_{k+1}}^n, \eta)$, then $f_j(c) \in J(G)$. Furthermore, each critical pair of f sticks to at most one of all these components $\text{Comp}(f_{\omega|_k}(z), f_{\omega|_{k+1}}^n, \eta)$.

Proof. The first part is obvious by the choice of β . In order to prove the second part suppose that

$$(c, \omega_{k+1}) \sim \text{Comp}(f_{\omega|_k}(z), f_{\omega|_{k+1}}^n, \eta) \quad \text{and} \quad (c, \omega_{l+1}) \sim \text{Comp}(f_{\omega|_l}(z), f_{\omega|_{l+1}}^n, \eta)$$

with some $0 \leq k < l \leq n - 1$ and $\omega_{k+1} = \omega_{l+1}$. Then both c and $f_{\omega|_{k+1}}^l(c)$ belong to $\text{Comp}(f_{\omega|_l}(z), f_{\omega|_{l+1}}^n, \eta)$, and therefore,

$$|f_{\omega|_{k+1}}^l(c) - c| \leq \text{diam}(\text{Comp}(f_{\omega|_l}(z), f_{\omega|_{l+1}}^n, \eta)) \leq \beta,$$

contrary to the choice of β . □

Let

$$\kappa = \prod_{(c,j) \in \text{CP}(f)} q(c, j)^{-1}.$$

Lemma 3.5. *If $g \in G$ and $z \in g^{-1}(J(G))$, then*

$$\text{Mod}(\text{Comp}(z, g, 2\gamma) \setminus \overline{\text{Comp}(z, g, \gamma)}) \geq \frac{\kappa \log 2}{\#\text{CP}(f)}.$$

Proof. By Lemma 3.4 there exists a geometric annulus $R \subset B(g(z), 2\gamma) \setminus B(g(z), \gamma)$ centered at $g(z)$ and with modulus $\geq \log 2 / \#\text{CP}(f)$ and such that $g^{-1}(R) \cap \text{Comp}(z, g, 2\gamma) \cap \text{Crit}(f) = \emptyset$. Since covering maps increase moduli of annuli by factors at most equal to their degrees, we conclude that

$$\text{Mod}(\text{Comp}(z, g, 2\gamma) \setminus \overline{\text{Comp}(z, g, \gamma)}) \geq \text{Mod}(R_g) \geq \frac{\log 2 / \#\text{CP}(f)}{\prod_{(c,j) \in \text{CP}(f)} q(c, j)} = \frac{\kappa \log 2}{\#\text{CP}(f)},$$

where $R_g \subset \text{Comp}(z, g, 2\gamma)$ is the connected component of R enclosing $\text{Comp}(z, g, \gamma)$. \square

As an immediate consequence of this lemma and Theorem 2.4 we get the following.

Lemma 3.6. *Let $\omega \in \Sigma_u^*$ and suppose $f_\omega(z) \in J(G)$. If $0 \leq k \leq |\omega|$ and the map $f_{\omega|_k} : \text{Comp}(z, f_\omega, 2\gamma) \rightarrow \text{Comp}(f_{\omega|_k}(z), f_{\omega|_{k+1}}^{\omega|_k}, 2\gamma)$ is univalent, then*

$$\frac{|f'_{\omega|_k}(y)|}{|f'_{\omega|_k}(x)|} \leq \text{const}$$

for all $x, y \in \text{Comp}(z, f_\omega, \gamma)$, where *const* is a number depending only on $\#\text{CP}(f)$ and κ .

Lemma 3.7. *Suppose that $g \in G$ and $g(z) \in J(\tilde{f})$. Suppose also that $Q^{(1)} \subset Q^{(2)} \subset B(g(z), \gamma)$ are connected sets. If $Q_g^{(2)}$ is a connected component of $g^{-1}(Q^{(2)})$ contained in $\text{Comp}(z, g, \gamma)$ and $Q_g^{(1)}$ is a connected component of $g^{-1}(Q^{(1)})$ contained in $Q_g^{(2)}$, then*

$$\frac{\text{diam}(Q_g^{(1)})}{\text{diam}(Q_g^{(2)})} \geq \Gamma \frac{\text{diam}(Q^{(1)})}{\text{diam}(Q^{(2)})}.$$

with some universal constant $\Gamma > 0$.

Proof. Write $g = f_\omega$, where $\omega \in \Sigma_u^*$ and put $n = |\omega|$. For every $0 \leq j \leq n$, set

$$Q_j^{(1)} = f_{\omega|_{n-j}}(Q_g^{(1)}) \quad \text{and} \quad Q_j^{(2)} = f_{\omega|_{n-j}}(Q_g^{(2)}).$$

Let $1 \leq n_1 \leq \dots \leq n_v \leq n$ be all the integers k between 1 and n such that

$$\text{Crit}(f_{\omega_{n-k+1}}) \cap \text{Comp}(f_{\omega|_{n-k}}(z), f_{\omega_{n-k+1}}^{\omega_{n-k}}, 2\gamma) \neq \emptyset.$$

Fix $1 \leq i \leq v$. If $j \in [n_i, n_{i+1} - 1]$ (we set $n_{v+1} = n - 1$), then by Lemma 3.6 there exists a universal constant $T > 0$ such that

$$(3.2) \quad \frac{\text{diam}(Q_j^{(1)})}{\text{diam}(Q_j^{(2)})} \geq T \frac{\text{diam}(Q_{n_i}^{(1)})}{\text{diam}(Q_{n_i}^{(2)})}.$$

Since, in view of Lemma 3.4, $v \leq \#\text{CP}(f)$, in order to conclude the proof it is enough to show the existence of a universal constant $E > 0$ such that for every $1 \leq i \leq v$

$$\frac{\text{diam}(Q_{n_i}^{(1)})}{\text{diam}(Q_{n_i}^{(2)})} \geq E \frac{\text{diam}(Q_{n_{i-1}}^{(1)})}{\text{diam}(Q_{n_{i-1}}^{(2)})}.$$

Indeed, let c be the critical point in $\text{Comp}(f_{\omega|_{n-n_i}}(z), f_{\omega|_{n_i+1}}, 2\gamma)$ and let q be its order. Since both sets $Q_{n_i}^{(1)}$ and $Q_{n_i}^{(2)}$ are connected, we get for $j = 1, 2$ that

$$\begin{aligned} \text{diam}(Q_{n_i-1}^{(j)}) &\asymp \text{diam}(Q_{n_i}^{(j)}) \sup\{|f'_{\omega_{n-n_i+1}}(x)| : x \in Q_{n_i}^{(j)}\} \\ &\asymp \text{diam}(Q_{n_i}^{(j)}) \text{Dist}(c, Q_{n_i}^{(j)}). \end{aligned}$$

Hence

$$\begin{aligned} \frac{\text{diam}(Q_{n_i}^{(1)})}{\text{diam}(Q_{n_i}^{(2)})} &\asymp \frac{\text{diam}(Q_{n_i-1}^{(1)})}{\text{Dist}(c, Q_{n_i}^{(1)})} \cdot \frac{\text{Dist}(c, Q_{n_i}^{(2)})}{\text{diam}(Q_{n_i-1}^{(2)})} \\ &\geq \frac{\text{diam}(Q_{n_i-1}^{(1)})}{\text{diam}(Q_{n_i-1}^{(2)})}. \end{aligned}$$

The proof is finished. \square

4. PARTIAL ORDER IN $\text{Crit}(f) \cap J(G)$ AND STRATIFICATION OF $J(G)$

In this section we introduce a partial order in the critical set $\text{Crit}(f) \cap J(G)$ and stratification of $J(G)$. They will be used to do the inductive steps in the proofs of the main theorems of our paper. We start with the following.

Lemma 4.1. *The set $\omega_G((\text{Crit}(f) \cap J(G))_+)$ is nowhere dense in $J(G)$.*

Proof. Suppose on the contrary that the interior (relative to $J(G)$) of $\omega_G((\text{Crit}(f) \cap J(G))_+)$ is not empty. Then, there exists a critical point $c \in \text{Crit}(f) \cap J(G)$ such that $\omega_G(c_+)$ has non-empty interior. But then, in virtue of Proposition 1.2 there would exist finitely many elements $g_1, g_2, \dots, g_n \in G$ such that $\omega_G(c_+) \supset g_1(\omega_G(c_+)) \cup g_2(\omega_G(c_+)) \cup \dots \cup g_n(\omega_G(c_+)) \subset J(G)$. Hence $c \in \omega_G(c_+)$, contrary to the non-recurrence condition (Theorem 3.2). \square

Now we introduce in $\text{Crit}(f) \cap J(G)$ a relation $<$ which, in view of Lemma 4.2 below, is an ordering relation. Put

$$c_1 < c_2 \Leftrightarrow c_1 \in \omega_G(c_{2+}).$$

Lemma 4.2. *If $c_1 < c_2$ and $c_2 < c_3$, then $c_1 < c_3$.*

Proof. Since $c_2 \in \omega_G(c_{3+})$, we have $\omega_G(c_{2+}) \subset \omega_G(c_{3+})$. Along with $c_1 \in \omega_G(c_{2+})$ this implies that $c_1 \in \omega_G(c_{3+})$, meaning that $c_1 < c_3$. \square

Lemma 4.3. *There exists no $c \in \text{Crit}(f) \cap J(G)$ such that $c < c$.*

Proof. Indeed, $c < c$ means that $c \in \omega_G(c_+)$, contrary to the non-recurrence condition. \square

Since the set $\text{Crit}(f) \cap J(G)$ is finite, as an immediate of this lemma and Lemma 4.2 we get the following.

Lemma 4.4. *There is no infinite linear subset of the partially ordered set $(\text{Crit}(f) \cap J(G), <)$.*

Now define inductively a sequence $\{Cr_i(f)\}$ of subsets of $\text{Crit}(f) \cap J(G)$ by setting $Cr_0(f) = \emptyset$ and

$$(4.1) \quad Cr_{i+1}(f) = \left\{ c \in (\text{Crit}(f) \cap J(G)) \setminus \bigcup_{j=0}^i Cr_j(f) : c' < c, \Rightarrow c' \in \bigcup_{j=0}^i Cr_j(f) \right\}.$$

Lemma 4.5. *The following four statements hold.*

- (a) *The sets $\{Cr_i(f)\}$ are mutually disjoint.*
- (b) $\exists_{p \geq 1} \forall_{i \geq p+1} Cr_i(f) = \emptyset$.
- (c) $Cr_0(f) \cup \dots \cup Cr_p(f) = \text{Crit}(f) \cap J(G)$.
- (d) $Cr_1(f) \neq \emptyset$.

Proof. By definition $Cr_{i+1}(f) \cap \bigcup_{j=0}^i Cr_j(f) = \emptyset$, whence disjointness in (a) is clear. As the set $\text{Crit}(f) \cap J(G)$ is finite, (b) follows from (a). Take p to be the minimal number satisfying (b) and suppose that $(\text{Crit}(f) \cap J(G)) \setminus \bigcup_{j=0}^p Cr_j(f) \neq \emptyset$. Take $c \in (\text{Crit}(f) \cap J(G)) \setminus \bigcup_{j=0}^p Cr_j(f)$. Since $Cr_{p+1} = \emptyset$, there would thus exist $c' \in (\text{Crit}(f) \cap J(G)) \setminus \bigcup_{j=0}^p Cr_j(f)$ such that $c' < c$. Iterating this procedure we would obtain an infinite sequence $c_1 = c > c' = c_2 > c_3 > \dots$, contrary to Lemma 4.4. Now, part (d) follows from (c) and (4.1). \square

For every $(\tau, z) \in J(\tilde{f})$ put

$$\text{Crit}(\tau, z) = \text{Crit}(\tilde{f}) \cap \omega(\tau, z) \quad \text{and} \quad \text{Crit}(\tau, z)_+ = p_2(\text{Crit}(\tilde{f}) \cap \omega(\tau, z))_+.$$

Lemma 4.6. *If $(\tau, z) \in J(\tilde{f})$, then*

$$p_2(\omega(\tau, z)) \not\subset \overline{G^*(\text{Crit}(\tau, z)_+)}.$$

Proof. Suppose on the contrary that

$$(4.2) \quad p_2(\omega(\tau, z)) \subset \overline{G^*(\text{Crit}(\tau, z)_+)}.$$

Consequently, $\text{Crit}(\tau, z) \neq \emptyset$. Let $(\tau^1, c_1) \in \text{Crit}(\tau, z)$. This means that $(\tau^1, c_1) \in \omega(\tau, z)$, and it follows from (4.2) that there exists $(\tau^2, c_2) \in \text{Crit}(\tau, z)$ such that either $c_1 \in \omega_G(c_{2+})$ or $c_1 = g_1(c_2)$ for some $g_1 \in G$ of the form f_ω with $f'_{\omega_1}(c_2) = 0$. Iterating this procedure we obtain an infinite sequence $((\tau^j, c_j))_{j=1}^\infty$ of points in $\text{Crit}(\tau, z)$ such that for every $j \geq 1$ either $c_j \in \omega_G(c_{j+1+})$ or $c_j = g_j(c_{j+1})$ for some $g_j \in G$ of the form f_ρ with $f'_{\rho_1}(c_{j+1}) = 0$. Consider an arbitrary block c_k, c_{k+1}, \dots, c_l such that $c_j = g_j(c_{j+1})$ for every $k \leq j \leq l-1$, and suppose that $l - (k-1) \geq \#(\text{Crit}(f) \cap J(G))$. Then there are two indices $k \leq a < b \leq l$ such that $c_a = c_b$. Hence $g_a \circ g_{a+1} \circ \dots \circ g_{b-1}(c_b) = c_a = c_b$ and $(g_a \circ g_{a+1} \circ \dots \circ g_{b-1})'(c_b) = 0$. This however contradicts our assumption that the Julia set of G contains no superstable fixed points. In consequence, the length of the block c_k, c_{k+1}, \dots, c_l is bounded above by $\#(\text{Crit}(f) \cap J(G))$. Therefore, there exists an infinite sequence $(j_n)_{n=1}^\infty$ such that $c_{j_n} \in \omega_G(c_{j_n+1+})$ for all $n \geq 1$. This however contradicts Lemma 4.4 and finishes the proof. \square

Now, for every $i = 0, 1, \dots, p$, set

$$S_i(f) = Cr_0(f) \cup Cr_1(f) \cup \dots \cup Cr_i(f).$$

Fix $i \in \{0, 1, \dots, p-1\}$ consider an arbitrary point $c' \in \bigcup_{c \in Cr_{i+1}(f)} \omega_G(c_+) \cap \text{Crit}(f) \cap J(G)$. Then there exists $c \in Cr_{i+1}(f)$ such that $c' \in \omega_G(c_+)$ which equivalently means that $c' < c$. Thus, by (4.1) we get $c' \in S_i(f)$. So,

$$(4.3) \quad \bigcup_{c \in Cr_{i+1}(f)} \omega_G(c_+) \cap ((\text{Crit}(f) \cap J(G)) \setminus S_i(f)) = \emptyset.$$

Since the set $\bigcup_{c \in Cr_{i+1}(f)} \omega_G(c_+)$ is compact and $(\text{Crit}(f) \cap J(G)) \setminus S_i(f)$ is finite, we therefore get

$$(4.4) \quad \delta_i = \text{dist} \left(\bigcup_{c \in Cr_{i+1}(f)} \omega_G(c_+), (\text{Crit}(f) \cap J(G)) \setminus S_i(f) \right) > 0.$$

Set

$$\rho = \min\{\delta_i/2 : i = 0, 1, \dots, p-1\}$$

and for every $i = 0, 1, \dots, p$ define

$$(4.5) \quad J_i(G) = \{z \in J(G) : \text{dist}(G^*(z), (\text{Crit}(f) \cap J(G)) \setminus S_i(f)) \geq \rho\}.$$

We end this section with the following two lemmas concerning the sets $J_i(G)$.

Lemma 4.7. $\emptyset \neq J_0(G) \subset J_1(G) \subset \dots \subset J_p(G) = J(G)$.

Proof. Since $S_{i+1}(f) \supset S_i(f)$, the inclusions $J_i(G) \subset J_{i+1}(G)$ are obvious. Since $S_p(f) = \text{Crit}(f) \cap J(G)$ (see Lemma 4.5), it holds $J_p(G) = J(G)$. We get from (4.4) that

$$\text{dist} \left(\bigcup_{c \in Cr_1(f)} \omega_G(c_+), (\text{Crit}(f) \cap J(G)) \setminus S_0(f) \right) = \delta_0 \geq 2\rho \geq \rho.$$

Thus, $\bigcup_{c \in Cr_1(f)} \omega_G(c_+) \subset J_0(G)$, and since $Cr_1(f) \neq \emptyset$ (see Lemma 4.5), we conclude that $J_0(G) \neq \emptyset$. The proof is complete. \square

Lemma 4.8. *There exists $l = l(f) \geq 0$ so large that for all $i = 0, 1, \dots, p-1$ we have*

$$\bigcup_{c \in Cr_{i+1}(f)} \omega_G(c_+) \cap J(G) \subset \overline{\bigcup_{|\tau| \geq l} f_\tau(Cr_{i+1}(f)_+) \cap J(G)} = \overline{G^* \left(\bigcup_{|\tau| = l} f_\tau(Cr_{i+1}(f)_+) \right)} \cap J(G) \subset J_i(G).$$

Proof. The left-hand inclusion is obvious regardless of what $l(f)$ is. The equal-ity part of the assertion is obvious. In order to prove the right-hand inclusion fix $i \in \{0, 1, \dots, p-1\}$. By the definition of the ω -limit sets of G there exists $l_i \geq 0$ such that for every $c \in Cr_{i+1}(f)$ we have $\text{dist}(\bigcup_{|\tau| \geq l_i} f_\tau(c_+), \bigcup_{c' \in Cr_{i+1}(f)} \omega_G(c'_+)) < \delta_i/2$. Thus, by (4.4), $\text{dist}(\bigcup_{|\tau| \geq l_i} f_\tau(c_+), (\text{Crit}(f) \cap J(G)) \setminus S_i(f)) > \delta_i/2 \geq \rho$. Hence, for every $\tau \in \Sigma_u^*$ with $|\tau| \geq l_i$, we have $\text{dist}(f_\tau(c_+), (\text{Crit}(f) \cap J(G)) \setminus S_i(f)) > \rho$. Thus $f_\tau(c_+) \subset J_i(G)$. Therefore, $\bigcup_{|\tau| \geq l_i} f_\tau(c_+) \subset J_i(G)$, and consequently, $\bigcup_{|\tau| \geq l_i} f_\tau(Cr_{i+1}(f)_+) \subset J_i(G)$. Since

$J_i(G)$ is a closed set, this yields that $\overline{\bigcup_{|\tau| \geq l_i} f_\tau(Cr_{i+1}(f)_+)} \subset J_i(G)$. Setting $l(f) = \max\{l_i : i = 0, 1, \dots, p-1\}$ completes the proof. \square

5. HOLOMORPHIC INVERSE BRANCHES.

In this section we prove the existence of suitable holomorphic inverse branches, our basic tools throughout the paper. Set

$$\text{Sing}(\tilde{f}) = \bigcup_{n \geq 0} \tilde{f}^{-n}(\text{Crit}(\tilde{f}))$$

and

$$\text{Sing}(f) = \bigcup_{g \in G^*} g^{-1}(\text{Crit}(f)).$$

Proposition 5.1. *For each $(\tau, z) \in J(\tilde{f}) \setminus \text{Sing}(\tilde{f})$, there exists a number $\eta(\tau, z)$ with $0 < \eta(\tau, z) < \gamma$, an increasing sequence $(n_j)_{j=1}^\infty$ of positive integers and a point $(\hat{\tau}, \hat{z}) \in \omega(\tau, z) \setminus p_2^{-1}(\overline{G^*(\text{Crit}(\tau, z)_+)})$ with the following two properties.*

- (a) $\lim_{j \rightarrow \infty} \tilde{f}^{n_j}(\tau, z) = (\hat{\tau}, \hat{z})$,
- (b) $\text{Comp}(z, f_{\tau|n_j}, \eta(\tau, z)) \cap \text{Crit}(f_{\tau|n_j}) = \emptyset$

Proof. In view of Lemma 4.6 there exists a point $(\hat{\tau}, \hat{z}) \in \omega(\tau, z)$ such that $\hat{z} \notin \overline{G^*(\text{Crit}(\tau, z)_+)}$. Let

$$\eta = \frac{1}{2} \text{dist}(\hat{z}, \overline{G^*(\text{Crit}(\tau, z)_+)}).$$

Then there exists an infinite increasing sequence $(n_j)_{j=1}^\infty$ of positive integers such that

$$(5.1) \quad \lim_{j \rightarrow \infty} \tilde{f}^{n_j}(\tau, z) = (\hat{\tau}, \hat{z})$$

and

$$(5.2) \quad f_{\tau|n_j}(z) \notin B(\overline{G^*(\text{Crit}(\tau, z)_+)}, \eta)$$

for all $j \geq 1$. We claim that there exists $\eta(\tau, z) > 0$ such that for all $j \geq 1$ large enough

$$\text{Comp}(z, f_{\tau|n_j}, \eta(\tau, z)) \cap \text{Crit}(f_{\tau|n_j}) = \emptyset.$$

Indeed, otherwise we find an increasing subsequence $(j_i)_{i=1}^\infty$ and a decreasing to zero sequence of positive numbers $\eta_i < \eta$ such that

$$\text{Comp}(z, f_{\tau|n_{j_i}}, \eta_i) \cap \text{Crit}(f_{\tau|n_{j_i}}) \neq \emptyset.$$

Let $\tilde{c}_i \in \text{Comp}(z, f_{\tau|n_{j_i}}, \eta_i) \cap \text{Crit}(f_{\tau|n_{j_i}})$. Then there exist $0 \leq p_i \leq n_{j_i} - 1$ and

$$(5.3) \quad c_i \in \text{Crit}(f_{\tau|p_i+1})$$

such that $c_i = f_{\tau|p_i}(\tilde{c}_i)$. Since $\lim_{i \rightarrow \infty} \eta_i = 0$, it follows from Theorem 3.3 that $\lim_{i \rightarrow \infty} \tilde{c}_i = z$. Since $(\tau, z) \notin \bigcup_{n \geq 0} \tilde{f}^{-n}(\text{Crit}(\tilde{f}))$, this implies that $\lim_{i \rightarrow \infty} p_i = +\infty$. But then, making use of Theorem 3.3 again and of the formula $(\sigma^{p_i}(\tau), c_i) = f^{p_i}(\tau, \tilde{c}_i)$, we conclude that the set

of accumulation points of the sequence $((\sigma^{p_i}(\tau), c_i))_1^\infty$ is contained in $\omega(\tau, z)$. Fix (τ^∞, c) to be one of these accumulation points. Since $\text{Crit}(f)$ is closed we conclude that

$$(5.4) \quad (\tau^\infty, c) \in \text{Crit}(\tau, z).$$

Since that set $\text{Crit}(f)$ is finite, passing to a subsequence, we may assume without loss of generality that $(c_i)_1^\infty$ is a constant sequence, so equal to c . Since $c = f_{\tau|_{p_i}}(\tilde{c}_i)$, we get

$$\left| f_{\tau|_{n_{j_i}}}(z) - f_{\tau|_{p_i+1}}^{n_{j_i}}(c) \right| = \left| f_{\tau|_{n_{j_i}}}(z) - f_{\tau|_{n_{j_i}}}(\tilde{c}_i) \right| < \eta_i < \eta.$$

But, looking at (5.3) and (5.4), we conclude that $f_{\tau|_{p_i+1}}^{n_{j_i}}(c) \in G^*(\text{Crit}(\tau, z)_+)$. We thus arrived at a contradiction with (5.2), and the proof is finished. \square

Corollary 5.2. *If $(\tau, z) \in J(\tilde{f}) \setminus \text{Sing}(\tilde{f})$, then $\limsup_{n \rightarrow \infty} |(f'_{\tau|_n}(z)| = +\infty$.*

Proof. Let $(n_j)_{j=1}^\infty$ and $\eta(\tau, z)$ be produced by Proposition 5.1. Then, by this proposition and Theorem 3.3, the family $\left\{ f_{\tau|_{n_j}, z}^{-1} : B(f_{\tau|_{n_j}}(z), \eta(\tau, z)) \rightarrow \mathbb{C} \right\}_{j=1}^\infty$ of holomorphic inverse branches of $f_{\tau|_{n_j}}$ sending $f_{\tau|_{n_j}}(z)$ to z is well defined and normal. As a matter of fact we mean here this family restricted to the disk $B(\hat{z}, \eta(\tau, z)/2)$ and $j \geq 1$ large enough. Therefore by Theorem 3.3 again, $\lim_{j \rightarrow \infty} |(f'_{\tau|_{n_j}}(z)|^{-1} = 0$ and we are done. \square

We end this section with the following. Let $\|\tilde{f}'\| = \sup_{w \in J(\tilde{f})} |\tilde{f}'(w)|$.

Proposition 5.3. *Fix $\theta \in (0, \min\{1, \gamma\})$. For all $(\tau, z) \in J(\tilde{f})$ and $r > 0$ there exists a minimal integer $s = s(\theta, (\tau, z), r) \geq 0$ with the following properties (a) and (b).*

- (a) $|(\tilde{f}^s)'(\tau, z)| \neq 0$.
- (b) *Either $r|(\tilde{f}^s)'(\tau, z)| > \|\tilde{f}'\|^{-1}$ or there exists $c \in \text{Crit}(f_{\tau_{s+1}})$ such that $f_{\tau_{s+1}}(c) \in J(G)$ and*

$$|f_{\tau|_s}(z) - c| \leq \theta r |f'_{\tau|_s}(z)|.$$

In addition, for this s , we have

- (c) $\theta r |f'_{\tau|_s}(z)| \leq \theta < \gamma$ and

$$\text{Comp}(z, f_{\tau|_s}, (KA_f^2)^{-1} 2^{-\#\text{Crit}(f)} \theta r |f'_{\tau|_s}(z)|) \cap \text{Crit}(f_{\tau|_s}) = \emptyset.$$

Proof. In view of Corollary 5.2 the set of integers (≥ 0) satisfying conditions (a) and (b) is not empty. Let s be the minimum of those numbers. Then conditions (a) and (b) are satisfied. If $s = 0$ then (c) is also satisfied since the identity map has no critical points. So, we may assume that $s \geq 1$. By the definition of s we have $r|(\tilde{f}^{s-1})'(\tau, z)| \leq \|\tilde{f}'\|^{-1}$, whence

$$\begin{aligned} \theta r |f'_{\tau|_s}(z)| &= \theta r |(\tilde{f}^s)'(\tau, z)| = \theta r |(\tilde{f}^{s-1})'(\tau, z)| \cdot |\tilde{f}'(\tilde{f}^{s-1}(\tau, z))| \\ &\leq \theta \|\tilde{f}'\|^{-1} \|\tilde{f}'\| = \theta < \gamma. \end{aligned}$$

Thus, (3.1) yield for all $0 \leq j \leq s$ that

$$\text{diam}(\text{Comp}(f_{\tau|_{s-j}}(z), f_{\tau|_{s-j+1}}^s, \theta r |f'_{\tau|_s}(z)|)) \leq \beta.$$

It therefore follows from Lemma 3.4 that there exist $0 \leq p \leq \#\text{Crit}(f)$, an increasing sequence of integers $1 \leq k_1 < k_2 < \dots < k_p \leq s$ and mutually distinct critical pairs $(c_1, \tau_{s-k_1+1}), (c_2, \tau_{s-k_2+1}), \dots, (c_p, \tau_{s-k_p+1})$ such that $f_{\tau_{s-k_l+1}}(c_l) \in J(G)$ and

$$\text{Comp}(f_{\tau|_{s-k_l}}(z), f_{\tau|_{s-k_l+1}}^s, \theta r|_{f'_{\tau|_s}(z)}) \cap \text{Crit}(f_{\tau_{s-k_l+1}}) = \{c_l\}$$

for every $l = 1, 2, \dots, p$, and, in addition, if $j \notin \{k_1, k_2, \dots, k_p\}$, then

$$(5.5) \quad \text{Comp}(f_{\tau|_{s-j}}(z), f_{\tau|_{s-j+1}}^s, \theta r|_{f'_{\tau|_s}(z)}) \cap \text{Crit}(f_{\tau_{s-j+1}}) = \emptyset.$$

Setting $k_0 = 0$, we shall prove by induction that for every $0 \leq l \leq p$, we have

$$(5.6) \quad \text{Comp}(f_{\tau|_{s-k_l}}(z), f_{\tau|_{s-k_l+1}}^s, (KA_f^2)^{-1}2^{-l}\theta r|_{f'_{\tau|_s}(z)}) \cap \text{Crit}(f_{\tau|_{s-k_l+1}}^s) = \emptyset,$$

where $f_{\tau_v}^s = Id$ if $s < v$. Indeed, for $l = 0$ there is nothing to prove. So, suppose that (5.6) is true for some $0 \leq l \leq p-1$. Then using (5.5) we get

$$\text{Comp}(f_{\tau|_{s-k_{l+1}+1}}(z), f_{\tau|_{s-k_{l+1}+2}}^s, (KA_f^2)^{-1}2^{-l}\theta r|_{f'_{\tau|_s}(z)}) \cap \text{Crit}(f_{\tau|_{s-k_{l+1}+2}}^s) = \emptyset.$$

So, if

$$c_{l+1} \in \text{Comp}(f_{\tau|_{s-k_{l+1}}}(z), f_{\tau|_{s-k_{l+1}+1}}^s, (KA_f^2)^{-1}2^{-(l+1)}\theta r|_{f'_{\tau|_s}(z)}),$$

then, by Lemma 2.6 applied to holomorphic maps $H = f_{\tau_{s-k_{l+1}+1}}$ and $Q = f_{\tau|_{s-k_{l+1}+2}}^s$, z being $f_{\tau|_{s-k_{l+1}}}(z)$ and the radius $R = (KA_f^2)^{-1}2^{-(l+1)}\theta r|_{f'_{\tau|_s}(z)} < \gamma$, we get

$$\begin{aligned} \left| f_{\tau|_{s-k_{l+1}}}(z) - c_{l+1} \right| &\leq KA_f^2 \left| f_{\tau|_{s-k_{l+1}+1}}^s(f_{\tau|_{s-k_{l+1}}}(z)) \right|^{-1} (KA_f^2)^{-1}2^{-(l+1)}\theta r|_{f'_{\tau|_s}(z)}| \\ &= 2^{-(l+1)}\theta r|_{f'_{\tau|_{s-k_{l+1}}}(z)}| \\ &\leq \theta r|_{f'_{\tau|_{s-k_{l+1}}}(z)}|, \end{aligned}$$

which along with the facts that $c_{l+1} \in \text{Crit}(f_{\tau|_{s-k_{l+1}+1}})$ and $f_{\tau_{s-k_{l+1}+1}}(c_{l+1}) \in J(G)$ contradicts the definition of s and proves (5.6) for $l+1$. In particular, it follows from (5.6) with $l = p$ and (5.5) with $j = k_p + 1, k_p + 2, \dots, s$, that

$$\text{Comp}(z, f_{\tau|_s}, (KA_f^2)^{-1}2^{-\#\text{Crit}(f)}\theta r|_{f'_{\tau|_s}(z)}) \cap \text{Crit}(f_{\tau|_s}) = \emptyset.$$

We are done. □

6. GEOMETRIC MEASURES THEORY AND CONFORMAL MEASURES; PRELIMINARIES

In this section we deal in detail with Hausdorff and packing measures and we also establish some geometrical properties of conformal measures.

6.1. Preliminaries from Geometric Measure Theory; Hausdorff and Packing Measures. Given a subset A of a metric space (X, d) , a countable family $\{B(x_i, r_i)\}_{i=1}^{\infty}$ of open balls centered at the set A is said to be a packing of A if and only if for any pair $i \neq j$

$$d(x_i, x_j) > r_i + r_j.$$

Given $t \geq 0$, the t -dimensional outer Hausdorff measure $H^t(A)$ of the set A is defined as

$$H^t(A) = \sup_{\varepsilon > 0} \inf \left\{ \sum_{i=1}^{\infty} r_i^t \right\}$$

where infimum is taken over all covers $\{B(x_i, r_i)\}_{i=1}^{\infty}$ of the set A by open balls centered at A with radii which do not exceed ε .

The t -dimensional outer packing measure $\Pi^t(A)$ of the set A is defined as

$$\Pi^t(A) = \inf_{\cup A_i = A} \left\{ \sum_i \Pi_*^t(A_i) \right\}$$

(A_i are arbitrary subsets of A), where

$$\Pi_*^t(A) = \sup_{\varepsilon > 0} \sup \left\{ \sum_{i=1}^{\infty} r_i^t \right\}.$$

Here the second supremum is taken over all packings $\{B(x_i, r_i)\}_{i=1}^{\infty}$ of the set A by open balls centered at A with radii which do not exceed ε . These two outer measures define countable additive measures on Borel σ -algebra of X .

The definition of the Hausdorff dimension $\text{HD}(A)$ of the set A is the following

$$\text{HD}(A) = \inf\{t : H^t(A) = 0\} = \sup\{t : H^t(A) = \infty\}.$$

Let ν be a Borel probability measure on X which is positive on open sets. Define the function $\rho = \rho_t(\nu) : X \times (0, \infty) \rightarrow (0, \infty)$ by

$$\rho(x, r) = \frac{\nu(B(x, r))}{r^t}$$

The following two theorems (see [26, 11], and [20]) are for our aims the key facts from geometric measure theory. Their proofs are an easy consequence of Besicovič covering theorem (see [26]) or a more elementary $4r$ -covering theorem (see [20]).

Theorem 6.1. *Let $X = \mathbb{R}^n$ for some $n \geq 1$. Then there exists a constant $b(n)$ depending only on n with the following properties. If A is a Borel subset of \mathbb{R}^n and $C > 0$ is a positive constant such that*

(1) *for all $x \in A$*

$$\limsup_{r \rightarrow 0} \rho(x, r) \geq C^{-1},$$

then for every Borel subset $E \subset A$ we have $H^t(E) \leq b(n)C\rho(E)$ and, in particular, $H^t(A) < \infty$,

or

(2) for all $x \in A$

$$\limsup_{r \rightarrow 0} \rho(x, r) \leq C^{-1},$$

then for every Borel subset $E \subset A$ we have $H^t(E) \geq C\rho(E)$.

(1)' If $t > 0$ then (1) holds under the weaker assumption that the hypothesis of part (1) is satisfied on the complement of a countable set.

Theorem 6.2. *Let $X = \mathbb{R}^n$ for some $n \geq 1$. Then there exists a constant $b(n)$ depending only on n with the following properties. If A is a Borel subset of \mathbb{R}^n and $C > 0$ is a positive constant such that*

(1) for all $x \in A$

$$\liminf_{r \rightarrow 0} \rho(x, r) \leq C^{-1},$$

then for every Borel subset $E \subset A$ we have $\Pi^t(E) \geq Cb(n)^{-1}\rho(E)$,

or

(2) for all $x \in A$

$$\liminf_{r \rightarrow 0} \rho(x, r) \geq C^{-1},$$

then $\Pi^t(E) \leq C\rho(E)$ and, consequently, $\Pi^t(A) < \infty$.

(1)' If ρ is non-atomic then (1) holds under the weaker assumption that the hypothesis of part (1) is satisfied on the complement of a countable set.

7. CONFORMAL MEASURES; EXISTENCE, UNIQUENESS, AND CONTINUITY

For every $t \geq 0$ and every function $\phi : J(\tilde{f}) \rightarrow \mathbb{C}$ let $\mathcal{L}_t\phi : J(\tilde{f}) \rightarrow \mathbb{C}$ be defined by the following formula:

$$\mathcal{L}_t\phi(y) = \sum_{x \in \tilde{f}^{-1}(y)} |\tilde{f}'(x)|^{-t} \phi(x).$$

$\mathcal{L}_t\phi(y)$ is finite if and only if $y \notin \text{Crit}(\tilde{f})$. Otherwise $\mathcal{L}_t\phi(y)$ is declared to be $+\infty$. Iterating this formula we get for all $n \geq 1$ that

$$\mathcal{L}_t^n\phi(y) = \sum_{x \in \tilde{f}^{-n}(y)} |(\tilde{f}^n)'(x)|^{-t} \phi(x).$$

If $y \in J(\tilde{f}) \setminus p_2^{-1}(G^*(\text{Crit}(f)_+))$, then $\mathcal{L}_t^n\mathbb{1}(y)$ is finite for all $n \geq 0$. If $\psi : \hat{\mathbb{C}} \rightarrow \mathbb{C}$, then define $\mathcal{L}_t\psi : \hat{\mathbb{C}} \rightarrow \mathbb{C}$ by the formula

$$\mathcal{L}_t\psi(z) = \sum_{i=1}^s \sum_{x \in f_i^{-1}(z)} |f_i'(x)|^{-t} \psi(x).$$

It will be always clear from the context whether \mathcal{L}_t is applied to a function defined on $J(\tilde{f})$ or on a compact neighborhood \mathcal{A} of $J(G)$. Iterating this formula we get for all $n \geq 1$ that

$$(7.1) \quad \mathcal{L}_t^n\psi(z) = \sum_{|\omega|=n} \sum_{x \in f_\omega^{-1}(z)} |f_\omega'(x)|^{-t} \psi(x).$$

Note that if $\tilde{\psi} : J(\tilde{f}) \rightarrow \mathbb{C}$ is defined by the formula $\tilde{\psi}(\tau, z) = \psi(z)$, then

$$\mathcal{L}_t^n \tilde{\psi}(\tau, z) = \mathcal{L}_t^n \psi(z)$$

for all $(\tau, z) \in J(\tilde{f})$. Without confusion we put $\tilde{\mathbb{1}} = \mathbb{1}$. Note that $\mathcal{L}_t^n \psi(z)$ is finite for all $z \in \mathcal{A} \setminus G^*(\text{Crit}(f)_+)$. For all $z \in \mathcal{A} \setminus G^*(\text{Crit}(f)_+)$ set

$$P_z(t) = \limsup_{n \rightarrow \infty} \frac{1}{n} \log \mathcal{L}_t^n \mathbb{1}(z) \in (-\infty, +\infty].$$

Definition 7.1. Denote by $\text{PCV}(\tilde{f})$ the closure of the postcritical set of \tilde{f} , i.e.

$$\text{PCV}(\tilde{f}) = \overline{\bigcup_{n=1}^{\infty} \tilde{f}^n(\text{Crit}(\tilde{f}))}.$$

Lemma 7.2. $\overline{G^*(\text{Crit}(f)_+)} \cap J(G)$ is a nowhere dense subset of $J(G)$ and $\text{PCV}(\tilde{f})$ is a nowhere dense subset of $J(\tilde{f})$.

Proof. Since, by Lemma 4.1, $\omega_G(\text{Crit}(f)_+) \cap J(G)$ is nowhere dense in $J(G)$ and since the set $\overline{G^*(\text{Crit}(f)_+)}$ is countable, it follows from the Baire Category Theorem the set $\overline{G^*(\text{Crit}(f)_+)} \cap J(G)$ is nowhere dense. In order to prove the second part of our lemma, suppose that $\text{PCV}(\tilde{f})$ is not nowhere dense in $J(\tilde{f})$. This means that $\text{PCV}(\tilde{f})$ has non-empty interior, and therefore, because of its forward invariance and topological exactness of the map $\tilde{f} : J(\tilde{f}) \rightarrow J(\tilde{f})$, we have $\text{PCV}(\tilde{f}) = J(\tilde{f})$. Hence $J(G) = p_2(J(\tilde{f})) = p_2(\text{PCV}(\tilde{f})) \subset \overline{G^*(\text{Crit}(f)_+)} \cap J(G)$, contrary to, the already proved, first part of the lemma. \square

We shall prove the following.

Lemma 7.3. The function $z \mapsto P_z(t)$ is constant throughout a neighborhood of $J(G) \setminus \overline{G^*(\text{Crit}(f)_+)}$.

Proof. For every $z \in J(G) \setminus \overline{G^*(\text{Crit}(f)_+)}$ fix $U_z = \{w \mid |w - z| < r\}$, an open round disk centered at z and such that $\{w \mid |w - z| < 2r\}$ is disjoint from $\overline{G^*(\text{Crit}(f)_+)}$. It then directly follows from Koebe's Distortion Theorem that the function $w \mapsto P_w(t)$ is constant on U_z . Now, fix $z_1, z_2 \in J(G) \setminus \overline{G^*(\text{Crit}(f)_+)}$. By [14, Lemma 3.2], there exists $g = f_\omega \in G$ such that $g(U_{z_1}) \cap U_{z_2} \cap J(G) \neq \emptyset$. Fix $x \in U_{z_1}$ such that $g(x) \in U_{z_2} \cap J(G)$. Then $x \in J(G)$ and for every $n \geq 1$, $\mathcal{L}_t^{n+|\omega|} \mathbb{1}(g(x)) \geq |g'(x)|^{-t} \mathcal{L}_t^n \mathbb{1}(x)$. Therefore, $P_{g(x)} t \geq P_x(t)$. Hence $P_{z_2}(t) \geq P_{z_1}(t)$. Exchanging the roles of z_1 and z_2 , we get $P_{z_1}(t) \geq P_{z_2}(t)$, and we are done. \square

By Lemma 7.2 the set $J(G) \setminus \overline{G^*(\text{Crit}(f)_+)}$ is not empty. Denote by $P(t)$ the constant common value of the function $z \mapsto P_z(t)$ on $J(G) \setminus \overline{G^*(\text{Crit}(f)_+)}$. $P(t)$ is called the topological pressure of t . Its basic properties are contained in the following.

Lemma 7.4. The function $t \mapsto P(t)$, $t \geq 0$, has the following properties.

- (a) $P(t)$ is non-increasing. In particular $P(t) < +\infty$ as clearly $P(0) < +\infty$.
- (b) $P(t)$ is convex and, hence, continuous.
- (c) $P(0) \geq \log 2 > 0$.
- (d) $P(2) \leq 0$.

Proof. Fix $z \in J(G) \setminus \overline{G^*(\text{Crit}(f)_+)}$. Since the family of all analytic inverse branches of all elements of G is normal in some neighborhood of z (see [36, Lemma 4.5]) and all its limit functions are constant (see Theorem 3.3), $\lim_{n \rightarrow \infty} \max\{|f'_\omega(x)| : |\omega| = n, x \in f_\omega^{-1}(z)\} = \infty$. So, item (a) follows directly from (7.1). Item (b), that is convexity of $P(t)$ follows directly from (7.1) and Hölder inequality. Item (c) follows from the fact that $\max\{u, \max\{\deg(f_j) : 1 \leq j \leq u\}\} \geq 2$. For the proof of item (d) let $U \subset \hat{\mathbb{C}}$ be the set coming from the nice open set condition. Fix $z \in J(G) \setminus \overline{G^*(\text{Crit}(f)_+)}$. Let $U_z = B(z, \frac{1}{2} \text{dist}(z, G^*(\text{Crit}(f)_+)))$. It follows from Koebe's Distortion Theorem that

$$|(g_*^{-1})'(z)|^2 \leq CK^2 \frac{l_2(g_*^{-1}(U_z \cap U))}{l_2(U_z \cap U)}$$

for all $g \in G$ and all analytic inverse branches g_*^{-1} of g defined on $B(z, \text{dist}(z, G^*(\text{Crit}(f)_+)))$, where $C > 0$ is a constant independent of g . Since, by the open set condition, all the sets $g_*^{-1}(U_z \cap U)$ are mutually disjoint, we thus get

$$\mathcal{L}_2^n \mathbb{1}(z) \leq CK^2 \frac{l_2(\bigcup g_*^{-1}(U_z \cap U))}{l_2(U_z \cap U)} \leq \frac{CK^2 l_2(U)}{l_2(U_z \cap U)}.$$

Hence $P(2) = P_z(2) \leq 0$ and we are done. \square

We say that a measure \tilde{m}_t on $J(\tilde{f})$ is $e^{P(t)}|\tilde{f}'|^t$ -conformal provided that

$$\tilde{m}_t(\tilde{f}(A)) = \int_A e^{P(t)}|\tilde{f}'|^t d\tilde{m}_t$$

for all Borel sets $A \subset J(\tilde{f})$ such that $\tilde{f}|_A$ is injective. If $P(t) = 0$, the measure \tilde{m}_t is simply referred to as t -conformal. Fix $z \in J(G) \setminus \overline{G^*(\text{Crit}(f)_+)}$. Observe that the critical parameter for the series

$$S_s(z) = \sum_{n=1}^{\infty} e^{-sn} \mathcal{L}_t^n \mathbb{1}(z)$$

is equal to the topological pressure $P(t)$, i.e. $S_s(z) = +\infty$ if $s < P(t)$ and $S_s(z) < +\infty$ if $s > P(t)$. For every σ -finite Borel measure m on $J(\tilde{f})$ let $\mathcal{L}_t^{*n} m$ be given by the formula

$$\mathcal{L}_t^{*n} m(A) = m(\mathcal{L}_t^n \mathbb{1}_A), \quad A \subset J(\tilde{f}).$$

Notice that if $(\tau, \xi) \in J(\tilde{f}) \setminus \bigcup_{n=1}^{\infty} \tilde{f}^n(\text{Crit}(\tilde{f}))$, then for all Borel sets $A \subset J(\tilde{f})$ we have

$$\mathcal{L}_t^{*n} \delta_{(\tau, \xi)}(A) = \delta_{(\tau, \xi)}(\mathcal{L}_t^n \mathbb{1}_A) = \mathcal{L}_t^n \mathbb{1}_A(\tau, \xi) = \sum_{|\omega|=n} \sum_{x \in A \cap f_\omega^{-1}(\xi)} |f'_\omega(x)|^{-t} \leq \mathcal{L}_t^n \mathbb{1}(\xi) < \infty.$$

In particular, $\mathcal{L}_t^{*n} \delta_{(\tau, \xi)}(J(\tilde{f})) \leq \mathcal{L}_t^n \mathbb{1}(\xi) < \infty$. Hence, if $s > P(t)$, then

$$(7.2) \quad \tilde{v}_s = S_s^{-1}(\xi) \sum_{n=1}^{\infty} e^{-sn} \mathcal{L}_t^{*n} \delta_{(\tau, \xi)}$$

is a Borel probability measure on $J(\tilde{f})$. Now, for every Borel set $A \subset J(\tilde{f})$ we have

$$\mathcal{L}_t^{*n} \delta_{(\tau, \xi)}(A) = \delta_{(\tau, \xi)}(\mathcal{L}_t^n \mathbf{1}_A) = \mathcal{L}_t^n \mathbf{1}_A(\tau, \xi) = \sum_{(\gamma, z) \in \tilde{f}^{-n}(\tau, \xi) \cap A} |(\tilde{f}^n)'(\gamma, z)|^{-t}.$$

So, $\mathcal{L}_t^{*n} \delta_{(\tau, \xi)}(\tilde{f}^{-n}(\tau, \xi)) = 1$. Hence, denoting

$$\nu_s = \tilde{\nu}_s \circ p_2^{-1},$$

we get the following.

Lemma 7.5. *We have $\tilde{\nu}_s(\bigcup_{n=0}^{\infty} f^{-n}(\tau, \xi)) = 1$ and $\nu_s(G^{-1}(\xi)) = 1$.*

In what follows that we are in the divergence type, i.e. $S_{P(t)}(\xi) = +\infty$. For the convergence type situation the usual modifications involving slowly varying functions have to be done, the details can be found in [9]. The following lemma is proved by a direct straightforward calculations.

Lemma 7.6. *For every $s > P(t)$ the following hold.*

- (a) $\tilde{\nu}_s$ is a Borel probability measure.
- (b) For every continuous function $g : J(\tilde{f}) \rightarrow \mathbb{R}$, we have

$$\int g d\tilde{\nu}_s = S_s^{-1}(\xi) \sum_{n=1}^{\infty} e^{-sn} \mathcal{L}_t^n g d\delta_{(\tau, \xi)} = S_s^{-1}(\xi) \sum_{n=1}^{\infty} e^{-sn} \mathcal{L}_t^n g(\tau, \xi).$$

(c)

$$e^{-s} \mathcal{L}_t^* \tilde{\nu}_s = S_s^{-1}(\xi) \sum_{n=1}^{\infty} e^{-s(n+1)} \mathcal{L}_t^{*(n+1)} \delta_{(\tau, \xi)} = \tilde{\nu}_s - S_s^{-1}(\xi) (e^{-s} \mathcal{L}_t^* \delta_{(\tau, \xi)}).$$

Now we can easily prove the following.

Proposition 7.7. *For every $t \geq 0$ there exists an $e^{P(t)} |\tilde{f}'|^t$ -conformal measure \tilde{m}_t for the map $\tilde{f} : J(\tilde{f}) \rightarrow J(\tilde{f})$.*

Proof. Since $\lim_{s \searrow P(t)} S_s(\xi) = +\infty$, it suffices to take as \tilde{m}_t any weak limit of $\tilde{\nu}_s$ when $s \searrow P(t)$, and to apply Lemma 7.6(c). \square

Consider now a Borel set $A \subset J(\tilde{f})$ such that $\tilde{f}|_A$ is injective. It then follows from Lemma 7.6(c) that

$$\begin{aligned}
(7.3) \quad \tilde{\nu}_s(A) &= e^{-s} \mathcal{L}_t^* \tilde{\nu}_s(\mathbb{1}_A) + S_s^{-1}(\xi) e^{-s} \mathcal{L}_t^* \delta_{(\tau, \xi)}(\mathbb{1}_A) \\
&= e^{-s} \int \mathcal{L}_t(\mathbb{1}|_A) d\tilde{\nu}_s + e^{-s} S_s^{-1}(\xi) \int \mathcal{L}_t(\mathbb{1}|_A) d\delta_{(\tau, \xi)} \\
&= e^{-s} \int \sum_{y \in \tilde{f}^{-1}(x)} |\tilde{f}'(y)|^{-t} \mathbb{1}_A(y) d\tilde{\nu}_s(x) + e^{-s} S_s^{-1}(\xi) \mathcal{L}_t(\mathbb{1}|_A)(\tau, \xi) \\
&= e^{-s} \int_{\tilde{f}(A)} |(\tilde{f}|_A^{-1})'(x)|^t d\tilde{\nu}_s(x) + \begin{cases} 0 & \text{if } A \cap \tilde{f}^{-1}(\tau, \xi) = \emptyset \\ e^{-s} S_s^{-1}(\xi) |\tilde{f}'(y)|^{-t} & \text{if } A \cap \tilde{f}^{-1}(\tau, \xi) = \{y\}. \end{cases}
\end{aligned}$$

Suppose now that $(\omega, x) \in J(\tilde{f})$ and there exists a (unique) continuous inverse branch $\phi_{(\omega, x)}^{-1} : \Sigma_u \times B(f_{\omega_1}(x), 2R) \rightarrow \Sigma_u \times \mathbb{C}$ of f sending $(\sigma(\omega), f_{\omega_1}(x))$ to (ω, x) . It then follows from (7.3) and Lemma 7.6(c) that for every set $A \subset \Sigma_u \times B(f_{\omega_1}(x), 2R)$, we have that

$$(7.4) \quad \tilde{\nu}_s(\phi_{(\omega, x)}^{-1}(A)) = e^{-s} \int_A |(\phi_{(\omega, x)}^{-1})'|^t d\tilde{\nu}_s + e^{-s} S_s^{-1}(\xi) |(\phi_{(\omega, x)}^{-1})'(\tau, \xi)|^t \delta_{(\tau, \xi)}(A)$$

From now on throughout the paper we assume that

$$(7.5) \quad P(t) \geq 0.$$

We also require that

$$(7.6) \quad \xi \notin \overline{G^*(\text{Crit}(f)_+)}.$$

Our goal now is to show that the measure

$$m_t = \tilde{m}_t \circ p_2^{-1}$$

is uniformly upper t -estimable. For every critical point $c \in \text{Crit}(f)$ let

$$I(c) = \{1 \leq i \leq u : f'_i(c) = 0\}$$

and let

$$\Sigma(c) = \bigcup_{i \in I(c)} [i] \subset \Sigma_u.$$

Now suppose that Γ is a closed subset of $J(G)$ such that $g(\Gamma) \cap J(G) \subset \Gamma$ for each $g \in \Gamma$, and that \tilde{m} is a Borel probability measure on $J(\tilde{f})$.

Definition 7.8. *The measure \tilde{m} is said to be nearly upper t -conformal relative to Γ provided that there exists an $S > 0$ such that the following conditions are satisfied.*

(a) For every $z \in \Gamma$

$$\tilde{m}(\tilde{f}(A)) \geq \int_A |\tilde{f}'|^t d\tilde{m}$$

for every Borel sets $A \subset J(\tilde{f}) \cap p_2^{-1}(B(z, S))$ such that $\tilde{f}|_A$ is injective.

- (b) For every $c \in \text{Crit}(f)$ such that $\bigcup_{|\tau|=l} f_\tau(c_+) \cap J(G) \subset \Gamma$ (the integer $l = l(f) \geq 0$ coming from Lemma 4.8) and every $1 \leq j \leq l+1$,

$$\tilde{m}(\tilde{f}^j(A)) \geq \int_A |(\tilde{f}^j)'|^t d\tilde{m}$$

for every Borel sets $A \subset J(\tilde{f}) \cap p_2^{-1}(B(c, S))$ such that $\tilde{f}^j|_A$ is injective.

- (c) $\tilde{m}(\Sigma(c) \times \{c\}) = 0$ for every point $c \in \Gamma \cap \text{Crit}(f)$.

The constant S is said to be the nearly upper conformality radius. If $\Gamma = J(G)$, we simply say that \tilde{m} is nearly upper t -conformal. In any case put

$$m = \tilde{m} \circ p_2^{-1}.$$

Let us prove the following.

Lemma 7.9. *Suppose that Γ is a closed subset of $J(G)$ such that $g(\Gamma) \cap J(G) \subset \Gamma$ for each $g \in G$, and that \tilde{m} is a Borel probability nearly upper t -conformal measure on $J(\tilde{f})$ respective to Γ . Fix $i \in \{0, 1, \dots, p\}$ and suppose that for every critical point $c \in S_i(f) \cap \Gamma$ the measure $\tilde{m}|_{\Sigma(c) \times \hat{c}} \circ p_2^{-1}$ is upper t -estimable at c . Then the measure m is uniformly upper t -estimable at all points $z \in J_i(G) \cap \Gamma$.*

Proof. Since Γ is a closed set and $\text{Crit}(f)$ is finite, the number $\Delta = \text{dist}(\Gamma, \text{Crit}(f) \setminus \Gamma)$ is positive (if $\text{Crit}(f) \setminus \Gamma = \emptyset$ then we put $\Delta = \infty$). Fix $\theta \in (0, \min\{1, \gamma\})$ so small that

$$(7.7) \quad \theta \|f'\|^{-1} < \min\{\Delta, \rho\}.$$

Put

$$\alpha = \theta(KA_f^2)^{-1} 2^{-\#\text{Crit}(f)}.$$

Let $z \in J_i(G) \cap \Gamma$. Fix $\tau \in \Sigma_u$ such that $(\tau, z) \in J(\tilde{f})$, i.e. $\tau \in p_1(J(\tilde{f}) \cap p_2^{-1}(z))$. Assume $r \in (0, R_f]$ to be sufficiently small. Let $s(\tau, r) = s(\theta, (\tau, z), 8\alpha^{-1}r) \geq 0$ be the integer produced in Proposition 5.3. Set $R_{\tau|_{s(\tau, r)+1}} = 4r|f'_{\tau|_{s(\tau, r)}}(z)|$. It then follows from Proposition 5.3 that the family

$$\mathcal{F}(z, r) = \{\tau|_{s(\tau, r)+1} : \tau \in p_1(J(\tilde{f}) \cap p_2^{-1}(z))\}$$

is $(4, \gamma, V)$ -essential for the pair (z, r) , where $V = \bigcup\{\tau|_{s+1} : \tau \in p_1(J(\tilde{f}) \cap p_2^{-1}(z))\}$. Keep $\tau \in p_1(J(\tilde{f}) \cap p_2^{-1}(z))$ and $s = s(\tau, r)$. Suppose that the first alternative of (b) in Proposition 5.3 holds. Then $8\alpha^{-1}r|f'_{\tau|_s}(z)| > \|f'\|^{-1}$. So, using Koebe's Distortion Theorem, and assuming θ is small enough, we get from nearly upper t -conformality of \tilde{m} respective to Γ that

$$(7.8) \quad \begin{aligned} \tilde{m}(f_{\tau|_s, z}^{-s}([\tau_{s+1}] \times B(f_{\tau|_s}(z), R_{\tau|_{s+1}}))) &\leq \tilde{m}(f_{\tau|_s, z}^{-s}(p_2^{-1}(B(f_{\tau|_s}(z), R_{\tau|_{s+1}})))) \\ &\leq K^t |f'_{\tau|_s}(z)|^{-t} \tilde{m} p_2^{-1}(B(f_{\tau|_s}(z), R_{\tau|_{s+1}})) \\ &\leq K^t |f'_{\tau|_s}(z)|^{-t} \\ &\leq (8K\alpha^{-1} \|f'\|)^t r^t. \end{aligned}$$

Now Suppose $8\alpha^{-1}r|f'_{\tau|_s}(z)| \leq \|\tilde{f}'\|^{-1}$ which particular implies that the second alternative of (b) in Proposition 5.3 holds. Let $c \in \text{Crit}(f_{\tau|_{s+1}})$ such that $f_{\tau|_{s+1}}(c) \in J(G)$ come from item (b) of this proposition. Since $z \in J_i(G)$ (and $\theta\|\tilde{f}'\|^{-1} < \rho$), it follows from (4.5) and Proposition 5.3 that $c \in S_i(f)$. Since $8\alpha^{-1}r|f'_{\tau|_s}(z)| \leq \|f'\|^{-1}$, it follows from Proposition 5.3(b) and (7.7) that $|f_{\tau|_s}(z) - c| \leq \theta\|f'\|^{-1} < \Delta$. Thus $c \in \Gamma$. Hence, making use of Proposition 5.3(b), (c), as well as Koebe's Distortion Theorem, nearly upper t -conformality of \tilde{m} , and our t -upper estimability assumption, and assuming θ is small enough, we get with some universal constant C_1 that

$$\begin{aligned}
& \tilde{m}(f_{\tau|_s, z}^{-s}([\tau_{s+1}] \times B(f_{\tau|_s}(z), R_{\tau|_{s+1}}))) \\
& \leq K^t |f'_{\tau|_s}(z)|^{-t} \tilde{m}([\tau_{s+1}] \times B(f_{\tau|_s}(z), R_{\tau|_{s+1}})) \\
& \leq K^t |f'_{\tau|_s}(z)|^{-t} \tilde{m}|_{\Sigma(c) \times \hat{\mathbb{C}}} \circ p_2^{-1}(B(f_{\tau|_s}(z), R_{\tau|_{s+1}})) \\
& \leq K^t |f'_{\tau|_s}(z)|^{-t} \tilde{m}|_{\Sigma(c) \times \hat{\mathbb{C}}} \circ p_2^{-1}(B(c, R_{\tau|_{s+1}} + 8\theta\alpha^{-1}r|f'_{\tau|_s}(z)|)) \\
& \leq K^t |f'_{\tau|_s}(z)|^{-t} \tilde{m}|_{\Sigma(c) \times \hat{\mathbb{C}}} \circ p_2^{-1}(B(c, 4(1 + 2\theta\alpha^{-1})r|f'_{\tau|_s}(z)|)) \\
& \leq K^t |f'_{\tau|_s}(z)|^{-t} C_1 (4(1 + 2\theta\alpha^{-1})r|f'_{\tau|_s}(z)|)^t \\
& = C_1 (4K(1 + 2\theta\alpha^{-1}))^t r^t.
\end{aligned}$$

Combining this with (7.8) and applying Proposition 2.15, we get that

$$(7.9) \quad m(B(z, r)) \leq \#_{4, \gamma} C_1 \max\{8K\alpha^{-1}\|\tilde{f}'\|, 4K(1 + 2\theta\alpha^{-1})\}^t r^t.$$

We are done. \square

Lemma 7.10. *There are two functions $(R, S) \mapsto R^*$ and $L \mapsto \hat{L}$ with the following property.*

- *Suppose that Γ is a closed subset of $J(G)$ such that $g(\Gamma) \cap J(G) \subset \Gamma$ for each $g \in G$, and that \tilde{m} is a Borel probability nearly upper t -conformal measure on $J(\tilde{f})$ respective to Γ with nearly upper conformality radius S . Fix $i \in \{0, 1, \dots, p\}$ and suppose that the measure m is uniformly upper t -estimable at all points $z \in J_i(G) \cap \Gamma$ with corresponding estimability constant L and estimability radius R . Then the measure $\tilde{m}|_{\Sigma(c) \times \hat{\mathbb{C}}} \circ p_2^{-1}$ is t -upper estimable, with upper estimability constant \hat{L} and radius R^* at every point $c \in Cr_{i+1}(f)$ such that $\bigcup_{|\omega|=l} f_\omega(c_+) \cap J(G) \subset \Gamma$.*

Proof. Fix $c \in Cr_{i+1}(f)$ such that $\bigcup_{|\omega|=l} f_\omega(c_+) \subset \Gamma$ and also $j \in \{0, 1, \dots, u\}$ such that $f'_j(c) = 0$. Consider an arbitrary $\tau \in \Sigma_u$ such that $\tau_1 = j$ and $(\tau, c) \in J(\tilde{f})$. In view of Lemma 4.8

$$f_{\tau|_{l+1}}(c) \in J_i(G) \cap \Gamma.$$

Let $R > 0$ (sufficiently small) be the radius resulting from uniform t -upper estimability at all points of $J_i(G) \cap \Gamma$. Let $D_{\tau|_{l+1}}(c)$ be the connected component of $f_{\tau|_{l+1}}^{-1}(B(f_{\tau|_{l+1}}(c), R))$ containing c . Set

$$\nu_{\tau|_{l+1}} = \tilde{m}|_{[\tau|_{l+1}] \times \hat{\mathbb{C}}} \circ p_2^{-1}|_{D_{\tau|_{l+1}}(c)}.$$

Applying nearly upper t -conformality of \tilde{m} we get for every Borel set $A \subset D_{\tau|_{l+1}}(c) \setminus \{c\}$ such that $f_{\tau|_{l+1}}|_A$ is injective, the following.

$$m(f_{\tau|_{l+1}}(A)) = \tilde{m}(\Sigma_u \times f_{\tau|_{l+1}}(A)) = \tilde{m}(f^{l+1}([\tau|_{l+1}] \times A)) \geq \int_A |f'_{\tau|_{l+1}}(x)|^t d\nu_{\tau|_{l+1}}(x).$$

It therefore follows from Lemma 2.10 and item (c) of Definition 7.8 that the measure $\nu_{\tau|_{l+1}}$ is upper t -estimable at c with upper estimability constant L_0 and radius R_0 independent of \tilde{m} (but possibly R_0 depends on (R, S) and L_0 depends on L). Let

$$\mathcal{F} = \{\tau|_{l+1} : (\tau, c) \in J(\tilde{f}) \text{ and } f'_{\tau_1}(c) = 0\}.$$

Let $D_c = \bigcap_{\omega \in \mathcal{F}} D_\omega(c)$. Since $\#\mathcal{F} \leq u^{l+1}$ and since

$$\tilde{m}|_{\Sigma(c) \times \hat{c}} \circ p_2^{-1}|_{D_c} = \sum_{\omega \in \mathcal{F}} \nu_\omega|_{D_c},$$

we conclude that the measure $\tilde{m}|_{\Sigma(c) \times \hat{c}} \circ p_2^{-1}$ is t -upper estimable at the point c with upper estimability constant \hat{L} and radius R^* independent of \tilde{m} . We are done. \square

Now, a straightforward inductive reasoning based on Lemma 7.9 and (7.9), (which also give the base of induction since $S_0(f) = \emptyset$), and Lemma 7.10 yields the following.

Lemma 7.11. *Suppose that Γ is a closed subset of $J(G)$ such that $g(\Gamma) \cap J(G) \subset \Gamma$ for each $g \in G$, and that \tilde{m} is a Borel probability nearly upper t -conformal measure on $J(\tilde{f})$ relative to Γ with nearly upper conformality radius S . Then the measure $m = \tilde{m} \circ p_2^{-1}$ is uniformly upper t -estimable at every point of Γ and $\tilde{m}|_{\Sigma(c) \times \hat{c}} \circ p_2^{-1}$ is upper t -estimable, with upper estimability constants and radii independent of the measure \tilde{m} (but possibly dependent on S), at every point $c \in \Gamma \cap \text{Crit}(f)$.*

Now we are in the position to prove the following.

Lemma 7.12. *If $P(t) \geq 0$, then the measure $m_t = \tilde{m}_t \circ p_2^{-1}$ is uniformly upper t -estimable.*

Proof. Fix $s > P(t) \geq 0$ and consider the measure $\tilde{\nu}_s$ defined in (7.2). We want to apply Lemma 7.11 with $\Gamma = \overline{G^*(\text{Crit}(f)_+ \cap J(G))} \cap J(G)$ and $\tilde{m} = \tilde{\nu}_s$. For this we have to check that $\tilde{\nu}_s$ is nearly upper t -conformal relative to Γ . Condition (c) of Definition 7.8 follows directly from Lemma 7.5 and the fact that $\xi \notin G(\text{Crit}(f))$ (see (7.6)). Since $\xi \notin \Gamma$ and $G(\Gamma) \cap J(G) \subset \Gamma$, there exists an $S_0 > 0$ such that $\xi \notin \bigcup_{j=1}^u f_j(B(\Gamma, S_0)) \cap J(G)$. Formula (7.4) then yields that for every $z \in \Gamma$,

$$\tilde{\nu}_s(\tilde{f}(A)) = e^s \int_A |\tilde{f}'|^t d\tilde{\nu}_s \geq \int_A |\tilde{f}'|^t d\tilde{\nu}_s$$

for every Borel set $A \subset \Sigma_u \times B(z, S_0)$ such that $\tilde{f}|_A$ is injective. Thus, condition (a) of Definition 7.8 is also verified. Condition (b) of this definition follows by iterating the above argument $l + 1$ times and keeping in mind that $\xi \notin \overline{G^*(\text{Crit}(f)_+)}$. Hence, there exists a

constant S such that for each $s > P(t)$, $\tilde{\nu}_s$ is nearly upper t -conformal respective to Γ with nearly upper conformality radius S . Therefore, Lemma 7.11 applies and we conclude that all measures $\tilde{\nu}_s|_{\Sigma(c) \times \hat{c}} \circ p_2^{-1}$ are upper t -estimable at respective points $c \in \text{Crit}(f) \cap J(G)$ with estimability constants and radii independent of $s > P(t)$. Therefore, \tilde{m}_t , a weak limit of measures $\tilde{\nu}_s$, $s > P(t)$, (see the proof of Proposition 7.7) also enjoys the property that $\tilde{m}_t|_{\Sigma(c) \times \hat{c}} \circ p_2^{-1}$ is upper t -estimable at respective points $c \in \text{Crit}(f) \cap J(G)$. Consequently $\tilde{m}_t(\Sigma(c) \times \{c\}) = 0$. Having this we immediately see from Proposition 7.7 that the measure \tilde{m}_t is nearly upper t -conformal, i.e. respective to $\Gamma = J(G)$. So, applying Lemma 7.11, we conclude that the measure $m_t = \tilde{m}_t \circ p_2^{-1}$ is uniformly upper t -estimable at every point of $\Gamma = J(G)$. We are done. \square

Now we assume that $t = h$, i.e. $P(t) = 0$ and we deal with the problem of lower estimability. It is easier than the upper one. We start with the following.

Lemma 7.13. *Fix $i \in \{0, 1, \dots, p\}$ and suppose that for every critical point $c \in S_i(f)$ and every $j \in I(c)$ the measure $\tilde{m}_h|_{[j] \times \hat{c}} \circ p_2^{-1}$ is strongly lower h -estimable at c with sufficiently small lower estimability size. Then m_h is uniformly strongly lower h -estimable at all points of $J_i(G)$.*

Proof. Let $\theta \in (0, \min\{1, \gamma\})$ be such that $\theta \|\tilde{f}'\|^{-1} < \rho$. Put $\alpha := \theta^{-1} K A_f^2 2^{\#\text{Crit}(f)+5}$.

$$\lambda = \max\{\lambda(c) : c \in S_i(f)\},$$

where all $\lambda(c)$ are lower estimability sizes at respective critical points c . Fix $z \in J_i(G) \setminus S_i(f)$ and take $\tau \in \Sigma_u$ such that $(\tau, z) \in J(\tilde{f})$. Assume $r > 0$ to be sufficiently small. Let $s = s(\theta, (\tau, z), \alpha r) \geq 0$ be the integer produced in Proposition 5.3 for the point z and radius r . A straightforward calculation based on Proposition 7.7 shows that

$$\nu_1 = \tilde{m}_h|_{[\tau|_s] \times f_{(\tau|_s, z)}^{-1}(B(f_{\tau|_s}(z), 32r|f'_{\tau|_s}(z)|))} \circ p_2^{-1} \quad \text{and} \quad \nu_2 = m_h|_{B(f_{\tau|_s}(z), 32r|f'_{\tau|_s}(z)|)}$$

form an h -conformal pair of measures with respect to the map

$$f_{\tau|_s} : f_{\tau|_s, z}^{-1}(B(f_{\tau|_s}(z), 32r|f'_{\tau|_s}(z)|)) \rightarrow B(f_{\tau|_s}(z), 32r|f'_{\tau|_s}(z)|).$$

By Koebe's $\frac{1}{4}$ -Theorem (Theorem 2.1) for every $x \in B(z, r)$ we have

$$(7.10) \quad B(x, r) \subset f_{\tau|_s, z}^{-1}(B(f_{\tau|_s}(z), 8r|f'_{\tau|_s}(z)|)).$$

So,

$$B(f_{\tau|_s}(x), r|f'_{\tau|_s}(z)|) \subset B(f_{\tau|_s}(z), 9r|f'_{\tau|_s}(z)|).$$

By Koebe's Distortion Theorem we also get (with small enough λ)

$$B(x, \lambda r) \supset f_{\tau|_s, z}^{-1}(B(f_{\tau|_s}(x), K^{-1}\lambda r|f'_{\tau|_s}(z)|)).$$

In virtue of Koebe's Distortion Theorem and t -conformality of the pair (ν_1, ν_2) , we get as a consequence of all of this that

$$\begin{aligned} m_h(B(x, \lambda r)) &\geq \nu_1(B(x, \lambda r)) \geq \nu_1(f_{\tau|s,z}^{-1}(B(f_{\tau|s}(x), K^{-1}\lambda r|f'_{\tau|s}(z)|))) \\ &= \int_{B(f_{\tau|s}(x), K^{-1}\lambda r|f'_{\tau|s}(z)|)} |(f_{\tau|s,z}^{-1})'|^h d\nu_2 \\ &\geq K^{-h}|f'_{\tau|s}(z)|^{-h}\nu_2(B(f_{\tau|s}(x), K^{-1}\lambda r|f'_{\tau|s}(z)|)). \end{aligned}$$

Suppose now that the first alternative in Proposition 5.3(b) holds. We then can continue the above estimate as follows.

$$(7.11) \quad \begin{aligned} m_h(B(x, \lambda r)) &\geq K^{-h}|f'_{\tau|s}(z)|^{-h}\nu_2(B(f_{\tau|s}(x), K^{-1}\lambda||f'|^{-1})) \\ &= K^{-h}|f'_{\tau|s}(z)|^{-h}m_h(B(f_{\tau|s}(x), K^{-1}\lambda||f'|^{-1})) \end{aligned}$$

By conformality the measure \tilde{m}_h is positive on open subsets of $J(\tilde{f})$, and so, the measure m_h is positive on open subsets of $J(G)$. Therefore, for every $R > 0$,

$$M_R = \inf\{m_h(B(w, R)) : w \in J(G)\} > 0.$$

Hence, (7.11) gives that

$$m_h(B(x, \lambda r)) \geq K^{-h}M_{K^{-1}\lambda||f'|^{-1}}|f'_{\tau|s}(z)|^{-h}$$

By minimality of $s = s(\theta, (\tau, z), \alpha r)$ we have $\alpha r|f'_{\tau|s-1}(z)| \leq ||f'|^{-1}$ ($s \geq 1$ assuming $r > 0$ to be sufficiently small). Hence $|f'_{\tau|s}(z)| \leq (\alpha r)^{-1}$, and therefore

$$m_h(B(x, \lambda r)) \geq K^{-h}M_{K^{-1}\lambda||f'|^{-1}}\alpha^h r^h.$$

So suppose that $\alpha r||(\tilde{f}^s)'(\tau, z)|| \leq ||f'|^{-1}$ and the second alternative in Proposition 5.3(b) holds. Let $c \in \text{Crit}(f_{\tau_{s+1}})$ be such that $f_{\tau_{s+1}}(c) \in J(G)$ and $|f_{\tau|s}(z) - c| \leq \theta\alpha r|f'_{\tau|s}(z)| \leq \theta||\tilde{f}'||^{-1} < \rho$. Since $z \in J_i(G)$, we obtain $c \in S_i(f)$. Then, using (7.10), we get

$$f_{\tau|s}(x) \in B(f_{\tau|s}(z), 8r|f'_{\tau|s}(z)|) \subset B(c, (\theta\alpha + 8)r|f'_{\tau|s}(z)|)$$

and

$$B(f_{\tau|s}(x), \lambda(\theta\alpha + 8)r|f'_{\tau|s}(z)|) \subset B(f_{\tau|s}(z), (8 + \lambda(\theta\alpha + 8))r|f'_{\tau|s}(z)|) \subset B(f_{\tau|s}(z), 9r|f'_{\tau|s}(z)|)$$

if $\lambda > 0$ is small enough. Hence, using conformality of the pair (ν_1, ν_2) , Koebe's Distortion Theorem, the fact that $\tau_{s+1} \in I(c)$, and the lower h -estimability $\tilde{m}_h|_{[\tau_{s+1}] \times \hat{C}} \circ p_2^{-1}$ at the point c , we get that

$$\begin{aligned} m_h(B(x, K\lambda(\theta\alpha + 8)r)) &\geq \nu_1(B(x, K\lambda(\theta\alpha + 8)r)) \geq \nu_1(f_{(\tau|s,z)}^{-1}(B(f_{\tau|s}(x), \lambda(\theta\alpha + 8)r|f'_{\tau|s}(z)|))) \\ &\geq K^{-h}|f'_{\tau|s}(z)|^{-h}\nu_2(B(f_{\tau|s}(x), \lambda(\theta\alpha + 8)r|f'_{\tau|s}(z)|)) \\ &\geq K^{-h}|f'_{\tau|s}(z)|^{-h}\tilde{m}_h|_{[\tau_{s+1}] \times \hat{C}} \circ p_2^{-1}(B(f_{\tau|s}(x), \lambda(\theta\alpha + 8)r|f'_{\tau|s}(z)|)) \\ &\geq K^{-h}|f'_{\tau|s}(z)|^{-h}L_0((\theta\alpha + 8)r|f'_{\tau|s}(z)|)^h \\ &= L_0((\theta\alpha + 8)K^{-1})^h r^h, \end{aligned}$$

where L_0 is a constant independent of x and r . So, we are done with the lower estimability size $K\lambda(\theta\alpha + 8)$. \square

Now we shall prove the following.

Lemma 7.14. *Fix $i \in \{0, 1, \dots, p\}$ and suppose that the measure m_h is uniformly strongly lower h -estimable at all points of $J_i(G)$. Then the measure $\tilde{m}_h|_{[j] \times \hat{\mathbb{C}}} \circ p_2^{-1}$ is strongly lower h -estimable at every critical point $c \in Cr_{i+1}(f)$ and every $j \in I(c)$.*

Proof. Fix $c \in Cr_{i+1}(f)$ and then an arbitrary $j \in I(c)$. Next consider an arbitrary $\tau \in \Sigma_u$ such that $\tau_1 = j$ and $(\tau, c) \in J(\tilde{f})$. Now, ignoring Γ , follow the proof of Lemma 7.10 up to the definition of the measure $\nu_{\tau|_{l+1}}$. It follows from conformality of \tilde{m}_h that the measure $\nu_{\tau|_{l+1}}$ on $D_{\tau|_{l+1}}(c)$ and $\tilde{m}_h|_{\Sigma_u \times B(f_{\tau|_{l+1}}(c), R)} \circ p_2^{-1} = m_h|_{B(f_{\tau|_{l+1}}(c), R)}$ form an h -conformal pair of measures for the map $f_{\tau|_{l+1}} : D_{\tau|_{l+1}}(c) \rightarrow B(f_{\tau|_{l+1}}(c), R)$. So the measure $\nu_{\tau|_{l+1}}$ is strongly lower h -estimable at c in virtue of our assumption and Lemma 2.11. Since $D_{\tau|_{l+1}}(c)$ is an open neighborhood of c and $[\tau]_{l+1} \times D_{\tau|_{l+1}}(c) \subset [j] \times \hat{\mathbb{C}}$, we thus see that the measure $\tilde{m}_h|_{[j] \times \hat{\mathbb{C}}} \circ p_2^{-1}$ is strongly lower h -estimable at c . We are done. \square

The second main result of this section is this.

Lemma 7.15. *The measure $m_h = \tilde{m}_h \circ p_2^{-1}$ is uniformly strongly lower h -estimable.*

Proof. Having $J_p(G) = J(G)$ (Lemma 4.7) the proof of this lemma is the obvious mathematical induction based on Lemma 7.13 and Lemma 7.14 as inductive steps and Lemma 7.13 with $i = 0$ (then $S_i(G) = \emptyset$ and its hypothesis are vacuously fulfilled) serving as the base of induction. \square

Since every uniformly strongly lower h -estimable measure is uniformly lower h -estimable, as an immediate consequence of Lemma 7.12, Lemma 7.15, and [11, 19, 26], we obtain the following main result of this section and one of the two main results of the entire paper.

Theorem 7.16. *Under Assumption (*), we have the following.*

- (a) *The measure $m_h = \tilde{m}_h \circ p_2^{-1}$ is geometric meaning that there exists a constant $C \geq 1$ such that*

$$C^{-1} \leq \frac{m_h(B(z, r))}{r^h} \leq C$$

for all $z \in J(G)$ and all $r \in (0, 1]$.

Consequently,

- (b) $h = \text{HD}(J(G)) = \text{PD}(J(G)) = \text{BD}(J(G))$.
(c) $\text{HD}(J(G))$ is the unique zero of $t \mapsto P(t)$.
(d) All the measures H^h , P^h , and m_h are mutually equivalent with Radon-Nikodym derivatives uniformly separated away from zero and infinity.

In particular

- (e) $0 < H^h(J(G)), P^h(J(G)) < +\infty$.

Definition 7.17. *The unique zero of $t \mapsto P(t)$ is denoted by $h = h(f)$. Note that $h(f) = \text{HD}(J(G)) = \text{PD}(J(G)) = \text{BD}(J(G))$.*

Corollary 7.18. *Under Assumption (*), for each $z \in J(G) \setminus \overline{G^*(\text{Crit}(f)_+)}$, we have $h(f) = T_f(z) = t_0(f) = S_G(z) = s_0(G) = \text{HD}(J(G)) = \text{PD}(J(G)) = \text{BD}(J(G))$.*

Proof. Let $z \in J(G) \setminus \overline{G^*(\text{Crit}(f)_+)}$. Since G satisfies the open set condition, G is a free semigroup. Hence $T_f(z) = S_G(z)$ and $t_0(f) = s_0(G)$. Moreover, by [37, Theorem 5.7], we have $\text{HD}(J(G)) \leq s_0(G) \leq S_G(z)$. We now let $a > h(f)$. Since $h(f)$ is the unique zero of $P(t)$ and since $t \mapsto P(t)$ is non-increasing function, we have $P(a) < 0$. Hence there exists a number $v < 0$ such that for each $n \in \mathbb{N}$, $\sum_{|\omega|=n} \sum_{x \in f_\omega^{-1}(z)} |f'_\omega(x)|^{-a} \leq e^{nv}$. Therefore $T_f(z) \leq a$. Thus $T_f(z) \leq h(f)$. Since $h(f) = \text{HD}(J(G))$, it follows that $h(f) = T_f(z) = t_0(f) = S_G(z) = s_0(G) = \text{HD}(J(G)) = \text{PD}(J(G)) = \text{BD}(J(G))$. We are done. \square

It follows from Theorem 7.16 that the measure m_h is atomless. We thus get the following.

Corollary 7.19. *Under Assumption (*), we have $\tilde{m}_h(\text{Sing}(\tilde{f})) = 0$.*

Proof. Indeed, the set $\text{Crit}(f)$ is finite and so, $G^{-1}(\text{Crit}(f))$ is countable. For all $n \geq 0$ we have

$$\tilde{f}^{-n}(\text{Crit}(\tilde{f})) \subset p_2^{-1}(p_2(\tilde{f}^{-n}(\text{Crit}(\tilde{f})))) \subset p_2^{-1}(G^{-1}(\text{Crit}(f))).$$

Hence, $\tilde{m}_h(\tilde{f}^{-n}(\text{Crit}(\tilde{f}))) \leq m_h(G^{-1}(\text{Crit}(f))) = 0$. Since $\text{Sing}(\tilde{f}) = \bigcup_{n=0}^{\infty} \tilde{f}^{-n}(\text{Crit}(\tilde{f}))$, we are thus done. \square

8. INVARIANT MEASURES

In this section we prove that there exists a unique Borel probability \tilde{f} -invariant measure on $J(\tilde{f})$ which is absolutely continuous with respect to \tilde{m}_h . This measure is proved to be metrically exact, in particular ergodic.

Frequently in order to denote that a Borel measure μ is absolutely continuous with respect to ν we write $\mu \prec \nu$. We do not use any special symbol to record equivalence of measures (mutual absolute continuity).

We use some notations from [1]. Given a σ -finite measure space Let (X, \mathcal{F}, μ) be a σ -finite measure space and let $T : X \rightarrow X$ be a measurable almost everywhere defined transformation. T is said to be nonsingular if and only if for any $A \in \mathcal{F}$, $\mu(T^{-1}(A)) \Leftrightarrow \mu(A) = 0$. T is said to be ergodic with respect to μ , or μ is said to be ergodic with respect to T , if and only if $\mu(A) = 0$ or $\mu(X \setminus A) = 0$ whenever the measurable set A is T -invariant, meaning that $T^{-1}(A) = A$. For a nonsingular transformation $T : X \rightarrow X$, the measure μ is said to be conservative with respect to T or T conservative with respect to μ if and only if for every measurable set A with $\mu(A) > 0$,

$$\mu(\{z \in A : \sum_{n=0}^{\infty} 1_A \circ T^n(z) < +\infty\}) = 0.$$

Note that by [1, Proposition 1.2.2], for a nonsingular transformation $T : X \rightarrow X$, μ is ergodic and conservative with respect to T if and only if for any $A \in \mathcal{F}$ with $\mu(A) > 0$,

$$\mu(\{z \in X \mid \sum_{n=0}^{\infty} 1_A \circ T^n(z) < +\infty\}) = 0.$$

Finally, the measure μ is said to be T -invariant, or T is said to preserve the measure μ if and only if $\mu \circ T^{-1} = \mu$. It follows from Birkhoff's Ergodic Theorem that every finite ergodic T -invariant measure μ is conservative, for infinite measures this is not longer true. Finally, two ergodic invariant measures defined on the same σ -algebra are either singular or they coincide up to a multiplicative constant.

Definition 8.1. *Suppose that (X, \mathcal{F}, ν) is a probability space and $T : X \rightarrow X$ is a measurable map such that $T(A) \in \mathcal{F}$ whenever $A \in \mathcal{F}$. The map $T : X \rightarrow X$ is said to be weakly metricaly exact provided that $\limsup_{n \rightarrow \infty} \mu(T^n(A)) = 1$ whenever $A \in \mathcal{F}$ and $\mu(A) > 0$.*

We need the following two facts about weak metrical exactness, the first being straightforward (see the argument in [1, page 15]), the latter more involved (see [26]).

Fact 8.2. *If a nonsingular measurable transformation $T : X \rightarrow X$ of a probability space (X, \mathcal{F}, ν) is weakly metricaly exact, then it is ergodic and conservative.*

Fact 8.3. *A measure-preserving transformation $T : X \rightarrow X$ of a probability space (X, \mathcal{F}, μ) is weakly metricaly exact if and only if it is exact, which means that $\lim_{n \rightarrow \infty} \mu(T^n(A)) = 1$ whenever $A \in \mathcal{F}$ and $\mu(A) > 0$, or equivalently, the σ -algebra $\bigcap_{n \geq 0} T^{-n}(\mathcal{F})$ consists of sets of measure 0 and 1 only. Note that if $T : X \rightarrow X$ is exact, then the Rokhlin's natural extension $(\tilde{T}, \tilde{X}, \tilde{\mu})$ of (T, X, μ) is K -mixing.*

The precise formulation of our main result in this section is the following.

Theorem 8.4. *\tilde{m}_h is a unique h -conformal measure for the map $\tilde{f} : J(\tilde{f}) \rightarrow J(\tilde{f})$. There exists a unique Borel probability \tilde{f} -invariant measure $\tilde{\mu}_h$ on $J(\tilde{f})$ which is absolutely continuous with respect to \tilde{m}_h . The measure $\tilde{\mu}_h$ is metricaly exact and equivalent with \tilde{m}_h .*

The proof of this theorem will consist of several steps. We start with the following.

Lemma 8.5. *Every h -conformal measure ν for $\tilde{f} : J(\tilde{f}) \rightarrow J(\tilde{f})$ is equivalent to \tilde{m}_h .*

Proof. Fix an integer $v \geq 1$ and let

$$J_v = \{(\tau, z) \in J(\tilde{f}) \setminus \text{Sing}(f) : \eta(\tau, z) \geq 1/v\},$$

where $\eta(\tau, z) > 0$ is the number produced in Proposition 5.1. We may assume that $\eta(\tau, z) \leq 1$. Let also $(\hat{\tau}, \hat{z})$ and $(n_j)_1^\infty$ be the objects produced in this proposition. Fix $(\tau, z) \in J_v$. Disregarding finitely many j s we may assume without loss of generality that

$$|f_{\tau|n_j}(z) - \hat{z}| < \frac{1}{4}\eta(\tau, z).$$

Let

$$(8.1) \quad \begin{aligned} B_j(\tau, z) &= [\tau|n_j] \times f_{\tau|n_j, z}^{-1}(B(f_{\tau|n_j}(z), \frac{1}{2}\eta(\tau, z))) \\ \text{and} \\ r_j(\tau, z) &= \frac{1}{2}K\eta(\tau, z)|f'_{\tau|n_j}(z)|^{-1}. \end{aligned}$$

By Koebe's Distortion Theorem and Proposition 5.1 we get that,

$$(8.2) \quad \begin{aligned} \nu(B_j(\tau, z)) &= \nu(\tilde{f}_{\tau|n_j, z}^{-n_j}(\Sigma_u \times B(f_{\tau|n_j}(z), \frac{1}{2}\eta(\tau, z)))) \\ &\geq K^{-h}|f'_{\tau|n_j}(z)|^{-h}\nu(\Sigma_u \times B(f_{\tau|n_j}(z), \frac{1}{2}\eta(\tau, z))) \\ &= K^{-h}|f'_{\tau|n_j}(z)|^{-h}\nu \circ p_2^{-1}(B(f_{\tau|n_j}(z), \frac{1}{2}\eta(\tau, z))) \\ &\geq M_{(2v)^{-1}, \nu}K^{-h}|f'_{\tau|n_j}(z)|^{-h} \\ &\geq M_{(2v)^{-1}, \nu}(2K^{-1}\eta^{-1}(\tau, z))^h r_j^h(\tau, z) \\ &\geq 2^h M_{(2v)^{-1}, \nu}K^{-h}r_j^h(\tau, z), \end{aligned}$$

where $M_{R, \nu} := \inf\{\nu \circ p_2^{-1}(B(w, R)) \mid w \in J(G)\} > 0$. Now fix E , an arbitrary Borel set contained in J_v . Fix also $\varepsilon > 0$. Since the measure ν is regular, by Theorem 3.3 there exists $j(\tau, z) \geq 1$ such that, with $B(\tau, z) = B_{j(\tau, z)}(\tau, z)$ and $r(\tau, z) = r_{j(\tau, z)}(\tau, z)$, we have

$$(8.3) \quad \nu \left(\bigcup_{(\tau, z) \in E} B(\tau, z) \right) \leq \nu(E) + \varepsilon.$$

By the $4r$ -Covering Theorem ([20]), there exists a countable set $\hat{E} \subset E$ such that the balls $\{B(z, r(\tau, z)) : (\tau, z) \in \hat{E}\}$ are mutually disjoint and

$$\bigcup_{(\tau, z) \in \hat{E}} B(z, 4r(\tau, z)) \supset \bigcup_{(\tau, z) \in E} B(z, r(\tau, z)) \supset p_2(E).$$

Hence, by Theorem 7.16 and (8.2), we get

$$\begin{aligned}
\tilde{m}_h(E) &\leq \tilde{m}_h(p_2^{-1}(p_2(E))) \leq \sum_{(\tau,z) \in \hat{E}} \tilde{m}_h \circ p_2^{-1}(B(z, 4r(\tau, z))) \\
&= \sum_{(\tau,z) \in \hat{E}} m_h(B(z, 4r(\tau, z))) \\
(8.4) \quad &\leq C4^h \sum_{(\tau,z) \in \hat{E}} r^h(\tau, z) \\
&\leq C(2K)^h M_{(2v)^{-1}, \nu}^{-1} \sum_{(\tau,z) \in \hat{E}} \nu(B(\tau, z))
\end{aligned}$$

Now, since the sets $\{B(z, r(\tau, z)) : (\tau, z) \in \hat{E}\}$ are mutually disjoint and since $B(\tau, z) \subset p_2^{-1}(B(z, r(\tau, z)))$, so are disjoint the sets $\{B(\tau, z) : (\tau, z) \in \hat{E}\}$. Thus, using (8.3), we get

$$(8.5) \quad \tilde{m}_h(E) \leq C(2K)^h M_{(2v)^{-1}, \nu}^{-1} \nu \left(\bigcup_{(\tau,z) \in \hat{E}} B(\tau, z) \right) \leq C(2K)^h M_{(2v)^{-1}, \nu}^{-1} (\nu(E) + \varepsilon)$$

Letting $\varepsilon \searrow 0$ we thus get

$$\tilde{m}_h(E) \leq C(2K)^h M_{(2v)^{-1}, \nu}^{-1} \nu(E).$$

Consequently $\tilde{m}_h|_{J_v} \prec \nu|_{J_v}$. Since, in virtue of Proposition 5.1, $J(\tilde{f}) \setminus \text{Sing}(\tilde{f}) = \bigcup_{v=1}^{\infty} J_v$, we get that

$$(8.6) \quad \tilde{m}_h|_{J(\tilde{f}) \setminus \text{Sing}(\tilde{f})} \prec \nu|_{J(\tilde{f}) \setminus \text{Sing}(\tilde{f})}.$$

Now, suppose that $\nu(\text{Sing}(\tilde{f})) > 0$. Since \tilde{f}' vanishes on $\text{Crit}(\tilde{f})$, the measure $\nu_0 = (\nu(\text{Sing}(\tilde{f})))^{-1} \nu|_{\text{Sing}(\tilde{f})}$, is h -conformal for $\tilde{f} : J(\tilde{f}) \rightarrow J(\tilde{f})$. But then (8.6) would be true with ν replaced by ν_0 . We would thus have $\tilde{m}_h(J(\tilde{f}) \setminus \text{Sing}(\tilde{f})) = 0$. Since, by Corollary 7.19, $\tilde{m}_h(\text{Sing}(\tilde{f})) = 0$, we would get $\tilde{m}_h(J(\tilde{f})) = 0$. This contradiction shows that $\nu(\text{Sing}(\tilde{f})) = 0$. Consequently,

$$(8.7) \quad \tilde{m}_h \prec \nu.$$

Seeking contradiction, suppose that ν is not absolutely continuous with respect to \tilde{m}_h . Then, there exists a Borel set $X \subset J(\tilde{f}) \setminus \bigcup_{n=0}^{\infty} \tilde{f}^n(\text{Sing}(\tilde{f}))$ such that $\tilde{m}_h(X) = 0$ but $\nu(X) > 0$. But then the measure ν restricted to the forward and backward invariant set $\bigcup_{n,m \in \mathbb{N}} \tilde{f}^{-m}(\tilde{f}^n(X))$ and multiplied by the reciprocal of $\nu(\bigcup_{n,m \in \mathbb{N}} \tilde{f}^{-m}(\tilde{f}^n(X)))$ is h -conformal for $\tilde{f} : J(\tilde{f}) \rightarrow J(\tilde{f})$. But, by conformality of \tilde{m}_h , and as $X \subset J(\tilde{f}) \setminus \bigcup_{n=0}^{\infty} \tilde{f}^n(\text{Sing}(\tilde{f}))$, we conclude from $\tilde{m}_h(X) = 0$ that $\tilde{m}_h(\bigcup_{n,m \in \mathbb{N}} \tilde{f}^{-m}(\tilde{f}^n(X))) = 0$. Since, by (8.7), the measure \tilde{m}_h is absolutely continuous with respect to ν restricted to $\bigcup_{n,m \in \mathbb{N}} \tilde{f}^{-m}(\tilde{f}^n(X))$, we finally get that $\tilde{m}_h(J(\tilde{f})) = 0$. This contradiction show that $\nu \prec \tilde{m}_h$. Together with (8.7) this gives that ν and \tilde{m}_h are equivalent. We are done. \square

Combining inequalities (8.4) and (8.5) (with $\nu = \tilde{m}_h$) from the proof of Lemma 8.5, and letting $\varepsilon \searrow 0$ in (8.5), we get for every Borel set $E \subset J_v$, $v \geq 1$, such that $p_2(E)$ is measurable, that

$$m_h(p_2(E)) \leq C(2K)^h M_{(2v)^{-1}}^{-1} \tilde{m}_h(E).$$

Consequently, as $J(\tilde{f}) \setminus \text{Sing}(\tilde{f}) = \bigcup_{v=1}^{\infty} J_v$ and $\tilde{m}_h(\text{Sing}(\tilde{f})) = 0$, we get the following.

Lemma 8.6. *If E is a Borel subset of $J(\tilde{f})$ such that $p_2(E)$ is measurable and $\tilde{m}_h(E) = 0$, then $m_h(p_2(E)) = 0$. So, by Lemma 8.5, for any h -conformal measure ν for $\tilde{f} : J(\tilde{f}) \rightarrow J(\tilde{f})$, we have that $\nu \circ p_2^{-1}(p_2(E)) = 0$ whenever $\nu(E) = 0$.*

Now, given $(\tau, z) \in J(\tilde{f}) \setminus \text{Sing}(\tilde{f})$, let

$$\mathcal{B}_{(\tau, z)} = \left\{ \left((\tau, z), \overline{B}_j(\tau, z) \right) \right\}_{j=1}^{\infty},$$

where the sets $B_j(\tau, z)$ are defined by formula (8.1). Let

$$\mathcal{B} = \bigcup_{(\tau, z) \in J(\tilde{f}) \setminus \text{Sing}(\tilde{f})} \mathcal{B}_{(\tau, z)}$$

and, following notation from Federer's book [12], let

$$\mathcal{B}_2 := \mathcal{B}(J(\tilde{f}) \setminus \text{Sing}(\tilde{f})) = \{ \overline{B}_j(\tau, z) : (\tau, z) \in J(\tilde{f}) \setminus \text{Sing}(\tilde{f}), j \geq 1 \}.$$

We shall prove the following.

Lemma 8.7. *The family \mathcal{B} is a Vitali relation (in the sense of Federer (see [12], p. 151)) for the measure \tilde{m}_h on the set $J(\tilde{f}) \setminus \text{Sing}(\tilde{f})$.*

Proof. Fix $(\tau, z) \in J(\tilde{f}) \setminus \text{Sing}(\tilde{f})$. Since $p_2(\overline{B}_j(\tau, z)) \subset \overline{B}(z, r_j(\tau, z))$ and since

$$(8.8) \quad \lim_{j \rightarrow \infty} r_j(\tau, z) = 0,$$

we have

$$\lim_{j \rightarrow \infty} \text{diam}(\overline{B}_j(\tau, z)) = 0.$$

This means that the relation \mathcal{B} is fine at the point (τ, z) . Aiming to apply Theorem 2.8.17 from [12], we set

$$\delta((\overline{B}_j(\omega, x))) = r_j(\omega, x)$$

for every $\overline{B}_j(\omega, x) \in \mathcal{B}_2$. Fix $1 < \kappa < +\infty$ (a different notation for $1 < \tau < +\infty$ appearing in Theorem 2.8.17 from [12]). With the notation from page 144 in [12] we have

$$\widehat{\overline{B}}_j(\tau, z) = \bigcup \{ B : B \in \mathcal{B}_2, B \cap \overline{B}_j(\tau, z) \neq \emptyset, \delta(B) \leq \kappa \delta(\overline{B}_j(\tau, z)) \} \subset p_2^{-1} \left((\overline{B}(z, (1 + 2\kappa)r_j(\tau, z))) \right).$$

So, in virtue of Theorem 7.16 and (8.2), we obtain

$$\delta(\overline{B}_j(\tau, z)) + \frac{\tilde{m}_h(\widehat{\overline{B}}_j(\tau, z))}{\tilde{m}_h(\overline{B}_j(\tau, z))} \leq r_j(\tau, z) + \frac{C((1 + 2\kappa)r_j(\tau, z))^h}{C^{-1}r_j^h(\tau, z)} = C^2(1 + 2\kappa)^h + r_j(\tau, z),$$

where $C > 0$ is a constant independent of j . Hence, using (8.8), we get

$$\lim_{j \rightarrow \infty} \left(\delta(\overline{B}_j(\tau, z)) + \frac{\tilde{m}_h(\widehat{B}_j(\tau, z))}{\tilde{m}_h(\overline{B}_j(\tau, z))} \right) \leq C^2(1 + 2\kappa)^h < +\infty.$$

Thus, all the hypothesis of Theorem 2.8.17 in [12], p. 151 are verified and the proof of our lemma is complete. \square

As an immediate consequence of this lemma and Theorem 2.9.11, p. 158 in [12] we get the following.

Proposition 8.8. *For every Borel set $A \subset J(\tilde{f}) \setminus \text{Sing}(\tilde{f})$ let*

$$A_h = \left\{ (\tau, z) \in A : \lim_{j \rightarrow \infty} \frac{\tilde{m}_h(A \cap \overline{B}_j(\tau, z))}{\tilde{m}_h(\overline{B}_j(\tau, z))} = 1 \right\}.$$

Then $\tilde{m}_h(A_h) = \tilde{m}_h(A)$.

Now, we shall prove the following.

Lemma 8.9. *The measure \tilde{m}_h is weakly metrically exact for the map $\tilde{f} : J(\tilde{f}) \rightarrow J(\tilde{f})$. In particular it is ergodic and conservative.*

Proof. Fix a Borel set $F \subset J(\tilde{f}) \setminus \text{Sing}(\tilde{f})$ with $\tilde{m}_h(F) > 0$. By Proposition 8.8 there exists at least one point $(\tau, z) \in F_h$. Our first goal is to show that

$$(8.9) \quad \lim_{j \rightarrow \infty} \frac{\tilde{m}_h(\tilde{f}^{n_j}(F) \cap p_2^{-1}(\overline{B}(f_{\tau|n_j}(z), \eta/2)))}{\tilde{m}_h(p_2^{-1}(\overline{B}(f_{\tau|n_j}(z), \eta/2)))} = 1,$$

where, we recall $\eta = \eta(\tau, z) > 0$ is the number produced in Proposition 5.1 and $(n_j)_1^\infty$ is the corresponding sequence produced there. Indeed, suppose for the contrary that

$$\kappa = \frac{1}{2} \liminf_{j \rightarrow \infty} \frac{\tilde{m}_h(p_2^{-1}(\overline{B}(f_{\tau|n_j}(z), \eta/2)) \setminus \tilde{f}^{n_j}(F))}{\tilde{m}_h(p_2^{-1}(\overline{B}(f_{\tau|n_j}(z), \eta/2)))} > 0.$$

Then, disregarding finitely many n s we may assume that

$$\frac{\tilde{m}_h(p_2^{-1}(\overline{B}(f_{\tau|n_j}(z), \eta/2)) \setminus \tilde{f}^{n_j}(F))}{\tilde{m}_h(p_2^{-1}(\overline{B}(f_{\tau|n_j}(z), \eta/2)))} \geq \kappa > 0$$

for all $j \geq 1$. But

$$\tilde{f}_{\tau|n_j, z}^{-n_j}(p_2^{-1}(\overline{B}(f_{\tau|n_j}(z), \eta/2)) \setminus \tilde{f}^{n_j}(F)) \subset ([\tau|n_j] \times \overline{B}(z, \frac{1}{2}K\eta |f'_{\tau|n_j}(z)|^{-1})) \setminus F = \overline{B}_j(\tau, z) \setminus F$$

and

$$\begin{aligned}
\tilde{m}_h(\tilde{f}_{\tau|n_j, z}^{-n_j}(p_2^{-1}(\overline{B}(f_{\tau|n_j}(z), \eta/2)) \setminus f^{n_j}(F))) &\geq \\
&\geq K^{-h} \left| f'_{\tau|n_j}(z) \right|^{-h} \tilde{m}_h(p_2^{-1}(\overline{B}(f_{\tau|n_j}(z), \eta/2)) \setminus f^{n_j}(F)) \\
&\geq \kappa K^{-h} \left| f'_{\tau|n_j}(z) \right|^{-h} \tilde{m}_h(p_2^{-1}(\overline{B}(f_{\tau|n_j}(z), \eta/2))) \\
&= \kappa K^{-h} \left| f'_{\tau|n_j}(z) \right|^{-h} m_h(\overline{B}(f_{\tau|n_j}(z), \eta/2)) \\
&\geq \kappa K^{-h} M_{\eta/2} \left| f'_{\tau|n_j}(z) \right|^{-h}.
\end{aligned}$$

Hence, making use of Theorem 7.16, we obtain

$$\begin{aligned}
\tilde{m}_h(\overline{B}_j(\tau, z) \setminus F) &\geq \kappa K^{-h} M_{\eta/2} \left| f'_{\tau|n_j}(z) \right|^{-h} \\
&= \kappa (K^2 \eta/2)^{-h} M_{\eta/2} r_j^h(\tau, z) \\
&\geq C^{-1} (K^2 \eta/2)^{-h} M_{\eta/2} \tilde{m}_h(\overline{B}_j(\tau, z)).
\end{aligned}$$

Thus,

$$\frac{\tilde{m}_h(\overline{B}_j(\tau, z) \setminus F)}{\tilde{m}_h(\overline{B}_j(\tau, z))} \geq C^{-1} (K^2 \eta/2)^{-h} M_{\eta/2} > 0.$$

Letting $j \rightarrow \infty$ this contradicts the fact that $(\tau, z) \in F_h$ and finishes the proof of (8.9). Now since $\tilde{f} : J(\tilde{f}) \rightarrow J(\tilde{f})$ is topologically exact, there exists $q \geq 0$ such that $\tilde{f}^q(p_2^{-1}(B(w, \eta/2))) \supset J(\tilde{f})$ for all $w \in J(G)$. It then easily follows from (8.9) and conformality of \tilde{m}_h that

$$\limsup_{k \rightarrow \infty} \tilde{m}_h(\tilde{f}^k(F)) \geq \limsup_{j \rightarrow \infty} \tilde{m}_h(\tilde{f}^{q+n_j}(F)) = 1.$$

Noting also that $\tilde{m}_h(\text{Sing}(\tilde{f})) = 0$ (by Corollary 7.19), the weak metric exactness of \tilde{m}_h is proved. Ergodicity and conservativity follow then from Fact 8.2. We are done. \square

Corollary 8.10. *\tilde{m}_h is the only h -conformal measure on $J(\tilde{f})$ for the map $\tilde{f} : J(\tilde{f}) \rightarrow J(\tilde{f})$.*

Proof. Let ν be an arbitrary h -conformal measure on $J(\tilde{f})$ for the map $\tilde{f} : J(\tilde{f}) \rightarrow J(\tilde{f})$. Since, by Lemma 8.5 the measure ν is absolutely continuous with respect to \tilde{m}_h , it follows from Theorem 2.9.7 in [12], p. 155 and Lemma 8.7 that for \tilde{m}_h -a.e. $(\tau, z) \in J(\tilde{f}) \setminus \text{Sing}(\tilde{f})$,

$$\begin{aligned}
\frac{d\nu}{d\tilde{m}_h}(\tilde{f}(\tau, z)) &= \lim_{j \rightarrow \infty} \frac{\nu(\overline{B}_j(\tilde{f}(\tau, z)))}{\tilde{m}_h(\overline{B}_j(\tilde{f}(\tau, z)))} = \lim_{j \rightarrow \infty} \frac{\nu(\tilde{f}(\overline{B}_j(\tau, z)))}{\tilde{m}_h(\tilde{f}(\overline{B}_j(\tau, z)))} \\
&= \lim_{j \rightarrow \infty} \frac{\int_{\overline{B}_j(\tau, z)} |\tilde{f}'|^h d\nu}{\int_{\overline{B}_j(\tau, z)} |\tilde{f}'|^h d\tilde{m}_h} = \lim_{j \rightarrow \infty} \frac{\nu(\overline{B}_j(\tau, z))}{\tilde{m}_h(\overline{B}_j(\tau, z))} = \frac{d\nu}{d\tilde{m}_h}(\tau, z).
\end{aligned}$$

Since, by Lemma 8.9, the measure \tilde{m}_h is ergodic, it follows that the Radon-Nikodym derivative $\frac{d\nu}{d\tilde{m}_h}$ is \tilde{m}_h -almost everywhere constant. Since ν and \tilde{m}_h are equivalent (by Lemma 8.5) this derivative must be almost everywhere, with respect to \tilde{m}_h as well as ν , equal to 1. Thus $\nu = \tilde{m}_h$ and we are done. \square

In order to prove the existence of a Borel probability \tilde{f} -invariant measure on $J(\tilde{f})$ equivalent to \tilde{m}_h , we will use Marco-Martens method originated in [18]. This means that we shall first produce a σ -finite \tilde{f} -invariant measure equivalent to \tilde{m}_h (this is the Marco-Martens method) and then we will prove this measure to be finite. The heart of the Martens' method is the following theorem which is a generalization of Proposition 2.6 from [18]. It is a generalization in the sense that we do not assume our probability space (X, \mathcal{B}, m) below to be a σ -compact metric space, neither assume we that our map is conservative, instead, we merely assume that item (6) in Definition 8.11 holds. Also, the proof we provide below is based on the concept of Banach limits rather than (see [18]) on the notion of weak limits.

Definition 8.11. *Suppose (X, \mathcal{B}, m) is a probability space. Suppose $T : X \rightarrow X$ is a measurable mapping, such that $T(A) \in \mathcal{B}$ whenever $A \in \mathcal{B}$, and such that the measure m is quasi-invariant with respect to T , meaning that $m \circ T^{-1} \prec m$. Suppose further that there exists a countable family $\{X_n\}_{n=0}^{\infty}$ of subsets of X with the following properties.*

- (1) For all $n \geq 0$, $X_n \in \mathcal{B}$.
- (2) $m(X \setminus \bigcup_{n=0}^{\infty} X_n) = 0$.
- (3) For all $m, n \geq 0$, there exists a $j \geq 0$ such that $m(X_m \cap T^{-j}(X_n)) > 0$.
- (4) For all $j \geq 0$ there exists a $K_j \geq 1$ such that for all $A, B \in \mathcal{B}$ with $A, B \subset X_j$ and for all $n \geq 0$,

$$m(T^{-n}(A))m(B) \leq K_j m(A)m(T^{-n}(B)).$$

- (5) $\sum_{n=0}^{\infty} m(T^{-n}(X_0)) = +\infty$.
- (6) $\lim_{l \rightarrow \infty} m(T(\bigcup_{j=l}^{\infty} Y_j)) = 0$, where $Y_j := X_j \setminus \bigcup_{i < j} X_i$.

Then the map $T : X \rightarrow X$ is called a Marco-Martens map and $\{X_j\}_{j=0}^{\infty}$ is called a Marco-Martens cover.

Remark 8.12. *Note that (6) is satisfied if the map $T : X \rightarrow X$ is finite-to-one. For, if T is finite-to-one, then $\bigcap_{l=1}^{\infty} T(\bigcup_{j=l}^{\infty} Y_j) = \emptyset$.*

Theorem 8.13. *Let (X, \mathcal{B}, m) be a probability space and let $T : X \rightarrow X$ be a Marco-Martens map with a Marco-Martens cover $\{X_j\}_{j=0}^{\infty}$. Then, there exists a σ -finite T -invariant measure μ on X equivalent to m . In addition, $0 < \mu(X_j) < +\infty$ for each $j \geq 0$. The measure μ is constructed in the following way: Let $l_B : l_{\infty} \rightarrow l_{\infty}$ be a Banach limit and let $Y_j := X_j \setminus \bigcup_{i < j} X_i$ for each $j \geq 0$. For each $A \in \mathcal{B}$, set*

$$m_n(A) := \frac{\sum_{k=0}^n m(T^{-k}(A))}{\sum_{k=0}^n m(T^{-k}(X_0))}.$$

If $A \in \mathcal{B}$ and $A \subset Y_j$ with some $j \geq 0$, then we obtain $(m_n(A))_{n=1}^{\infty} \in l_{\infty}$. We set

$$\mu(A) := l_B((m_n(A))_{n=1}^{\infty}).$$

For a general measurable subset $A \subset X$, set

$$\mu(A) := \sum_{j=0}^{\infty} \mu(A \cap Y_j).$$

In addition, if for a measurable subset $A \subset X$, the sequence $(m_n(A))_{n=1}^{\infty}$ is bounded, then we have the following formula.

$$(8.10) \quad \mu(A) = l_B((m_n(A))_{n=1}^{\infty}) - \lim_{l \rightarrow \infty} l_B((m_n(A \cap \bigcup_{j=l}^{\infty} Y_j))_{n=0}^{\infty}).$$

Furthermore, if the transformation $T : X \rightarrow X$ is ergodic (equivalently with respect to the measure m or μ), then the T -invariant measure μ is unique up to a multiplicative constant.

In order to prove Theorem 8.13, we need several lemmas.

Lemma 8.14. *If (Z, \mathcal{F}) is a σ -algebra of sets, $Z = \bigcup_{j=0}^{\infty} Z_j$ is a disjoint union of measurable sets, and for each $j \geq 0$, ν_j is a finite measure on Z_j , then the function $A \mapsto \nu(A) := \sum_{j=0}^{\infty} \nu_j(A \cap Z_j)$, is a σ -finite measure on Z .*

Proof. Let $A \in \mathcal{F}$ and let $(A_n)_{n=1}^{\infty}$ be a partition of A into sets in \mathcal{F} . Then

$$\begin{aligned} \nu(A) &= \sum_{j=0}^{\infty} \nu_j(A \cap Z_j) = \sum_{j=0}^{\infty} \nu_j\left(\bigcup_{n=1}^{\infty} (A_n \cap Z_j)\right) \\ &= \sum_{j=0}^{\infty} \sum_{n=1}^{\infty} \nu_j(A_n \cap Z_j) = \sum_{n=1}^{\infty} \sum_{j=0}^{\infty} \nu_j(A_n \cap Z_j) = \sum_{n=1}^{\infty} \nu(A_n), \end{aligned}$$

where we could have changed the order of summation since all terms involved were non-negative. Thus, we have completed the proof of our lemma. \square

We now suppose that we have the assumption of Theorem 8.13.

Lemma 8.15. *For every $j \geq 0$, the sequence $(m_n(X_j))_{n=1}^{\infty}$ is bounded and $\mu(Y_j) \leq \mu(X_j) < +\infty$.*

Proof. In virtue of (3) of Definition 8.11 there exists a $q \geq 0$ such that $m(X_j \cap T^{-q}(X_0)) > 0$. By (4) of Definition 8.11, we have for all $n \geq 0$ that

$$\begin{aligned} m_n(Y_j) \leq m_n(X_j) &\leq K_j \frac{m(X_j)}{m(X_j \cap T^{-q}(X_0))} m_n(X_j \cap T^{-q}(X_0)) \\ &\leq K_j \frac{m(X_j)}{m(X_j \cap T^{-q}(X_0))} m_{n+q}(X_0) \frac{\sum_{k=0}^{n+q} m(T^{-k}(X_0))}{\sum_{k=0}^n m(T^{-k}(X_0))} \\ &= K_j \frac{m(X_j)}{m(X_j \cap T^{-q}(X_0))} \left(1 + \frac{\sum_{k=n+1}^{n+q} m(T^{-k}(X_0))}{\sum_{k=0}^n m(T^{-k}(X_0))} \right) \\ &\leq K_j \frac{m(X_j)}{m(X_j \cap T^{-q}(X_0))} \left(1 + \frac{q}{\sum_{k=0}^n m(T^{-k}(X_0))} \right). \end{aligned}$$

It follows from (5) of Definition 8.11 that $(m_n(X_j))_{n=1}^\infty \in l_\infty$ and

$$\mu(Y_j) \leq K_j m(X_j) / m(X_j \cap T^{-q}(X_0)) < \infty.$$

Since $X_j = \bigcup_{i=0}^j Y_i$, we are therefore done. \square

Now, for every $j \geq 0$, set $\mu_j := \mu|_{Y_j}$.

Lemma 8.16. *For every $j \geq 0$ such that $\mu(Y_j) > 0$, and for every measurable set $A \subset Y_j$, we have*

$$K_j^{-1} \frac{\mu(Y_j)}{m(Y_j)} m(A) \leq \mu_j(A) \leq K_j \frac{\mu(Y_j)}{m(Y_j)} m(A).$$

Proof. This is an immediate consequence of (4) of Definition 8.11 and the definition of the measure μ . \square

Lemma 8.17. *For any $j \geq 0$, μ_j is a (countably additive) measure on Y_j .*

Proof. Let $j \geq 0$. We may assume without loss of generality that $\mu_j(Y_j) > 0$. Let $A \subset Y_j$ be a measurable set and let $(A_k)_{k=1}^\infty$ be a countable partition of A into measurable sets. For every $n \geq 1$ and for every $l \geq 1$, we have

$$(8.11) \quad \left(\sum_{k=1}^\infty m_n(A_k) \right)_{n=1}^\infty - \sum_{k=1}^l (m_n(A_k))_{n=1}^\infty = \left(\sum_{k=1}^\infty m_n(A_k) \right)_{n=1}^\infty - \left(\sum_{k=1}^l m_n(A_k) \right)_{n=1}^\infty \\ = \left(\sum_{k=l+1}^\infty m_n(A_k) \right)_{n=1}^\infty.$$

It therefore follows from (4) of Definition 8.11 that

$$\left\| \left(\sum_{k=1}^\infty m_n(A_k) \right)_{n=1}^\infty - \sum_{k=1}^l (m_n(A_k))_{n=1}^\infty \right\|_\infty = \left\| \left(\sum_{k=l+1}^\infty m_n(A_k) \right)_{n=1}^\infty \right\|_\infty \\ \leq \left\| \frac{K_j}{m(Y_j)} \left(m_n(Y_j) \sum_{k=l+1}^\infty m(A_k) \right)_{n=1}^\infty \right\|_\infty \\ = \frac{K_j}{m(Y_j)} \left\| \left(m_n(Y_j) \sum_{k=l+1}^\infty m(A_k) \right)_{n=1}^\infty \right\|_\infty.$$

Since, by Lemma 8.15, $(m_n(Y_j))_{n=1}^\infty \in l_\infty$, and since $\lim_{l \rightarrow \infty} \sum_{k=l+1}^\infty m(A_k) = 0$, we conclude that $\lim_{l \rightarrow \infty} \left\| \left(\sum_{k=1}^\infty m_n(A_k) \right)_{n=1}^\infty - \sum_{k=1}^l (m_n(A_k))_{n=1}^\infty \right\|_\infty = 0$. This means that in the Banach space l_∞ , we have $\left(\sum_{k=1}^\infty m_n(A_k) \right)_{n=1}^\infty = \sum_{k=1}^\infty (m_n(A_k))_{n=1}^\infty$. Hence, using continuity of the Banach limit $l_B : l_\infty \rightarrow l_\infty$, we get,

$$\mu(A) = l_B((m_n(A))_{n=1}^\infty) = l_B((m_n(\bigcup_{k=1}^\infty A_k))_{n=1}^\infty) = l_B((\sum_{k=1}^\infty m_n(A_k))_{k=1}^\infty) \\ = \sum_{k=1}^\infty l_B((m_n(A_k))_{n=1}^\infty) = \sum_{k=1}^\infty \mu(A_k).$$

We are done. \square

Combining Lemmas 8.14, 8.15, 8.16, and 8.17, and (3) of Definition 8.11, we get the following.

Lemma 8.18. μ is a σ -finite measure on X equivalent to m . Moreover, $\mu(Y_j) \leq \mu(X_j) < \infty$ and $0 < \mu(X_j)$ for all $j \geq 0$.

Lemma 8.19. The formula 8.10 holds.

Proof. Fix a measurable set $A \subset X$. Then, for every $l \geq 1$ we have that

$$\begin{aligned} l_B((m_n(A))_{n=1}^\infty) &= l_B\left(\sum_{j=0}^l (m_n(A \cap Y_j))_{n=1}^\infty\right) + l_B\left((m_n(\bigcup_{j=l+1}^\infty A \cap Y_j))_{n=1}^\infty\right) \\ &= \sum_{j=0}^l l_B((m_n(A \cap Y_j))_{n=1}^\infty) + l_B\left((m_n(A \cap \bigcup_{j=l+1}^\infty Y_j))_{n=1}^\infty\right). \end{aligned}$$

Hence, letting $l \rightarrow \infty$, we get

$$\begin{aligned} l_B((m_n(A))_{n=1}^\infty) &= \sum_{j=0}^\infty l_B((m_n(A \cap Y_j))_{n=1}^\infty) + \lim_{l \rightarrow \infty} l_B\left((m_n(A \cap \bigcup_{j=l+1}^\infty Y_j))_{n=1}^\infty\right) \\ &= \mu(A) + \lim_{l \rightarrow \infty} l_B\left((m_n(A \cap \bigcup_{j=l}^\infty Y_j))_{n=1}^\infty\right). \end{aligned}$$

We are done. \square

Lemma 8.20. The σ -finite measure μ is T -invariant.

Proof. Let $i \geq 0$ be such that $m(Y_i) > 0$. Fix a measurable set $A \subset Y_i$. Fix $l \geq 1$. We then have

$$\begin{aligned} m_n(T^{-1}(A) \cap \bigcup_{j=l}^\infty Y_j) &= \frac{\sum_{k=0}^n m(T^{-k}(T^{-1}(A) \cap \bigcup_{j=l}^\infty Y_j))}{\sum_{k=0}^n m(T^{-k}(Y_0))} \\ &\leq \frac{\sum_{k=0}^n m(T^{-(k+1)}(A \cap T(\bigcup_{j=l}^\infty Y_j)))}{\sum_{k=0}^n m(T^{-k}(Y_0))} \\ &\leq m_{n+1}(A \cap T(\bigcup_{j=l}^\infty Y_j)) \cdot \frac{\sum_{k=0}^{n+1} m(T^{-k}(Y_0))}{\sum_{k=0}^n m(T^{-k}(Y_0))} \\ &\leq K_i \frac{m_{n+1}(Y_i)}{m(Y_i)} \cdot m(A \cap T(\bigcup_{j=l}^\infty Y_j)) \cdot \frac{\sum_{k=0}^{n+1} m(T^{-k}(Y_0))}{\sum_{k=0}^n m(T^{-k}(Y_0))}, \end{aligned}$$

where the last inequality sign was written because of (4) of Definition 8.11 and since $A \subset Y_i$. Since, the limit when $n \rightarrow \infty$ at last quotient is 1, we get that

$$l_B\left((m_n(T^{-1}(A) \cap \bigcup_{j=l}^\infty Y_j))_{n=1}^\infty\right) \leq \frac{K_i \mu(Y_i)}{m(Y_i)} m(T(\bigcup_{j=l}^\infty Y_j)).$$

Hence, in virtue of (6) of Definition 8.11,

$$\lim_{l \rightarrow \infty} l_B \left((m_n(T^{-1}(A) \cap \bigcup_{j=l}^{\infty} Y_j))_{n=1}^{\infty} \right) \leq \frac{K_i \mu(Y_i)}{m(Y_i)} \lim_{l \rightarrow \infty} m(T(\bigcup_{j=l}^{\infty} Y_j)) = 0.$$

Thus, it follows from Lemma 8.19, and as $A \subset Y_i$, that

$$\mu(T^{-1}(A)) = l_B((m_n(T^{-1}(A)))_{n=1}^{\infty}) = l_B((m_n(A))_{n=1}^{\infty}) = \mu(A).$$

For an arbitrary $A \subset X$, write $A = \bigcup_{j=0}^{\infty} A \cap Y_j$ and observe that

$$\mu(T^{-1}(A)) = \mu\left(\bigcup_{j=0}^{\infty} T^{-1}(A \cap Y_j)\right) = \sum_{j=0}^{\infty} \mu(T^{-1}(A \cap Y_j)) = \sum_{j=0}^{\infty} \mu(A \cap Y_j) = \mu(A).$$

We are done. \square

We now give the proof of Theorem 8.13.

Proof of Theorem 8.13: Combining Lemmas 8.15, 8.18, 8.19, and 8.20, we obtain the statement of Theorem 8.13. We are done. \square

Applying Theorem 8.13 we shall prove Theorem 8.4.

Proof of Theorem 8.4. Since the topological support of \tilde{m}_h is equal to the Julia set $J(\tilde{f})$ and since, by Lemma 7.2, $\text{PCV}(\tilde{f})$ is a nowhere dense subset of $J(\tilde{f})$, we have $\tilde{m}_h(\text{PCV}(\tilde{f})) < 1$. Since the set $\text{PCV}(\tilde{f})$ is forward invariant under \tilde{f} , it follows from ergodicity and conservativity of \tilde{m}_h (see Lemma 8.9) that $\tilde{m}_h(\text{PCV}(\tilde{f})) = 0$. Therefore, in virtue of Lemma 8.6

$$(8.12) \quad \tilde{m}_h(p_2^{-1}(p_2(\text{PCV}(\tilde{f})))) = 0.$$

Now, for every $z \in J(G) \setminus p_2(\text{PCV}(\tilde{f}))$ take $r_z > 0$ such that $J(G) \cap B(z, 2r_z) \subset \hat{\mathbb{C}} \setminus p_2(\text{PCV}(\tilde{f}))$. Since $J(G) \setminus p_2(\text{PCV}(\tilde{f}))$ is a separable metric space, Lindelöf's Theorem yields the existence of a countable set $\{z_j\}_{j=0}^{\infty} \subset J(G) \setminus p_2(\text{PCV}(\tilde{f}))$ such that

$$\bigcup_{j=0}^{\infty} B(z_j, r_{z_j}) \supset J(G) \setminus p_2(\text{PCV}(\tilde{f})).$$

Set

$$A_j := p_2^{-1}(B(z_j, r_j)).$$

Verifying the conditions of Definition 8.11 (with $X = J(\tilde{f})$, $T = \tilde{f}$, $m = \tilde{m}_h$, $X_j = A_j$), \tilde{f} is nonsingular because of Corollary 7.19 and h -conformality of \tilde{m}_h . We immediately see that condition (1) is satisfied, that (2) holds because of (8.12), and that (3) holds because of h -conformality of \tilde{m}_h and topological exactness of the map $\tilde{f} : J(\tilde{f}) \rightarrow J(\tilde{f})$. Condition (5) follows directly from ergodicity and conservativity of the measure \tilde{m}_h . Condition (6) follows since $\tilde{f} : J(\tilde{f}) \rightarrow J(\tilde{f})$ is finite-to-one (see Remark 8.12). Let us prove condition

(4). Fix $j \geq 1$ and two arbitrary Borel sets $A, B \subset A_j$ with $\tilde{m}_h(A), \tilde{m}_h(B) > 0$. Since $B(z_j, 2r_{z_j}) \cap p_2(\text{PCV}(\tilde{f})) = \emptyset$, for all $n \geq 0$ all continuous inverse branches

$$\{\tilde{f}_*^{-n} : p_2^{-1}(B(z_j, 2r_{z_j})) \rightarrow \Sigma_u \times \hat{\mathbb{C}}\}_{* \in I_n}$$

of \tilde{f}^n are well-defined, where $I_n = \{1, \dots, u\}^n$, and because of Koebe's Distortion Theorem and h -conformality of the measure \tilde{m}_h , we have

$$\begin{aligned} \tilde{m}_h \circ \tilde{f}^{-n}(A) &= \tilde{m}_h \left(\bigcup_{* \in I_n} \tilde{f}_*^{-n}(A) \right) = \sum_{* \in I_n} \tilde{m}_h(\tilde{f}_*^{-n}(A)) \\ &\leq \sum_{* \in I_n} K^h |(f_*^{-n})'(\tau, z_j)|^h \tilde{m}_h(A) \\ &= K^{2h} \frac{\tilde{m}_h(A)}{\tilde{m}_h(B)} \sum_{* \in I_n} K^{-h} |(f_*^{-n})'(\tau, z_j)|^h \tilde{m}_h(B) \\ &\leq K^{2h} \frac{\tilde{m}_h(A)}{\tilde{m}_h(B)} \sum_{* \in I_n} \tilde{m}_h(\tilde{f}_*^{-n}(B)) \\ &= K^{2h} \frac{\tilde{m}_h(A)}{\tilde{m}_h(B)} \tilde{m}_h \left(\bigcup_{* \in I_n} \tilde{f}_*^{-n}(B) \right) \\ &= K^{2h} \tilde{m}_h \circ \tilde{f}^{-n}(B) \frac{\tilde{m}_h(A)}{\tilde{m}_h(B)}, \end{aligned}$$

where τ is an arbitrary element of Σ_u . Hence,

$$\frac{\tilde{m}_h \circ \tilde{f}^{-n}(A)}{\tilde{m}_h \circ \tilde{f}^{-n}(B)} \leq K^{2h} \frac{\tilde{m}_h(A)}{\tilde{m}_h(B)},$$

and consequently, condition (4) of Definition 8.11 is satisfied. Therefore, Theorem 8.13 produces a Borel σ -finite \tilde{f} -invariant measure μ on $J(\tilde{f})$, equivalent to \tilde{m}_h .

Now, let us show that the measure μ is finite. Indeed, by Theorem 3.3, there exists a $\delta > 0$ such that for all $g \in G^*$ and for all $x \in J(G)$, every connected component W of $g^{-1}(B(x, \delta))$ satisfies that $\text{diam}(W) < \gamma$ and that W is simply connected. Cover $p_2(\text{PCV}(\tilde{f}))$ with finitely many open balls $\{B(z, \delta) : z \in F\}$, where F is some finite subset of $p_2(\text{PCV}(\tilde{f}))$. for all $j \geq 1$. Since $J(G) \setminus \bigcup_{z \in F} B(z, \delta)$ is covered by finitely many balls $B(z_j, r_{z_j})$, $j \geq 1$, it therefore suffices to show that $\mu(p_2^{-1}(B(z, \delta))) < +\infty$ for all $z \in F$. So, fix $z \in F$. Since $z \in p_2(\text{PCV}(\tilde{f}))$, there thus exists $k \geq 1$ such that $B(z_k, r_{z_k}) \subset B(z, \delta)$. By Lemma 8.15 and the formula (8.10) of Theorem 8.13, it therefore suffices to show that

$$(8.13) \quad \limsup_{n \rightarrow \infty} \frac{\tilde{m}_h(\tilde{f}^{-n}(p_2^{-1}(B(z, \delta))))}{\tilde{m}_h(\tilde{f}^{-n}(A_k))} < +\infty.$$

In order to do this let for every $\tau \in \{1, 2, \dots, s\}^n$, the symbol Γ_τ denote the collection of all connected components of $f_\tau^{-1}(B(z, \delta))$. It follows from Theorem 7.16, Lemma 3.7 and

[37, Corollary 1.9] that for every $V \in \Gamma_\tau$, we have

$$(8.14) \quad \begin{aligned} \tilde{m}_h([\tau] \times V) &\leq m_h(V) \leq C \text{diam}^h(V) \leq C \Gamma^{-h} \left(\frac{\text{diam}(B(z, \delta))}{\text{diam}(B(z_k, r_{z_k}))} \right)^h \text{diam}^h(V_k) \\ &= C(\delta r_{z_k}^{-1} \Gamma^{-1})^h \text{diam}^h(V_k), \end{aligned}$$

where $C > 0$ is a constant independent of n and τ , V_k is a connected component of $f_\tau^{-1}(B(z_k, r_{z_k}))$ contained in V , and Γ is the constant in Lemma 3.7. But, from conformality of the measure \tilde{m}_h and from the fact that $V_k = f_{\tau^*}^{-1}(B(z_k, r_{z_k}))$, where $f_{\tau^*}^{-1} : B(z_k, 2r_{z_k}) \rightarrow \hat{\mathbb{C}}$ is an analytic inverse branch of f_τ , we see that

$$\begin{aligned} \tilde{m}_h([\tau] \times V_k) &\geq K^{-h} |(f_{\tau^*}^{-1})'(z_k)|^h \tilde{m}_h(A_k) \geq K^{-h} \left(K^{-1} \frac{\text{diam}(V_k)}{2r_{z_k}} \right)^h \tilde{m}_h(A_k) \\ &= (2K^2 r_{z_k})^{-h} \text{diam}^h(V_k) \tilde{m}_h(A_k). \end{aligned}$$

Combining this with (8.14) we get that

$$\tilde{m}_h([\tau] \times V) \leq C(2K^2 \delta \Gamma^{-1})^h (\tilde{m}_h(A_k))^{-1} \tilde{m}_h([\tau] \times V_k)$$

Therefore,

$$\begin{aligned} \tilde{m}_h(f^{-n}(p_2^{-1}(B(z, \delta)))) &= \sum_{|\tau|=n} \sum_{V \in \Gamma_\tau} \tilde{m}_h([\tau] \times V) \\ &\leq C(2K^2 \delta \Gamma^{-1})^h (\tilde{m}_h(A_k))^{-1} \sum_{|\tau|=n} \sum_{V \in \Gamma_\tau} \tilde{m}_h([\tau] \times V_k) \\ &\leq C(2K^2 \delta \Gamma^{-1})^h (\tilde{m}_h(A_k))^{-1} \tilde{m}_h(\tilde{f}^{-n}(A_k)). \end{aligned}$$

Thus, the upper limit in (8.13) is bounded above by $C(2K^2 \delta \Gamma^{-1})^h (\tilde{m}_h(A_k))^{-1} < +\infty$, and finiteness of the measure μ is proved.


Dividing μ by $\mu(J(\tilde{f}))$, we may assume without loss of generality that μ is a probability measure. Since for every Borel set $F \subset J(\tilde{f})$ the sequence $(\mu(\tilde{f}^n(F)))_{n=1}^\infty$ is (weakly) increasing, the metric exactness of μ follows from weak metrical exactness of \tilde{m}_h (Lemma 8.9) and the fact that μ and \tilde{m}_h are equivalent. Since, by metrical exactness, μ is ergodic, it is a unique Borel probability measure absolutely continuous with respect to \tilde{m}_h . The proof is complete. \square

9. EXAMPLES

In this section, we give some examples of semi-hyperbolic rational semigroups with nice open set condition.

Example 9.1 ([37, 39]). *Let $f_1(z) = z^2 + 2$, $f_2(z) = z^2 - 2$, and $f = (f_1, f_2)$. Let $G = \langle f_1, f_2 \rangle$. Moreover, let $U := \{z \in \mathbb{C} \mid |z| < 2\}$. Then, G is semi-hyperbolic but not hyperbolic ([37, Example 5.8]). Moreover, G satisfies the nice open set condition with U . Since $J(G) \subset f_1^{-1}(\bar{U}) \cup f_2^{-1}(\bar{U}) \subsetneq \bar{U}$, [39, Theorem 1.25] implies that $J(G)$ is porous and $\text{HD}(J(G)) < 2$. Moreover, by Theorem 1.11, we have $h(f) = \text{HD}(J(G)) = \text{PD}(J(G)) = \text{BD}(J(G))$. Furthermore, $f_1^{-1}(\bar{U}) \cap f_2^{-1}(\bar{U}) \neq \emptyset$. See figure 1.*

FIGURE 1. The Julia set of $\langle f_1, f_2 \rangle$, where $f_1(z) = z^2 + 2$, $f_2(z) = z^2 - 2$.



Proposition 9.2. (See [42]) *Let f_1 be a semi-hyperbolic polynomial with $\deg(f_1) \geq 2$ such that $J(f_1)$ is connected. Let $K(f_1)$ be the filled-in Julia set of f_1 and suppose that $\text{int}K(f_1)$ is not empty. Let $b \in \text{int}K(f_1)$ be a point. Let d be a positive integer such that $d \geq 2$. Suppose that $(\deg(f_1), d) \neq (2, 2)$. Then, there exists a number $c > 0$ such that for each $\lambda \in \{\lambda \in \mathbb{C} : 0 < |\lambda| < c\}$, setting $f_\lambda = (f_{\lambda,1}, f_{\lambda,2}) = (f_1, \lambda(z-b)^d + b)$ and $G_\lambda := \langle f_1, f_{\lambda,2} \rangle$, we have that G_λ is semi-hyperbolic and f_λ satisfies the nice open set condition with an open set U_λ , $J(G_\lambda)$ is porous, $\text{HD}(J(G_\lambda)) = h(f_\lambda) < 2$, and $P(G_\lambda) \setminus \{\infty\}$ is bounded in \mathbb{C} .*

Proof. We will follow the argument in [42]. Conjugating f_1 by a Möbius transformation, we may assume that $b = 0$ and the coefficient of the highest degree term of f_1 is equal to 1.

Let $r > 0$ be a number such that $\overline{B(0, r)} \subset \text{int}K(f_1)$. We set $d_1 := \deg(f_1)$. Let $\alpha > 0$ be a number. Since $d \geq 2$ and $(d, d_1) \neq (2, 2)$, it is easy to see that $(\frac{r}{\alpha})^{\frac{1}{d}} > 2 \left(2(\frac{1}{\alpha})^{\frac{1}{d-1}}\right)^{\frac{1}{d_1}}$ if and only if


$$(9.1) \quad \log \alpha < \frac{d(d-1)d_1}{d+d_1-d_1d} \left(\log 2 - \frac{1}{d_1} \log \frac{1}{2} - \frac{1}{d} \log r \right).$$

We set

$$(9.2) \quad c_0 := \exp \left(\frac{d(d-1)d_1}{d+d_1-d_1d} \left(\log 2 - \frac{1}{d_1} \log \frac{1}{2} - \frac{1}{d} \log r \right) \right) \in (0, \infty).$$

Let $0 < c < c_0$ be a small number and let $\lambda \in \mathbb{C}$ be a number with $0 < |\lambda| < c$. Put $f_{\lambda,2}(z) = \lambda z^d$. Then, we obtain $K(f_{\lambda,2}) = \{z \in \mathbb{C} \mid |z| \leq (\frac{1}{|\lambda|})^{\frac{1}{d-1}}\}$ and

$$f_{\lambda,2}^{-1}(\{z \in \mathbb{C} \mid |z| = r\}) = \{z \in \mathbb{C} \mid |z| = (\frac{r}{|\lambda|})^{\frac{1}{d}}\}.$$

FIGURE 2. The Julia set of $\langle f_1^2, f_2^2 \rangle$, where $f_1(z) = z^2 - 1$, $f_2(z) = z^2/4$.

Let $D_\lambda := \overline{B(0, 2(\frac{1}{|\lambda|})^{\frac{1}{d-1}})}$. Since $f_1(z) = z^{d_1}(1 + o(1))$ ($z \rightarrow \infty$), it follows that if c is small enough, then for any $\lambda \in \mathbb{C}$ with $0 < |\lambda| < c$,

$$f_1^{-1}(D_\lambda) \subset \left\{ z \in \mathbb{C} \mid |z| \leq 2 \left(2 \left(\frac{1}{|\lambda|} \right)^{d-1} \right)^{\frac{1}{d_1}} \right\}.$$

This implies that

$$(9.3) \quad f_1^{-1}(D_\lambda) \subset f_{\lambda,2}^{-1}(\{z \in \mathbb{C} \mid |z| < r\}).$$

Hence, setting $U_\lambda := \text{int}K(f_{\lambda,2}) \setminus K(f_1)$, $f_1^{-1}(U_\lambda) \cup f_{\lambda,2}^{-1}(U_\lambda) \subset U_\lambda$ and $f_1^{-1}(U_\lambda) \cap f_{\lambda,2}^{-1}(U_\lambda) = \emptyset$. Furthermore, since f_1 is semi-hyperbolic, $\hat{\mathbb{C}} \setminus K(f_1)$ is a John domain (see [6]). Hence, U_λ satisfies (osc3). Therefore, G_λ satisfies the nice open set condition with U_λ . We have $J(G_\lambda) \subset \overline{U_\lambda} \subset K(f_{\lambda,2}) \setminus \text{int}K(f_1)$. In particular, $\text{int}K(f_1) \cup (\hat{\mathbb{C}} \setminus K(f_{\lambda,2})) \subset F(G_\lambda)$. Furthermore, (9.3) implies that $f_{\lambda,2}(K(f_1)) \subset \text{int}K(f_1)$. Thus, we have $P(G_\lambda) \setminus \{\infty\} = \cup_{g \in G_\lambda} g(CV^*(f_1) \cup CV^*(f_{\lambda,2})) \subset K(f_1)$, where $CV^*(\cdot)$ denotes the set of all critical values in \mathbb{C} . Hence, $P(G_\lambda) \setminus \{\infty\}$ is bounded in \mathbb{C} . Since f_1 is semi-hyperbolic, there exist an $N \in \mathbb{N}$ and a $\delta_1 > 0$ such that for each $x \in J(f_1)$ and for each $n \in \mathbb{N}$, $\deg(f_1^n : V \rightarrow B(x, \delta_1)) \leq N$ for each connected component V of $f_1^{-n}(B(x, \delta_1))$. Moreover, $f_{\lambda,2}^{-1}(J(f_1)) \cap K(f_1) = \emptyset$ and so $f_{\lambda,2}^{-1}(J(f_1)) \subset \hat{\mathbb{C}} \setminus P(G_\lambda)$. From these arguments, it follows that there exists a $0 < \delta_2 (< \delta_1)$ such that for each $x \in J(f_1)$ and each $g \in G_\lambda$, $\deg(g : V \rightarrow B(x, \delta_2)) \leq N$ for each connected component V of $g^{-1}(B(x, \delta_2))$. Since $P(G_\lambda) \setminus \{\infty\} \subset K(f_1)$ again, we obtain that there exists a $0 < \delta_3 (< \delta_2)$ such that for each $x \in J(G_\lambda)$ and each $g \in G_\lambda$, $\deg(g : V \rightarrow B(x, \delta_3)) \leq N$ for each connected component V of $g^{-1}(B(x, \delta_3))$. Thus, G_λ is semi-hyperbolic. Since $J(G_\lambda) \subset f_1^{-1}(\overline{U_\lambda}) \cup f_{\lambda,2}^{-1}(\overline{U_\lambda}) \subsetneq \overline{U_\lambda}$, [39] implies that $J(G_\lambda)$ is porous and $\text{HD}(J(G_\lambda)) < 2$. Moreover, by Theorem 1.11, we have $h(f_\lambda) = \text{HD}(J(G_\lambda))$. We are done. \square

REFERENCES

- [1] J. Aaronson, *An Introduction to Infinite Ergodic Theory*, Mathematical Surveys and Monographs Vol. 50, American Mathematical Society, 1997.

- [2] R. Brück, *Geometric properties of Julia sets of the composition of polynomials of the form $z^2 + c_n$* , Pacific J. Math., **198** (2001), no. 2, 347–372.
- [3] R. Brück, M. Bürger and S. Reitz, *Random iterations of polynomials of the form $z^2 + c_n$: Connectedness of Julia sets*, Ergod. Th. and Dynam. Sys., **19**, (1999), No.5, 1221–1231.
- [4] M. Bürger, *Self-similarity of Julia sets of the composition of polynomials*, Ergod. Th. and Dynam. Sys. **17** (1997), 1289–1297.
- [5] M. Bürger, *On the composition of polynomials of the form $z^2 + c_n$* , Math. Ann. 310 (1998), no. 4, 661–683.
- [6] L. Carleson, P. W. Jones and J. -C. Yoccoz, *Julia and John*, Bol. Soc. Brazil. Math. **25** (1) 1994, 1-30.
- [7] M. Denker, M. Urbański, *On the existence of conformal measures*, Trans. A.M.S. **328** (1991), 563-587.
- [8] M. Denker, M. Urbański, *Ergodic theory of equilibrium states for rational maps*, Nonlinearity **4** (1991), 103-134.
- [9] M. Denker, M. Urbański, *On Sullivan’s conformal measures for rational maps of the Riemann sphere*, Nonlinearity **4** (1991) 365-384.
- [10] R. Devaney, *An Introduction to Chaotic Dynamical Systems*, Perseus Books, 1989.
- [11] K. Falconer, *Techniques in Fractal Geometry*, John Wiley & Sons, 1997.
- [12] H. Federer, *Geometric Measure Theory*, Springer, 1969.
- [13] J. E. Fornæss and N. Sibony, *Random iterations of rational functions*, Ergod. Th. and Dynam. Sys. **11**(1991), 687–708.
- [14] A. Hinkkanen and G. J. Martin, *The Dynamics of Semigroups of Rational Functions I*, Proc. London Math. Soc. (3) **73**(1996), 358-384.
- [15] A. Hinkkanen and G. J. Martin, *Julia Sets of Rational Semigroups*, Math. Z. **222**, 1996, no.2, 161–169.
- [16] Z. Gong, W. Qiu and Y. Li, *Connectedness of Julia sets for a quadratic random dynamical system*, Ergod. Th. and Dynam. Sys. (2003), **23**, 1807-1815.
- [17] M. Lyubich, *Entropy properties of rational endomorphisms of the Riemann sphere*. Ergod. Th. Dynam. Sys. **3** (1983), 351-386.
- [18] 23. M. Martens, *The existence of σ -finite invariant measures, Applications to real one-dimensional dynamics*. *Front for the Math. ArXiv*, <http://front.math.ucdavis.edu/math.DS/9201300>.
- [19] P. Mattila, *Geometry of sets and measures in Euclidean spaces. Fractals and rectifiability*. Cambridge Studies in Advanced Mathematics, 44. Cambridge University Press, Cambridge, 1995.
- [20] R. D. Mauldin and Szarek, *Graph directed Hilbert space*, Preprint 2008.
- [21] R. D. Mauldin and M. Urbański, *Dimensions and measures in infinite iterated function systems*, Proc. London Math. Soc. (3) **73** (1996) 105-154.
- [22] R. D. Mauldin, M. Urbański, *Graph Directed Markov Systems: Geometry and Dynamics of Limit Sets*, Cambridge Univ. Press (2003)
- [23] J. Milnor, *Dynamics in One Complex Variable (Third Edition)*, Annals of Mathematical Studies, Number 160, Princeton University Press, 2006.
- [24] V. Mayer, B. Skorulski, and M. Urbański, *Random Distance Expanding Mappings, Thermodynamic Formalism, Gibbs Measures, and Fractal Geometry*, Preprint 2008.
- [25] W. Parry, *Entropy And Generators In Ergodic Theory*, Mathematics Lecture Note Series, 1969, Benjamin Inc.
- [26] F. Przytycki and M. Urbański, *Fractals in the Plane - the Ergodic Theory Methods*, to be published from Cambridge University Press, see <http://www.math.unt.edu/~urbanski/>
- [27] D. Ruelle, *Thermodynamic formalism*, Encyclopedia of Math. and Appl., vol. 5, Addison - Wesley, Reading Mass., 1976.
- [28] R. Stankewitz, *Completely invariant Julia sets of polynomial semigroups*, Proc. Amer. Math. Soc., **127**, (1999), No. 10, 2889–2898.

- [29] R. Stankewitz, *Completely invariant sets of normality for rational semigroups*, Complex Variables Theory Appl., Vol 40.(2000), 199–210.
- [30] R. Stankewitz, *Uniformly perfect sets, rational semigroups, Kleinian groups and IFS's*, Proc. Amer. Math. Soc. **128**, (2000), No. 9, 2569–2575.
- [31] R. Stankewitz, T. Sugawa, and H. Sumi, *Some counterexamples in dynamics of rational semigroups*, Annales Academiae Scientiarum Fennicae Mathematica Vol. 29, 2004, 357–366.
- [32] R. Stankewitz and H. Sumi, *Dynamical properties and structure of Julia sets of postcritically bounded polynomial semigroups*, preprint, <http://arxiv.org/abs/0708.3187>
- [33] D. Steinsaltz, *Random logistic maps and Lyapunov exponents*, Indag. Mathem., N. S., 12 (4), 557–584, 2001.
- [34] H. Sumi, *On dynamics of hyperbolic rational semigroups*, J. Math. Kyoto Univ., Vol. 37, No. 4, 1997, 717–733.
- [35] H. Sumi, *On Hausdorff dimension of Julia sets of hyperbolic rational semigroups*, Kodai Mathematical Journal, **21**, (1), 1998, 10-28.
- [36] H. Sumi, *Skew product maps related to finitely generated rational semigroups*, Nonlinearity **13** (2000), 995-1019.
- [37] H. Sumi, *Dynamics of sub-hyperbolic and semi-hyperbolic rational semigroups and skew products*, Ergod. Th. and Dynam. Sys.(2001), **21**, 563-603.
- [38] H. Sumi, *Dimensions of Julia sets of expanding rational semigroups*, Kodai Mathematical Journal, **28**, No.2, 2005, pp390-422.
- [39] H. Sumi, *Semi-hyperbolic fibered rational maps and rational semigroups*, Ergod. Th. and Dynam. Sys. (2006), **26**, 893-922.
- [40] H. Sumi, *Random dynamics of polynomials and devil's-staircase-like functions in the complex plane*, Appl. Math. Comput. 187 (2007), no. 1, 489–500. (Proceedings paper.)
- [41] H. Sumi, *The space of postcritically bounded 2-generator polynomial semigroups with hyperbolicity*, RIMS Kokyuroku 1494, pp 62-86 (Proceedings paper), 2006.
- [42] H. Sumi, *Dynamics of postcritically bounded polynomial semigroups*, preprint 2007, <http://arxiv.org/abs/math/0703591>.
- [43] H. Sumi, *Interaction cohomology of forward or backward self-similar systems*, preprint, <http://arxiv.org/abs/0804.3822>.
- [44] H. Sumi, *Random complex dynamics and semigroups of holomorphic maps*, in preparation.
- [45] H. Sumi and M. Urbański, *The equilibrium states for semigroups of rational maps*, to appear in Monatsh. Math. <http://arxiv.org/abs/0707.2444>.
- [46] H. Sumi and M. Urbański, *Real analyticity of Hausdorff dimension for expanding rational semigroups*, preprint 2007, <http://arxiv.org/abs/0707.2447>.
- [47] M. Urbański, *Rational functions with no recurrent critical points*, Ergod. Th. and Dynam. Sys. 14 (1994), 391-414.
- [48] M. Urbański, *Geometry and ergodic theory of conformal non-recurrent dynamics*, Ergod. Th. and Dynam. Sys. 17 (1997), 1449-1476.
- [49] P. Walters, *An Introduction To Ergodic Theory*, Springer-Verlag, 1982.
- [50] W. Zhou, F. Ren, *The Julia sets of the random iteration of rational functions*, Chinese Sci. Bulletin, **37**(12), 1992, 969-971.