

THE BISHOP-JONES RELATION FOR METRICALLY PROPER ISOMETRIC ACTIONS ON REAL INFINITE-DIMENSIONAL HYPERBOLIC SPACE

TUSHAR DAS, BERND O. STRATMANN, AND MARIUSZ URBAŃSKI

CONTENTS

1. Introduction	1
2. Infinite-dimensional models of hyperbolic geometry	3
3. Classification of isometries	6
4. Discrete Groups of Isometries	7
5. Poincaré Series and Its Exponent of Convergence	9
6. Limit sets	10
7. Convex-cobounded groups of compact type	12
8. The theorem of Bishop and Jones	14
9. Convex-Cobounded Groups	26
10. Acknowledgements	27
References	27

1. INTRODUCTION

Dennis Sullivan, in his IHÉS *Seminar on Conformal and Hyperbolic Geometry* [40] that ran during the late 1970's and early '80s, indicated a possibility¹ of developing the theory of discrete groups acting by hyperbolic isometries on the open unit ball of a separable infinite-dimensional real Hilbert space. Later in the early '90s, Misha Gromov lamented the paucity of results regarding such actions in his seminal lectures *Asymptotic Invariants of Infinite Groups* [19, 6A.III] where he encouraged their investigation in memorable terms: “The spaces like this [infinite-dimensional symmetric spaces] . . . look as cute and sexy to me as their finite dimensional siblings but they have been for years shamefully neglected by geometers and algebraists alike”.

Gromov's lament had not fallen to deaf ears and the geometry and representation theory of infinite-dimensional hyperbolic space \mathbb{H}^∞ and its isometry group have been studied in the

Date: July 23, 2014.

1991 Mathematics Subject Classification. Primary: 37F30, 37F35; Secondary: 30F40, 30F45, 58B20.

Key words and phrases. Fractal geometry, hyperbolic geometry, Hilbert manifold, Poincaré exponent, discrete group, Kleinian group, Fuchsian group, Hilbert space, indefinite metric, Poincaré series, critical exponent, Hausdorff dimension, geodesic dynamics.

The research was supported in part by the NSF Grant DMS 1001874.

¹This was the earliest instance of such a proposal that we could find in the literature. It would be of interest to know whether such an idea might have been discussed prior to that.

last decade by a handful of mathematicians. See, for example, the work by Burger-Iozzi-Monod [3], Delzant-Py [12], and Monod-Py [31]. However, infinite-dimensional hyperbolic space has come into prominence most spectacularly through the recent resolution of a longstanding conjecture in algebraic geometry due to Enriques from the late nineteenth century. Cantat-Lamy [5] proved that the Cremona group (of birational transformations of the complex projective plane) has uncountably many non-isomorphic normal subgroups, i.e. it is not a simple group. Key to their enterprise is the fact, due to Manin [30], that the Cremona group admits a faithful isometric action on an infinite-dimensional hyperbolic space called the Picard-Manin space.

We will be interested in subgroups of $\text{Isom}(\mathbb{H}^\infty)$ whose natural actions are metrically proper, i.e. the orbit of an arbitrary point meets every bounded set in a set of finite cardinality. We call such groups *strongly discrete*. Now by a result of Gromov [6, Theorem 7.4.3] abstract groups that admit such actions correspond to those with the *Haagerup property*². They include amenable groups, Coxeter groups and free groups, and are connected to various lines of investigation within geometric group theory, ergodic theory, representation theory and operator algebras, see [6]. For instance, it is an outstanding problem in geometric group theory to determine whether mapping class groups have the Haagerup property.

To make the connection with subgroups of $\text{Isom}(\mathbb{H}^\infty)$ note that the boundary of infinite-dimensional hyperbolic space is conformally equivalent to Hilbert space $\mathcal{H} := \partial\mathbb{H}^\infty \cup \{\infty\}$. As in finite dimensions, any isometry of \mathcal{H} with respect to the Euclidean metric extends uniquely to an isometry of \mathbb{H}^∞ which fixes ∞ . Therefore there exists a correspondence between parabolic subgroups of the stabilizer $\text{Stab}(\text{Isom}(\mathbb{H}^\infty); \infty)$ and subgroups of $\text{Isom}(\mathcal{H})$ whose orbits are unbounded. However, unlike in finite dimensions, such groups are not necessarily virtually nilpotent. Furthermore, even cyclic subgroups of $\text{Isom}(\mathcal{H})$ are quite different from cyclic subgroups of $\text{Isom}(\mathbb{R}^d)$ for $d \in \mathbb{N}$. Indeed, there is a well-known example of M. Edelstein [13] of a cyclic subgroup of $\text{Isom}(\mathcal{H})$ whose orbits are unbounded but which is not strongly discrete.

This short note describes some of the first investigations regarding the Hausdorff geometry of limit sets of metrically proper isometric actions on real infinite-dimensional hyperbolic space. Our goal is to present a generalization of the Bishop-Jones formula, equating the Poincaré exponent of the underlying group to the Hausdorff dimensions of the uniformly-radial and radial limit sets. To give a dynamical picture of what the Bishop-Jones relation is saying in terms of the geodesic flow on the underlying manifold, let us recall that radial limit points³ correspond to geodesic excursions that return infinitely often to some bounded subset of the manifold, whereas uniformly radial directions correspond to geodesics that never leave a bounded region on the manifold. A priori, there seems to be no reason to believe that the Hausdorff dimensions of these sets are equal and their elegant result significantly generalized a large collection of previously known special cases, see for instance the work of Patterson [34], Sullivan [38] and Dani [8]. Our proof was inspired by Stratmann's

²Such groups are also known as *a-T-menable* groups, since they are *morally* diametrically opposite to groups with Kazhdan's property (T).

³These were introduced in 1936 by Hedlund as *points of approximation*, where he proved that the geodesic flow on compact surfaces of constant negative curvature was topological mixing in [20].

presentation in [37]. Although the original proof and those of various generalizations thereafter (for instance [35, 7, 21]) crucially use the compactness of the sphere at infinity, our proof avoids such a dependence. We hope that it will shed some light on what aspects of this equality are “dimension-free” and follow from the presence of negative curvature. Finally, we indicate the robustness of strongly discrete convex-cobounded groups by showing such groups are finitely generated and of divergence type with finite Poincaré exponent. Further, these groups have compact limit sets, and the Hausdorff and packing measures on the limit sets are finite and positive and coincide with the conformal Patterson measure, up to a multiplicative constant.

The basic ideas behind these results were obtained by the authors during the summer of 2009 at the end of a productive conference, Dynamical Systems II, hosted at the University of North Texas in Denton. These investigations have continued to develop and the reader is encouraged to follow-up this note with the work being done in collaboration with Lior Fishman and David S. Simmons in [17, 10, 11]. The more flexible concept of *partition structures* in these papers generalize the basic mass-redistribution principle that is used in this article. However, the ideas of the proof are more transparent in the setting of this paper and the authors are grateful for the gentle insistence of various colleagues to write such up.

2. INFINITE-DIMENSIONAL MODELS OF HYPERBOLIC GEOMETRY

The infinite-dimensional hyperbolic space is an infinite-dimensional Riemannian manifold, see [29]. Let

$$\mathcal{H} = \ell_2(\mathbb{N}) := \left\{ \mathbf{x} = (x_i)_{i=1}^{\infty} \in \mathbb{R}^{\mathbb{N}} \mid \sum_{i=1}^{\infty} x_i^2 < +\infty \right\}$$

and for $\mathbf{x} \in \mathcal{H}$ let

$$\|\mathbf{x}\| := \left(\sum_{i=1}^{\infty} x_i^2 \right)^{1/2}.$$

Example 2.1. Hilbert space \mathcal{H} can be considered by itself as an infinite-dimensional Riemannian manifold, with at each point the standard inner product

$$\langle \mathbf{u}, \mathbf{v} \rangle_{\mathbf{x}} := \sum_{i=1}^{\infty} u_i v_i.$$

Example 2.2. The space $\widehat{\mathcal{H}} := \mathcal{H} \cup \{\infty\}$ may be given the structure of a Hilbert manifold. The topology on $\widehat{\mathcal{H}}$ is defined as follows: a subset $U \subseteq \widehat{\mathcal{H}}$ is open if and only if $U \cap \mathcal{H}$ is open and if

$$\infty \in U \Rightarrow \mathcal{H} \setminus U \text{ is bounded.}$$

Warning 2.3. The topology on $\widehat{\mathcal{H}}$ is not a one-point compactification. Indeed, $\widehat{\mathcal{H}}$ with the topology defined above is not a compact space, since \mathcal{H} is not locally compact.

2.0.1. *The ball model and the upper half-space model.* There are several models of hyperbolic geometry, which are isometric as infinite-dimensional Riemannian manifolds but which reflect different aspects of hyperbolic geometry. The models we will be interested in are the ball model (\mathbb{B}) and the upper half-space model (\mathbb{H}) and when we do not wish to specify a model we will write \mathbb{H} (for “hyperbolic”). The ball model is the set

$$\mathbb{B} := \{\mathbf{x} \in \mathcal{H} : \|\mathbf{x}\| < 1\}$$

together with the Riemannian metric

$$\langle \mathbf{u}, \mathbf{v} \rangle_{\mathbf{x}, \mathbb{B}} := \frac{4\langle \mathbf{u}, \mathbf{v} \rangle}{(1 - \|\mathbf{x}\|^2)^2}.$$

The upper half-space model is the set

$$\mathbb{H} := \{\mathbf{x} \in \mathcal{H} : x_1 > 0\}$$

together with the Riemannian metric

$$\langle \mathbf{u}, \mathbf{v} \rangle_{\mathbf{x}, \mathbb{H}} := \frac{\langle \mathbf{u}, \mathbf{v} \rangle}{x_1^2}.$$

Remark 2.4. In most references, the $(d + 1)$ -dimensional upper half-space model is defined to be the set $\{\mathbf{x} \in \mathbb{R}^{d+1} : x_{d+1} > 0\}$. When $d = \infty$, this does not make sense since there is no ∞ th coordinate. Thus we have decided to use the first coordinate instead.

Note that the topological boundaries of \mathbb{H} and \mathbb{B} are also Hilbert manifolds (although it requires slightly more work to come up with coordinate charts):

$$\begin{aligned} \partial\mathbb{B} &= \{\mathbf{x} \in \mathcal{H} : \|\mathbf{x}\| = 1\}, \\ \partial\mathbb{H} &= \{\mathbf{x} \in \mathcal{H} : x_1 = 0\} \cup \{\infty\}. \end{aligned}$$

Further note that we have taken the boundary of \mathbb{H} with respect to the Hilbert manifold $\widehat{\mathcal{H}}$ defined in Example 2.2. Henceforth we shall always respect this convention. We note that the closures $\overline{\mathbb{B}}$ and $\overline{\mathbb{H}}$ are not Hilbert manifolds, but they are Hilbert manifolds with boundary (see [29]). We will be content with considering them as topological subspaces of $\widehat{\mathcal{H}}$.

Finally, let us say a word about the geometric significance of \mathbb{B} and \mathbb{H} . The ball model is best if you want to figure out what the world looks like if you are “at a point in X ”; whereas the upper half-space model is best if you are “at a point on the boundary ∂X ”.

2.0.2. *Equivalence of models.* A C^∞ diffeomorphism $\Psi : X \rightarrow Y$ between infinite-dimensional Riemannian manifolds is an *isomorphism* if $\langle \Psi'(x)[u], \Psi'(x)[v] \rangle_{\Psi(x), Y} = \langle u, v \rangle_{x, X}$ for all $x \in X$ and for all $u, v \in T_x X$. Note that every isomorphism of Riemannian manifolds $\Psi : X \rightarrow Y$ is also an *isometry*, i.e.

$$(2.1) \quad d_Y(\Psi(x_1), \Psi(x_2)) = d_X(x_1, x_2) \quad \forall x_1, x_2 \in X.$$

The converse (known in finite dimensions as the Myers-Steenrod theorem) is also true, but much less trivial:

Theorem 2.5 (Theorem 7 of [18]). *Let X and Y be infinite-dimensional Riemannian manifolds, and let $\Psi : X \rightarrow Y$ be a bijection. If Ψ satisfies (2.1), then Ψ is an isomorphism (and in particular is C^∞).*

It may be shown by direct calculation (see [4]) that the map

$$e_{\mathbb{B},\mathbb{H}}(\mathbf{x}) = -\mathbf{e}_1 + 2\frac{\mathbf{x} + \mathbf{e}_1}{\|\mathbf{x} + \mathbf{e}_1\|^2}$$

is an isomorphism of Riemannian manifolds and in particular an isometry. Furthermore, the map $e_{\mathbb{B},\mathbb{H}}$ extends uniquely to a homeomorphism between $\overline{\mathbb{B}}$ and $\overline{\mathbb{H}}$.

2.0.3. Comparison with the classical theory. In contrast with the infinite-dimensional setting of this article, we make a few brief remarks in this subsection about analogous considerations in finite dimensions. In particular about conformal maps, Möbius transformations and the notion of preserving orientation. To define conformal maps we first need the notion of a similarity.

Definition 2.6. Let $T : \mathcal{H} \rightarrow \mathcal{H}$ be a bounded linear operator. T is a *similarity* if it can be written as the product of a positive real number (called the *scaling constant* of T) and a linear isometry. An affine map $A : \mathcal{H} \rightarrow \mathcal{H}$ is a *similarity* if its linear part $\mathbf{x} \mapsto A(\mathbf{x}) - A(\mathbf{0})$ is a similarity. The group of similarities of \mathcal{H} will be denoted $\text{Sim}(\mathcal{H})$.

Definition 2.7. Let X and Y be infinite-dimensional Riemannian manifolds, and let $f : X \rightarrow Y$ be a diffeomorphism. We say that f is *conformal* if for each $x \in X$, $f'(x) : T_x X \rightarrow T_{f(x)} Y$ is a similarity.

As in the finite-dimensional case, the quintessential (non-linear) conformal map is the inversion with respect to a sphere.⁴ The following theorem generalizes the classical result known as Liouville's theorem, which tells us that for $d \geq 3$, any conformal diffeomorphism between two open connected subsets of \mathbb{R}^d is the restriction of a Möbius transformation.

Theorem 2.8 (Liouville's theorem in Hilbert space). *Let $U, V \subseteq \mathcal{H}$ be nonempty open connected sets and let $\phi : U \rightarrow V$ be a conformal diffeomorphism. Then one of the following two cases holds:*

- (NL) ϕ is the composition of an inversion and an affine similarity, or
- (L) ϕ is an affine similarity.

Note that in either case the map ϕ extends uniquely to a conformal map $\widehat{\phi} : \widehat{\mathcal{H}} \rightarrow \widehat{\mathcal{H}}$. As in the finite-dimensional case, the map $\widehat{\phi}$ is called a *Möbius transformation* and we denote the class of such maps by $\text{Mob}(\widehat{\mathcal{H}})$. The nonlinear (NL) case corresponds to the case when ∞ is not preserved, and the linear (L) case corresponds to when ∞ is preserved. Theorem 2.8 follows from the observation (see [23]) that R. Nevanlinna's proof of the finite-dimensional Liouville's theorem [32] extends to infinite dimensions.

⁴ Fix $\mathbf{p} \in \mathcal{H}$ and $\alpha > 0$, and let $S(\mathbf{p}, \alpha)$ denote the sphere around \mathbf{p} of radius α . The *inversion with respect to the sphere $S(\mathbf{p}, \alpha)$* is the map

$$i_{\mathbf{p},\alpha} : \mathbf{x} \mapsto \alpha^2 \frac{\mathbf{x} - \mathbf{p}}{\|\mathbf{x} - \mathbf{p}\|^2} + \mathbf{p}.$$

We make the conventions that $i_{\mathbf{p}} = i_{\mathbf{p},1}$ and $i = i_{\mathbf{0}}$.

Remark 2.9. Notice that if, motivated by the finite-dimensional theory, we restricted to the subclass of $\text{Mob}(\widehat{\mathcal{H}})$ defined by

$$\text{Mob}^*(\widehat{\mathcal{H}}) := \{ g \in \text{Mob}(\widehat{\mathcal{H}}) \mid g \text{ is the composition of finitely many inversions} \},$$

then, unlike the finite-dimensional case, we have

$$\text{Mob}^*(\widehat{\mathcal{H}}) \subsetneq \text{Mob}(\widehat{\mathcal{H}}).$$

In fact, the map g can be written as a finite composition of inversions if and only if $\text{Fix}(g)$ has finite codimension. For example, the shift map on $\ell_2(\mathbb{Z})$ cannot be written as a finite composition of inversions. We give an indication of why this is true. Say $\text{Fix}(g)$ has finite codimension, then one can find a finite-dimensional subspace V such that the entire map is the Poincaré extension of its restriction to V . This reduces us to the finite-dimensional statement [36] that every Möbius map is a composition of finitely many inversions which may then be re-extended. On the other hand if one computes the composition of two inversions it can be shown that $\text{Fix}(g)$ has codimension 1 and so composing finitely many inversions only adds one finitely many times to the codimension.

Remark 2.10. One cannot make sense of *orientation-preserving* transformations in infinite dimensions as one cannot define a meaningful notion of orientation. If one wanted to define orientation-preserving via the kernel of a continuous homomorphism $\mathcal{O} : \mathcal{O}(\mathcal{H}) \rightarrow \mathbb{Z}_2$ one would easily fall into a trap. (Here \mathbb{Z}_2 is the group with two elements.) For example, any reflection in a hyperplane on $\ell_2(\mathbb{Z})$ would be orientation-preserving. For a construction of such a map, take the commutator of the shift map squared and the map that switches consecutive pairs of nonnegative coordinates, i.e. 0 and 1, 2 and 3, etc.

3. CLASSIFICATION OF ISOMETRIES

In [3] one may find the following classification of isometries of \mathbb{H} based on results in [25] and [26]. Every isometry $g \in \text{Isom}(\mathbb{H})$ is exactly one of the following three types: if it has bounded orbits then it is called *elliptic*; if its orbits (one or equivalently all) are unbounded and it has one fixed point on the boundary then it is called *parabolic*, and if its orbits are unbounded and it fixes two points on the boundary it is called *hyperbolic*. We may conjugate each $g \in \text{Isom}(\mathbb{H})$ to a “normal form” whose geometrical significance is clearer. The normal form will depend on the classification of g as elliptic, parabolic, or hyperbolic. We will not prove the remaining propositions in this section, but proofs may be found in [10]. We first must introduce some

Notation 3.1. If G is a group acting on a space X , then for each $x \in X$ we will denote its stabilizer by

$$\text{Stab}(G; x) := \{ g \in G : g(x) = x \}.$$

For any Hilbert space \mathcal{H} , by $\mathcal{O}(\mathcal{H})$ we denote the group of linear isometries of \mathcal{H} . Let us write $\partial\mathbb{H} = \mathcal{E} \cup \{\infty\}$, where $\mathcal{E} := \{ \mathbf{x} \in \mathcal{H} : x_1 = 0 \}$. As in finite dimensions, for any $g \in \text{Sim}(\mathcal{E})$ there exists a unique $\widehat{g} \in \text{Sim}(\mathcal{H})$ so that $\widehat{g} \upharpoonright \mathcal{E} = g$ and so that $\widehat{g}(\mathbb{H}) = \mathbb{H}$. The map \widehat{g} is called the *Poincaré extension* of g .

Proposition 3.2. *Fix $g \in \text{Isom}(\mathbb{H})$.*

- (i) If g is elliptic, then g is conjugate to a map of the form $T \upharpoonright \mathbb{B}$ for some linear isometry $T \in \mathcal{O}(\mathcal{H})$.
- (ii) If g is parabolic, then g is conjugate to a map of the form $\mathbf{x} \mapsto \widehat{T}[\mathbf{x}] + \mathbf{p} : \mathbb{H} \rightarrow \mathbb{H}$, where $T \in \mathcal{O}(\mathcal{E})$ and $\mathbf{p} \in \mathcal{E}$.
- (iii) If g is hyperbolic, then g is conjugate to a map of the form $\mathbf{x} \mapsto \lambda \widehat{T}[\mathbf{x}] : \mathbb{H} \rightarrow \mathbb{H}$, where $0 < \lambda < 1$ and $T \in \mathcal{O}(\mathcal{H})$.

In the first (*elliptic*) case the orbit $(g^n(\mathbf{0}))_1^\infty$ remains fixed forever, and in the third (*hyperbolic*) case it diverges to the boundary. In the latter, there is a pair of fixed points at infinity: one is attracting and the other repelling. In this case, every orbit is unbounded and the forward orbit approaches the attractive fixed point while the backward orbit approaches the repelling fixed point. Further, there exists a unique fixed geodesic connecting the two fixed points that is invariant under the action of g . On the other hand, things can get far more interesting in the second case when g is *parabolic*: then the orbit can oscillate, both accumulating at infinity and returning infinitely often to a bounded region. Note that this is forbidden in finite dimensions. We record this phenomena in the following

Proposition 3.3. *There exists a parabolic $g \in \text{Stab}(\text{Isom}(\mathbb{H}); \infty)$ whose orbit $(g^n(\mathbf{0}))_1^\infty$ is unbounded but returns infinitely often to a bounded region and in fact accumulates at $\mathbf{0}$.*

We remark that this proposition is equivalent to a construction of M. Edelstein⁵, who in Theorem 2.1 of [13] showed that there exists a fixed-point-free isometry $g \in \text{Isom}(\ell_2(\mathbb{C}))$ and a sequence $(n_k)_1^\infty$ so that $g^{n_k}(\mathbf{0}) \xrightarrow[k]{} \mathbf{0}$.

4. DISCRETE GROUPS OF ISOMETRIES

Let $X = \mathbb{H}$ and let G be a subgroup of the isometry group $\text{Isom}(X)$. In finite dimensions, i.e. when $X = \mathbb{H}^n$ the following definitions are equivalent:

- (1) For every bounded $B \subseteq X$, $\#[g \in G : gB \cap B \neq \emptyset] < \infty$.
- (2) For every $x \in X$, there exists an open set $U \ni x$ with

$$gU \cap U \neq \emptyset \Rightarrow gx = x.$$

- (3) G is a discrete subset of $\text{Isom}(X)$ w.r.t. compact-open topology.

Any one of them may be taken as a definition of a *discrete* group of isometries. However in infinite dimensions one must proceed more carefully. Notice that although (1) \Rightarrow (2) even in infinite dimensions, there exist natural examples of groups that show us (2) \nRightarrow (1).

Example 4.1. Consider \mathbb{H}_∞ and let

$$V := \bigcup_{m \geq 2} \{(0, n_2, n_3, \dots, n_m, 0, 0, \dots) : n_i \in \mathbb{Z} \ \forall i = 2, \dots, m\} \subseteq \mathcal{E}$$

Then v is a \mathbb{Z} -vector space and the group

$$G := \langle x \mapsto x + v : v \in V \rangle$$

is an example of one that satisfies (2) but not (1).

⁵See [42] for a recent presentation.

It may be somewhat harder to imagine the right infinite-dimensional analogue of (3). Let us start with the following definitions:

Definition 4.2. The group G is *strongly discrete* if (1) holds, i.e. for every bounded $B \subseteq X$,

$$\#[g \in G : gB \cap B \neq \emptyset] < \infty.$$

The group G is *weakly discrete* if (2) holds, i.e. for every $x \in X$, there exists an open set $U \ni x$ with

$$gU \cap U \neq \emptyset \Rightarrow gx = x.$$

Definition 4.3. An group G acts *properly discontinuously* if for every $x \in X$, there exists an open set $U \ni x$ with

$$gU \cap U \neq \emptyset \Rightarrow g = \text{id}.$$

Equivalently, if there exists $r > 0$ such that

$$B(0, r) \cap \bigcup_{g \in G \setminus \{\text{id}\}} g(B(0, r)) = \emptyset.$$

A group is *torsion-free* if every element of finite order is the identity. A group action is *free* if $\text{Fix}(g) \neq \emptyset \Rightarrow g = \text{id}$.

We remark that unlike strong discreteness which turns out to be a rather robust notion in infinite dimensions, the notion of being properly discontinuous is far more fragile. Now we summarize the connections between our various notions in the following

Observation 4.4. Let (X, d) be a metric space and let $G < \text{Isom}(X)$. Then:

1. Strongly discrete actions are weakly discrete.
2. Torsion-free strongly discrete actions are free.
3. Properly discontinuous actions are weakly discrete and free.
4. Torsion-free strongly discrete actions are properly discontinuous.
5. If X is a CAT(0) space, then free actions are torsion-free.

We make a couple of remarks before proving the observation.

Remark 4.5. Strongly discrete torsion-free groups are always properly discontinuous. In the reverse direction

- In finite dimensions, or when X is proper, properly discontinuous groups are strongly discrete via 3., since $\text{WD} \Rightarrow \text{SD}$ in such a situation.
- On the other hand 5. tells that that in CAT(0) setting properly discontinuous groups are torsion-free.

We also remark that in a CAT(0) setting, if your group has torsion elements then it has no chance of being properly discontinuous. However for properly discontinuous groups:

- (1) The fixed points of elements $g \in G$ do not occur in the interiors of our models of hyperbolic Hilbert space.
- (2) If $g \in G$ has three fixed points, then $g = \text{id}$.

Proof. [Of Observation 4.4]

- Part 1. [SD \Rightarrow WD]
Fix $x \in X$. Then $\varepsilon := \min\{d(x, gx) : gx \neq x\} > 0$ by strong discreteness. So set $U := B(x, \varepsilon/2)$. Then we have that for every g with $gx \neq x$ we have $d(x, gx) \geq \varepsilon$ which implies $gU \cap U = \emptyset$.
- Part 2. [SD + TF \Rightarrow F]
By way of contradiction if G were not free then $gx = x$ for some $g \neq \text{id}$. Then for every n we have $d(g^n x, x) = 0$. Now strong discreteness implies that $\#[g^n : n \in \mathbb{Z}] < \infty$ which in turn produces an N for which $g^N = \text{id}$. This contradicts the assumption that G was torsion-free.
- Part 3. [Properly discontinuous \Leftrightarrow WD + F]
This follows straight from the definitions.
- Part 4. [SD + TF \Rightarrow PD]
Follows from Parts 1., 2. and 3.
- Part 5. [If X is a CAT(0) space, then F \Rightarrow TF.]
Suppose not, then $g^n = \text{id}$ for some $g \neq \text{id}$. Then $\#[H] < \infty$ where $H := \langle g \rangle$. Therefore the orbit $H(0)$ is bounded and by the Cartan's lemma there exists an $x \in \text{Fix}(H) = \text{Fix}(g)$. But we had assumed that G was free.

□

We make the simplifying assumption that our groups are without torsion-elements.

Remark 4.6. A discrete group G is strongly discrete if and only if for every sequence $(g_n)_1^\infty$ of distinct elements of G ,

$$\limsup_{n \rightarrow \infty} d_{\mathcal{H}}(g_n(0), \partial\mathbb{B}) = 0.$$

This may be observed from the following equivalences

$$\limsup_{n \rightarrow \infty} d_{\mathcal{H}}(g_n(0), \partial\mathbb{B}) = 0 \Leftrightarrow \liminf_{n \rightarrow \infty} d_{\mathcal{H}}(g_n(0), 0) = 1 \Leftrightarrow \liminf_{n \rightarrow \infty} d_{\mathbb{B}}(g_n(0), 0) = +\infty.$$

5. POINCARÉ SERIES AND ITS EXPONENT OF CONVERGENCE

Definition 5.1. Fix $x \in \mathbb{K}_\infty$ and $s > 0$. The *Poincaré series* of the group G is defined by

$$(5.1) \quad \Sigma_{x,y}(s) = \sum_{g \in G} e^{-sd_{\mathbb{K}}(x, g(y))}.$$

With this notation, the *Poincaré exponent* of G is given by

$$(5.2) \quad \delta_G = \inf\{s > 0 : \Sigma_{x,y}(s) < +\infty\}.$$

It may be shown by using the triangle inequality that the definition is independent of our choice in x and y .

Observation 5.2. In the finite-dimensional case the Poincaré exponent of a discrete group is always finite. However in the Hilbertian case one may construct a strongly discrete (Schottky) group G such that $\delta_G = +\infty$. Construct a sequence of positive reals (a_n) with $a_n \rightarrow 0$ and $\sum_{n \geq 1} a_n^s = \infty$ for every s . Then with some care one may choose infinitely many generators $(g_n)_n$ such that for every n we have that $e^{-d(g_n(0), 0)} \asymp a_n$ and so that each generator has a distinct coordinate axis as its hyperbolic axis.

However the following fact remains true even in infinite dimensions.

Observation 5.3. If G is a subgroup of $\text{Isom}(\mathbb{K})$ and $\delta_G < +\infty$, then the group G is strongly discrete.

Proof. Suppose that G is not strongly discrete. Then there exists some bounded set W such that $\#W \cap \overline{G(0)} = \infty$. Now the fact that W is bounded implies there exists some $B_{\mathbb{K}}(0, K) \supseteq W$ and therefore for every $g(0) \in W$, we have that

$$\sum_{g(0) \in W} e^{-td(0, g(0))} \geq \sum_{g(0) \in W} e^{-tK} = +\infty.$$

Since t was arbitrary we are done. \square

6. LIMIT SETS

We now will define the limit sets of our group actions and then move on to describe basic properties of such sets. When not specified we may assume that G is a subgroup of $\text{Isom}(\mathbb{H})$ that acts properly discontinuously on \mathbb{H} . In most cases though this is more than we need and it suffices to assume that G is weakly discrete.

Definition 6.1. The limit set of a group $G < \text{Isom}(\mathbb{H})$ is defined to be

$$L(G) := \{\alpha \in \partial\mathbb{H} \mid \exists (g_n)_n \lim_{n \rightarrow \infty} g_n(x) = \alpha\}.$$

This definition is independent of the choice of $x \in \mathbb{H}$ and it is also clear from the definition that $L(G)$ is closed and G -invariant.

Definition 6.2. A group G is called *elementary* whenever $\#L(G) \in \{0, 1, 2\}$.

Theorem 6.3 (No Global Fixed Points). *Any non-elementary weakly discrete group G has no global fixed points.*

Proof. Assume by way of contradiction that we have a global fixed point and then conjugate to send it to ∞ . Then by Liouville's theorem g must be of the form

$$g(x) = \lambda_g T[x] + b.$$

The proof now splits into two cases, viz.

Case 1: [$\lambda_g = 1$ for every $g \in G$.]

Then each hyperplane $\{x : x_0 = \alpha\}$ is fixed by G for every $\alpha > 0$ and therefore $L(G) = \{\infty\}$. This leads to a contradiction since we assumed that G was non-elementary.

Case 2: [There exists a $g \in G$ with $\lambda_g < 1$.]

Without loss of generality we assume that $g(0) = 0$ since g is hyperbolic and so $l_{0, \infty}$ is the invariant axis. Then g is of the form

$$g(x) = g_1(x) = \lambda_1 T_1[x]$$

with $\lambda_1 < 1$. We claim that there exists a $g_2(x) := \lambda_2 T_2[x] + b$ with $\lambda_2 < 1$ and $b \neq 0$. Indeed, since G is non-elementary, there exists a third limit point say ξ that is neither 0 nor ∞ . Now fix a point on the invariant axis at distance greater than one from the boundary $\partial\mathbb{H}_\infty$. Then there exists an element of the group that brings

it arbitrarily close to ξ and this element is the one we were after. Note that since $\xi \notin \{0, \infty\}$ its translation vector is non-zero as claimed.

Now let us calculate the commutator of g_1 and g_2 and call it $g_3 := [g_2, g_1]$. Let

$$y = g_3(x) = g_2^{-1} \circ g_1^{-1} \circ g_2 \circ g_1(x).$$

One may compute the following form

$$y = T_2^{-1} \circ T_1^{-1} \circ T_2 \circ T_1[x] + T_2^{-1} \circ T_1^{-1} \frac{I - \lambda_1 T_1}{\lambda_1 \lambda_2} [b]$$

which we rewrite as $g_3(x) = \hat{T}[x] + \hat{b}$. Note that $\hat{b} \neq 0$ [suppose not, then $T_2^{-1} \circ T_1^{-1} (I - \lambda_1 T_1)[b] = 0 \Leftrightarrow \lambda_1 T_1[b] = b$ which leads to a contradiction since $\lambda_1 < 1$ and $b \neq 0$]. Recall that $\uparrow = e_1$. Now consider

$$\begin{aligned} g_1^n g_3 g_1^{-n}(\uparrow) &= g_1^n g_3(\lambda_1^{-n} T_1^{-n} \uparrow) \\ &= g_1^n (\lambda_1^{-n} \hat{T}[T_1^{-n} \uparrow] + \hat{b}) \\ &= T_1^n \hat{T}[T_1^{-n} \uparrow] + \lambda_1^n T_1^n [\hat{b}] \\ &= \uparrow + \lambda_1^n T_1^n [\hat{b}]. \end{aligned}$$

Note that the last term goes to zero in norm as $n \rightarrow \infty$ since $\lambda_1 < 1$. But this contradicts the fact that G is weakly discrete. □

Definition 6.4. A properly discontinuous group G is said to be of *compact type* when $L(G)$ is compact.

Theorem 6.5. For a properly discontinuous group G acting on \mathbb{B}_∞ , the following are equivalent:

- (1) G is of compact type.
- (2) Every infinite subset of $G(0)$ contains an accumulation point.
- (3) Each sequence $(g_n(0))_{n \in \mathbb{N}}$ with $\lim_{n \rightarrow \infty} \|g_n(0)\| = 1$ has a converging subsequence, which necessarily accumulates at an element in $L(G)$.

Proof. Notice that (2) \Rightarrow (3) is immediate. Let's start by proving (3) \Rightarrow (1). Suppose that $(\xi_n)_n \subseteq L(G)$. Then for every n , there exists a m_n such that $\|g_{m_n}(0) - \xi_n\| \leq 1/n$. Since we have a discontinuous action, orbits must accumulate on the boundary, i.e. $\|g_{m_n}(0)\| \rightarrow 1$ and $n \rightarrow \infty$. By hypothesis, there exists a subsequence $(n_k)_k$ and $\xi \in L(G)$ such that $g_{m_{n_k}}(0) \rightarrow \xi$ as $k \rightarrow \infty$. We now have that

$$\|\xi_{n_k} - \xi\| \leq \|g_{m_{n_k}}(0) - \xi_{n_k}\| + \|g_{m_{n_k}}(0) - \xi\|,$$

where the first term is bounded by $1/n_k$ and therefore it and the second term both vanish as $k \rightarrow \infty$.

Next we show that (1) \Rightarrow (2). Let $(g_n(0))_n$ be our infinite subset of $G(0)$. Then for every n , $g_n \in G$. Now pick two points $\alpha, \beta \in L(G)$ and let $l_{\alpha, \beta}$ be the geodesic between them. Now fix some $z \in l_{\alpha, \beta}$ and pick some common subsequence such that $g_n(\alpha) \rightarrow \gamma$ and $g_n(\beta) \rightarrow \delta$ as $n \rightarrow \infty$. Now one may verify via calculation (or geometric intuition) that for every $\epsilon > 0$ there exists N_ϵ such that for every $n \geq N_\epsilon$, $g_n(l_{\alpha, \beta}) \subseteq B_{\mathcal{H}}(l_{\gamma, \delta}, \epsilon)$. One

may check that the Hausdorff distance between the $g_n(l_{\alpha,\beta})$ is dependent on the distances $\|\gamma - g_n(\alpha)\|$ and $\|\delta - g_n(\beta)\|$ and vanishes as they decrease to zero. Therefore there exists a subsequence such that $g_n(l_{\alpha,\beta}) \subseteq B_{\mathcal{H}}(l_{\alpha,\beta}, 1/n)$ for every n . It now follows that for every n there exists $w_n \in l_{\gamma,\delta}$ such that $d_{\mathcal{H}}(g_n(z), w_n) \leq 1/n$. Now there exists a subsequence $(w_n)_n$ (by compactness of $l_{\gamma,\delta}$) such that $w_n \rightarrow w \in l_{\gamma,\delta}$ as $n \rightarrow \infty$. Finally one can see that $g_n(z) \rightarrow w$ as $n \rightarrow \infty$ since

$$\|g_n(z) - w\| \leq \|g_n(z) - w_n\| + \|w_n - w\|.$$

□

Remark 6.6. This theorem may be extended to non-proper CAT(-1) spaces modulo the construction of geodesics between two boundary points, see [10].

7. CONVEX-COBOUNDED GROUPS OF COMPACT TYPE

Fix $\alpha \in \partial\mathbb{K}_{\infty}$ and let S_{α} refer to the *hyperbolic ray* from 0, the origin, to α . As in finite dimensions the Busemann function may be extended to $\partial\mathbb{K}_{\infty} \times \mathbb{K}_{\infty} \times \mathbb{K}_{\infty}$ via

$$\lim_{z \rightarrow \alpha} \mathcal{B}_z(x, y) =: \mathcal{B}_{\alpha}(x, y),$$

where $\mathcal{B}_z(x, y) := d(x, z) - d(y, z)$.

Definition 7.1. Let us define the following distinguished subsets of $L(G)$:

- $\alpha \in L(G)$ is a *horospherical limit point* of G if there exists a sequence of orbit points $g(0)$ with

$$\mathcal{B}_{\alpha}(0, g(0)) \rightarrow -\infty.$$

The set of such points is called the *horospherical limit set* and is denoted by $L_h(G)$ or by L_h .

- $\alpha \in L(G)$ is a *radial limit point* of G if there exists a constant $c = c(\alpha) > 0$ such that

$$S_{\alpha} \cap B(g(0), c) \neq \emptyset$$

for infinitely many $g(0)$ for every $g \in G$. The set of such points is called the *radial limit set* and is denoted by $L_r(G)$ or by L_r .

- $\alpha \in L(G)$ is a *uniformly radial limit point* of G if there exists a positive constant $c = c(\alpha) > 0$ such that

$$S_{\alpha} \subseteq \cup_{g \in G} B(g(0), c).$$

The set of such points is called the *uniformly radial limit set* and is denoted by $L_{ur}(G)$ or by L_{ur} .

In other words, we may write

$$L_{ur}(G) = \cup_{\sigma > 0} L_{ur,\sigma}(G)$$

and

$$L_r(G) = \cup_{\sigma > 0} L_{r,\sigma}(G)$$

where $\alpha \in L_{ur,\sigma}(G)$ when the geodesic from 0 to α , S_{α} , is covered by hyperbolic balls $B(g(0), \sigma)$ over the G -orbit of the origin and similarly where $\alpha \in L_{r,\sigma}(G)$ when S_{α} intersects

infinitely many hyperbolic balls $B(g_n(0), \sigma)$ for some subsequence of the G -orbit of the origin.

Notice that we have the following inclusions:

$$L_{ur} \subseteq L_r \subseteq L_h \subseteq L.$$

Definition 7.2. We define the projection map $\Pi : X \setminus \{0\} \rightarrow \partial X$ to be the unique map so that for all $x \in X \setminus \{0\}$, x is on the geodesic joining 0 and $\Pi(x)$. For $x \in X$ and $\sigma > 0$, it is useful to consider the set $\Pi(B(x, \sigma))$, which is called the “shadow” of the ball $B(x, \sigma)$, which we may also denote by $\text{Shad}(x, \sigma)$. One may imagine shining a light from the point 0 onto the boundary.

Remark 7.3. In general there is nothing distinguished about the origin 0 and in fact we can similarly define Π_z and Shad_z when we are shining a light from $z \in \mathbb{H}$. Without the subscript we assume that the light is based at the origin. The reader is invited to verify that we would get the same sets $L_{ur}(G)$, $L_r(G)$, and $L_h(G)$ if in our constructions we replaced 0 by an arbitrary but frozen point $z \in \mathbb{H}$.




FIGURE 1. A useful estimate regarding the Euclidean diameter of shadows that will be used in the sequel is $\text{diam}(\text{Shad}(g(0), \sigma)) \asymp \sigma e^{-d(0, g(0))}$, see [37, Lemma 2.1].

Definition 7.4. Given $\alpha, \beta \in \overline{\mathbb{B}}_\infty$, define $[\alpha, \beta]$ and (α, β) respectively to be the unique geodesics joining α and β , the former with the endpoints α and β while the latter without them. Let

$$\mathcal{C} := \mathcal{C}(L(G)) = \bigcup_{\alpha, \beta \in L(G)} [\alpha, \beta]$$

A strongly discrete G is *convex-cobounded* if there exists $R > 0$ such that

$$\mathcal{C} \subseteq G(B(\mathbf{0}, R)).$$

Theorem 7.5. Suppose G is a strongly discrete group of compact type. Then the following are equivalent:

- (A) G is convex-cobounded.


$$(B) L(G) = L_{ur}(G)$$

$$(C) L(G) = L_r(G)$$

$$(D) L(G) = L_h(G)$$

Proof. Since $(B) \Rightarrow (C) \Rightarrow (D)$, it is enough to prove $(A) \Rightarrow (B)$ and $(D) \Rightarrow (A)$.

[Proof of $(A) \Rightarrow (B)$] Fix some $\xi \in L(G)$. In view of Remark 7.3, we may assume without loss of generality that $[\mathbf{0}\xi] \subseteq \mathcal{C}$. Therefore, for every $x \in [\mathbf{0}\xi]$ there exists some $g \in G$ with $d(g(\mathbf{0}), x) \leq R$, with R coming from the definition of convex-coboundedness. Choose $x_n \in [\mathbf{0}\xi]$ with $d(\mathbf{0}, x_n) = n$ and let $g_n \in G$ denote the corresponding elements that move the origin R -close to x_n .



Then notice that $d(g_n(\mathbf{0}), g_{n+1}(\mathbf{0})) \leq 2R + 1$ and so it is clear that $\xi \in L_{ur}(G)$.

[Proof of $(D) \Rightarrow (A)$] By way of contradiction suppose that G is not convex-cobounded. Let D be a Dirichlet domain centered at $\mathbf{0}$, i.e.

$$D := \{x : d(\mathbf{0}, x) < d(\mathbf{0}, g(x)) \quad \forall g \in G \text{ such that } g(\mathbf{0}) \neq \mathbf{0}\}$$

and let $\Delta := D \cap \mathcal{C}$. Fix a point $x \in \mathcal{C} \setminus G(B(\mathbf{0}, R))$ and note that the set of points in the orbit of $\mathbf{0}$ whose distance to x is less than the distance from x to $\mathbf{0}$ is finite, since G is strongly discrete. Pick the element $g(\mathbf{0}) \in G(\mathbf{0})$ which is closest to x and then $g^{-1}(x) \in \Delta \setminus B(\mathbf{0}, R)$. Therefore $\Delta \subsetneq B(\mathbf{0}, R)$ for every $R > 0$. Hence, we may construct a sequence $y_n \in G(x) \cap (\Delta \setminus B(\mathbf{0}, n))$, $n \in \mathbb{N}$. Let ξ_n and η_n be points in $L(G)$ such that $y_n \in [\xi_n, \eta_n]$. Since $\lim_{n \rightarrow \infty} (1 - \|y_n\|_{euc}) = 0$, and the group G has a compact limit set, there exists a convergent subsequence $(y_{n_k})_{k=1}^{\infty}$ to some point $\xi \in L(G)$, and also $\lim_{n \rightarrow \infty} \min\{\|y_n - \xi_n\|_{euc}, \|y_n - \eta_n\|_{euc}\} = 0$. Because of the symmetry of ξ_n and η_n , passing to yet another subsequence, we may assume without loss of generality that

$$\lim_{k \rightarrow \infty} \|y_{n_k} - \xi_{n_k}\|_{euc} = 0$$

and consequently that

$$\lim_{k \rightarrow \infty} \xi_{n_k} = \lim_{k \rightarrow \infty} y_{n_k} = \xi.$$

We now show that such a ξ must belong to $L(G) \setminus L_h(G)$, which will give a contradiction. Indeed, for all $g \in G$ and for every $y_n \in \Delta$ we have that $d(\mathbf{0}, y_n) \leq d(g(\mathbf{0}), y_n)$, which in turn implies that $\mathcal{B}_\xi(g(\mathbf{0}), \mathbf{0}) \geq 0$ and therefore $\xi \notin L_h(G)$. □

8. THE THEOREM OF BISHOP AND JONES

Throughout this section let (X, ρ) be a complete metric space. We start with a version of mass-redistribution principle without any reference to hyperbolic geometry or groups.

Definition 8.1. Given two sets $C, D \subseteq X$ and some $\kappa > 0$ we say that

$$C \subseteq_{\kappa} D \Leftrightarrow B(C, \kappa \text{diam}(C)) \subseteq D,$$

and read it as the κ -thickening of C is contained in D . We denote $B(C, \kappa \text{diam}(C)) := C_{\kappa}$ to be the κ -thickening of C .

We prove the following mass-redistribution result.

Proposition 8.2 (Mass-Redistribution). *Let (X, ρ) be a complete metric space and fix $t \geq 0$ and $\kappa \in (0, 1)$. For every $n \geq 1$ let E_n be a finite set. Set $E_i^j := E_i \times \dots \times E_j$ for $0 \leq i \leq j$ and to avoid clutter we write E^n for E_1^n . Suppose that $\Sigma \subseteq E^* := \bigcup_{n \geq 1} E^n$, has the property that*

$$\widehat{\Sigma} := \{\omega|_{|\omega|-1} \in E^{|\omega|-1} : \omega \in \Sigma\} \subseteq \Sigma \cup \{\emptyset\}.$$

We denote $\Sigma^n := \Sigma \cap E^n$. Suppose further that for every $\omega \in \Sigma$, there exists a closed subset $A(\omega) \subseteq X$ with the following properties:

- (a) For every $\omega \in \Sigma$, $A(\omega) \subseteq_{\kappa} A(\omega|_{|\omega|-1})$;
- (b) For every $\omega \in \Sigma$, $\text{diam}(A(\omega|_{|\omega|-1})) \leq \kappa^{-1} \text{diam}(A(\omega))$;
- (c) For $\omega, \tau \in E^n \cap \Sigma$, $\omega \neq \tau$, $A(\omega) \cap A(\tau) = \emptyset$;
- (d) For every $\omega \in \Sigma$

$$\sum_{\substack{e \in E_{|\omega|+1} \\ \omega e \in \Sigma}} \text{diam}^t(A(\omega e)) \geq \text{diam}^t(A(\omega));$$

and

- (e) $\lim_{n \rightarrow \infty} \max\{\text{diam}(A(\omega)) : \omega \in \Sigma^n\} = 0$.

Then it follows that

$$\text{HD} \left(\bigcap_{n=1}^{\infty} \bigcup_{\omega \in \Sigma^n} A(\omega) \right) \geq t.$$

Remark 8.3. We make a few small remarks, before starting the proof of the Proposition.

- The sets E_i may be thought of as index sets or as alphabets if we think in terms of symbolic dynamics, or IFSs. Note that if we take all of them to be equal to the same set E then E^n is simply the n -fold product of the set E .
- It may help to explain what each of the conditions are saying
 - (1) Condition (a) says that the κ -thickenings are decreasing, while
 - (2) Condition (b) tells us that they do not decrease too fast.
 - (3) Condition (d) is the appropriate redistribution of mass that leads to the appropriate measure in the limit
 - (4) Condition (c) is a natural disjointness condition which is necessary when building a measure.

We note that we may prove the same proposition by specifying rates, i.e. different κ 's for (a) and (b).

- In general metric spaces $\text{diam}(A_{\kappa}) \leq (1 + 2\kappa) \text{diam}(A)$. Whereas for Banach spaces, it turns out that we have equality. Therefore when in Hilbert spaces for instance, we have that (a) implies condition (e).

Proof. Decreasing Σ if necessary we may assume without loss of generality that $A(\omega) \neq \emptyset$ for every $\omega \in \Sigma$. For every $n \geq 1$, let

$$J_n := \bigcup_{\omega \in E^n \cap \Sigma} A(\omega) \quad \text{and} \quad J := \bigcap_{n=1}^{\infty} J_n.$$

Note that for every $\omega \in \Sigma$, we have that $J \cap A(\omega) \neq \emptyset$ and so we fix a point $x_\omega \in J \cap A(\omega)$. Now define inductively the following sequence of Borel probability measures $(\mu_n)_{n \geq 0}$ on J as follows. Let μ_0 be an arbitrary Borel probability measure on J . For the inductive step, suppose that $n \geq 0$ and $\mu_0, \mu_1, \dots, \mu_n$ (each a Borel probability measure on J) have already been defined. Then set

$$(8.1) \quad \mu_{n+1} := \sum_{\omega \in E^{n+1} \cap \Sigma} \frac{\text{diam}^t(A(\omega))}{\sum_{e \in E: \omega|_n e \in \Sigma} \text{diam}^t(A(\omega|_n e))} \mu_n(A(\omega|_n)) \delta_{x_\omega}.$$

Now since μ_n is a probability measure and by (c) all the sets $A(\tau), \tau \in E^n \cap \Sigma$ are pairwise disjoint, a straightforward direct computation shows that μ_{n+1} is a Borel probability measure. Now since $x_\omega \in J$ for every $\omega \in \Sigma$, we immediately get from (8.1) that

$$(8.2) \quad \mu_n(J) = 1 \quad \text{for every } n \geq 0.$$

Now suppose that $\tau \in E^n \cap \Sigma$. In view of (8.1) and (c) we then get

$$(8.3) \quad \begin{aligned} \mu_{n+1}(A(\tau)) &= \sum_{\omega \in E^{n+1} \cap \Sigma} \frac{\text{diam}^t(A(\omega))}{\sum_{e \in E: \omega|_n e \in \Sigma} \text{diam}^t(A(\omega|_n e))} \mu_n(A(\omega|_n)) \delta_{x_\omega}(A(\tau)) \\ &= \sum_{a \in E: \tau a \in \Sigma} \frac{\text{diam}^t(A(\tau a))}{\sum_{e \in E: \tau e \in \Sigma} \text{diam}^t(A(\tau e))} \mu_n(A(\tau)) \\ &= \mu_n(A(\tau)) \sum_{a \in E: \tau a \in \Sigma} \frac{\text{diam}^t(A(\tau a))}{\sum_{e \in E: \tau e \in \Sigma} \text{diam}^t(A(\tau e))} \\ &= \mu_n(A(\tau)) \frac{\sum_{a \in E: \tau a \in \Sigma} \text{diam}^t(A(\tau a))}{\sum_{e \in E: \tau e \in \Sigma} \text{diam}^t(A(\tau e))} \\ &= \mu_n(A(\tau)). \end{aligned}$$

Because of (c) and since $x_\omega \in J \cap A(\omega)$ for every $\omega \in \Sigma$, we get that for every $n \geq 0$ and for all $0 \leq i \leq j$ that

$$(8.4) \quad \mu_{n+j}(A(\tau)) = \mu_{n+j} \left(\bigcup_{\omega \in E_{n+1}^{n+i}: \tau \omega \in \Sigma} A(\tau \omega) \right) = \sum_{\omega \in E_{n+1}^{n+i}: \tau \omega \in \Sigma} \mu_{n+j}(A(\tau \omega)).$$

Having this formula, we now prove the following.

Observation 8.4. $\mu_{n+k}(A(\tau)) = \mu_n(A(\tau))$ for every $n, k \geq 0$ and all $\tau \in E^n \cap \Sigma$.

Proof. Fix $n \geq 0$. The proof now follows by induction on $k \geq 0$. For $k = 0$ it is a tautology. So suppose the claim is true for some $k \geq 0$ and all $\tau \in E^n \cap \Sigma$. Using (8.4) and (8.3) we

then get

$$\begin{aligned}
\mu_{n+(k+1)}(A(\tau)) &= \sum_{\omega \in E^k: \tau\omega \in \Sigma} \mu_{n+(k+1)}(A(\tau\omega)) = \sum_{\omega \in E^k: \tau\omega \in \Sigma} \mu_{(n+k)+1}(A(\tau\omega)) \\
&= \sum_{\omega \in E^k: \tau\omega \in \Sigma} \mu_{n+k}(A(\tau\omega)) \\
&= \mu_{n+k}(A(\tau)) \\
&= \mu_n(A(\tau)).
\end{aligned}$$

The inductive proof is complete. \square

Now let $c := [\sum_{e \in E_1} \text{diam}^t(A(e))]^{-1} < \infty$. Next we prove the following.

Observation 8.5. For every $n \geq 1$ and for every $\omega \in E^n \cap \Sigma$, we have that $\mu_n(A(\omega)) \leq c \text{diam}^t(A(\omega))$.

Proof. We prove this by induction on $n \geq 1$. If $e \in E \cap \Sigma$, then it follows from (8.1) that

$$\begin{aligned}
\mu_1(A(e)) &= \left[\sum_{a \in E_1} \text{diam}^t(A(a)) \right]^{-1} \text{diam}^t(A(e)) \cdot \mu_0(X) \\
&= c \cdot \text{diam}^t(A(e)).
\end{aligned}$$

So suppose that the claim holds for some $n \geq 1$. For every $\omega \in E^{n+1} \cap \Sigma$, we then have from (8.3) and (d) that

$$\begin{aligned}
\mu_{n+1}(A(\omega)) &= \left[\sum_{e \in E_{n+1}: \omega|_n e \in \Sigma} \text{diam}^t(A(\omega|_n e)) \right]^{-1} \text{diam}^t(A(\omega)) \cdot \mu_n(A(\omega|_n)) \\
&\leq [\text{diam}^t(A(\omega|_n))]^{-1} \text{diam}^t(A(\omega)) \cdot \mu_n(A(\omega|_n)) \\
&\leq c \cdot \text{diam}^t(A(\omega)).
\end{aligned}$$

We are done. \square

Now the set $J = \bigcap_{n \geq 1} \bigcup_{\omega \in E^n \cap \Sigma} A(\omega) =: \bigcap_{n \geq 1} J_n$ is closed as it is the intersection of closed sets J_n . Recall that (e) gives us that

$$\lim_{n \rightarrow \infty} \max\{\text{diam}(A(\omega)) : \omega \in E^n \cap \Sigma\} = 0,$$

and now because of the finiteness of the index sets E_n , the set J is totally bounded. Thus J is compact since X is complete. Therefore by the Banach-Alaoglu theorem, the sequence $(\mu_n)_1^\infty$ of Borel probability measures on J contains a weakly convergent subsequence. Denote its weak limit by μ . Since $A(\omega) \cap J$ is a clopen subset of J (with respect to the topology relative to J) for every $\omega \in \Sigma$, it follows from Observation 8.5 that for every $\omega \in \Sigma$,

$$(8.5) \quad \mu(A(\omega)) \leq c \cdot \text{diam}^t(A(\omega)).$$

Note that we have not yet used conditions (a) and (b) and we now do so in estimating from above the measures of balls centered at the points of J .

Let $z \in J = \bigcap_{n \geq 1} J_n$. Since for every $n \geq 1$, the sets $E_n \cap \Sigma$ are finite, it follows from König's Lemma that there exists $\omega \in E^{\mathbb{N}}$ such that $\omega|_n \in \Sigma$ for every $n \geq 1$ and $\{z\} = \bigcap_{n=1}^{\infty} A(\omega|_n)$. Because of (c) this $\omega \in E^{\mathbb{N}}$ is unique.

Now fix a radius $r \in (0, \kappa \min\{\text{diam}(A(\omega)) : \omega \in E^2 \cap \Sigma\})$. Then there exists a largest $n = n(\omega, r) \geq 2$ such that

$$(8.6) \quad r \leq \kappa \text{diam}(A(\omega|_n)).$$

Since $z \in A(\omega|_n)$ it follows from (a) that

$$B(z, r) \subseteq B(z, \kappa \text{diam}(A(\omega|_n))) \subseteq B(A(\omega|_n), \kappa \text{diam}(A(\omega|_n))) \subseteq A(\omega|_{n-1}).$$

Now (8.5) implies that

$$(8.7) \quad \mu(B(z, r)) \leq c \cdot \text{diam}^t(A(\omega|_{n-1})).$$

By the definition of n , we have that $\kappa \text{diam}(A(\omega|_{n+1})) < r$ and then applying (b) twice we get that

$$\text{diam}(A(\omega|_{n-1})) \leq \kappa^{-1} \text{diam}(A(\omega|_n)) \leq \kappa^{-2} \text{diam}(A(\omega|_{n+1})) \leq \kappa^{-3} r.$$

Inserting this into (8.7) we finally get

$$(8.8) \quad \mu(B(z, r)) \leq c \kappa^{-3t} r^t.$$

Note that $\mu(J) = 1$ (since its a probability measure on J) and thus by a direct application of Frostmann's Lemma, we have that $\text{HD}(J) \geq t$. We are done. \square

Remark 8.6. In the case when (X, ρ) is a finite-dimensional Euclidean space, condition (a) can be replaced by the requirement that all the sets $A(\omega)$ for $\omega \in \Sigma$ are uniformly undistorted balls and then a standard volume argument would work to get (8.8). Our Proposition 8.2 requires no extra structure on X and condition (a) will be proved to be satisfied in the course of our proof of the Bishop-Jones theorem for Hilbert spaces. We should note however that condition (a) is somewhat strong and for example it fails in the standard construction of C , the middle-third Cantor set if $X = [0, 1]$. Note however if $X = C$, then (a) is satisfied. Therefore one must take some care in the choice of X .

Notation 8.7. For $\xi \in \partial \mathbb{B}_{\infty}$, we denote $\widehat{B}(\xi, r)$ to be the union of all geodesics (in \mathbb{B}) with both endpoints in the ball in $\partial \mathbb{B}_{\infty}$ centered at ξ and with spherical radius equal to r . In particular $\Pi[\widehat{B}(\xi, r)]$ is equal to this ball.

Next we shall prove the following

Lemma 8.8. *If G is a non-elementary strongly discrete group acting on \mathbb{B} , then for every $t < \delta_G$ there exist two distinct points $\xi_1, \xi_2 \in L(G)$ such that for all $r > 0$,*

$$\sum_{\gamma(0) \in \widehat{B}(\xi_i, r)} e^{-t d(\gamma(0), 0)} = +\infty \quad \text{for } i = 1, 2.$$

We call such points ξ_i , t -divergent points of $L(G)$.

Proof. We assume that our group is non-elementary and so there exist at least two distinct hyperbolic elements. We prove the existence of one such point (as claimed in the Lemma) and the same argument will provide another point distinct from the first; as we have that these hyperbolic elements have distinct pairs of fixed points.

Suppose by way of contradiction, that there exist no t -divergent points in the limit set. Now pick a hyperbolic element g with axis l_g whose attracting and repelling endpoints respectively are ξ_g^+ and ξ_g^- on $L(G)$. Let's look at ξ_g^- and refer to it as simply ξ . Then there exists an r_ξ such that the sum over the G -orbit of 0 within $A := \widehat{B}(\xi, r_\xi)$ is finite, i.e.

$$\sum_{\gamma(0) \in A} e^{-td(\gamma(0), 0)} < +\infty .$$

Then we have that

$$\sum_{\gamma(0) \in \mathbb{B} \setminus A} e^{-td(\gamma(0), 0)} = +\infty .$$

Now for an arbitrary $\varepsilon > 0$, there exists $n \geq 0$ large enough such that $g^n(\mathbb{B} \setminus A) \subseteq \widehat{B}(\xi_g^+, \varepsilon)$. It is enough to show that for such n , we have that

$$\sum_{\gamma(0) \in \mathbb{B} \setminus A} e^{-td(g^n \gamma(0), 0)} = +\infty ,$$

since then we would have shown ξ_g^+ to be t -divergent and thus derived a contradiction. Notice that

$$\begin{aligned} \sum_{\gamma(0) \in \mathbb{B} \setminus A} e^{-td(g^n \gamma(0), 0)} &\geq \sum_{\gamma(0) \in \mathbb{B} \setminus A} e^{-td(0, g^n(0))} e^{-td(g^n(0), g^n \gamma(0))} \\ &\quad \text{(by the triangle inequality)} \\ &= e^{-td(0, g^n(0))} \sum_{\gamma(0) \in \mathbb{B} \setminus A} e^{-td(g^n(0), g^n \gamma(0))} \\ &= e^{-td(0, g^n(0))} \sum_{\gamma(0) \in \mathbb{B} \setminus A} e^{-td(0, \gamma(0))} \\ &\quad \text{[since } g \text{ is an isometry]} \\ &= +\infty . \end{aligned}$$

We are done. □

Definition 8.9. Fix $\tau > 0$. For every integer $n \geq 0$ let

$$A_n(\tau) := \{z \in G(0) : \tau n \leq d(z, 0) \leq \tau(n+1)\}.$$

The set $A_n(\tau)$ is called the *hyperbolic* (n, G) -annulus centered at 0 and of width τ .

We now prove the following lemma

Lemma 8.10. *Let G be a strongly discrete group. Fix $\tau > 0$ and $0 \leq s < t < \delta_G$. Let $\xi \in L(G)$ be a t -divergent point. Then for every $M > 0$ and for every $r > 0$ there exists*

$(n_j(\xi))_{j=1}^\infty$ an increasing sequence of positive integers such that

$$\sum_{g(0) \in \widehat{B}(\xi, r) \cap A_{n_j(\xi)}(\tau)} e^{-sd(g(0), 0)} \geq M \text{ for every } j \geq 1.$$

Proof. Suppose by way of contradiction that there exist $M, r > 0$ and an integer $q \geq 0$ such that

$$\sum_{z \in \widehat{B}(\xi, r) \cap A_n(\tau)} e^{-sd(z, 0)} < M \text{ for every } n \geq q + 1.$$

Take $r_* \in (0, r]$ so small that $A_q(\tau) \cap \widehat{B}(\xi, r_*) = \emptyset$. Then $A_n(\tau) \cap \widehat{B}(\xi, r_*) = \emptyset$ for all $n = 0, 1, 2, \dots, q$ and thus we get that

$$\sum_{z \in \widehat{B}(\xi, r_*) \cap A_n(\tau)} e^{-sd(z, 0)} = 0 \leq M \text{ if } n \leq q$$

and that

$$\sum_{z \in \widehat{B}(\xi, r_*) \cap A_n(\tau)} e^{-sd(z, 0)} \leq \sum_{z \in \widehat{B}(\xi, r) \cap A_n(\tau)} e^{-sd(z, 0)} \leq M \text{ if } n \geq q + 1.$$

Therefore we have that $\sum_{z \in \widehat{B}(\xi, r_*) \cap A_n(\tau)} e^{-sd(z, 0)} \leq M$ for all $n \geq 0$. Hence

$$\begin{aligned} \sum_{z \in G(0) \cap \widehat{B}(\xi, r_*)} e^{-td(z, 0)} &= \sum_{n=0}^{\infty} \sum_{z \in \widehat{B}(\xi, r_*) \cap A_n(\tau)} e^{-(t-s)d(z, 0)} e^{-sd(z, 0)} \\ &\leq \sum_{n=0}^{\infty} \sum_{z \in \widehat{B}(\xi, r_*) \cap A_n(\tau)} e^{-(t-s)\tau n} e^{-sd(z, 0)} \\ &= \sum_{n=0}^{\infty} e^{-(t-s)\tau n} \sum_{z \in \widehat{B}(\xi, r_*) \cap A_n(\tau)} e^{-sd(z, 0)} \\ &\leq M \sum_{n=0}^{\infty} e^{-(t-s)\tau n} \\ &< +\infty. \end{aligned}$$

The last inequality follows since s was chosen strictly smaller than t and so we have a geometric series that converges. But then this contradicts the hypothesis that ξ is a t -divergent point and finishes the proof. \square

We sometimes, for emphasis, will use the shorthand B_e, d_e and similarly B_h, d_h to distinguish between the Euclidean/Hilbertian and hyperbolic settings respectively. In general the absence of subscripts refers to the hyperbolic setting, though diam without a subscript will refer to Euclidean diameter.

Lemma 8.11. *There exists $\alpha > 0$ such that for all $\sigma > \log 2$ we have*

$$\Pi[B_h(z, \alpha\sigma)] \subseteq B_e\left(\Pi[z], \frac{1}{8} \text{diam}_e \Pi[B_h(z, \sigma)]\right),$$

for every z with $d(z, 0) > \sigma$.

Proof. The proof immediately follows from the estimate

$$\text{diam}_e(\Pi[B(z, \sigma)]) = \text{diam}_e(\text{Shad}(z, \sigma)) \asymp \sigma e^{-d(z, g(z))}$$

described in Figure 7.2 above and [37, Lemma 2.1]. \square

For every $\sigma > 0$, let $r_\sigma > 0$ be chosen so small that

$$(8.9) \quad r_\sigma < \pi e^{-\alpha_\sigma \sigma}.$$

From this point on, for the remainder of the section, fix an arbitrary $\tau > 0$. The main ingredient, forming the inductive step in our proof of the Bishop-Jones Theorem, is the following lemma, whose proof is illustrated on the Figure 2 below and is provided after formulation of the lemma.




FIGURE 2. The strategy for the proof of Lemma 8.12 is to construct a collection of “children” of the point $g(0)$. We “pull back” the entire picture via g^{-1} . In the pulled-back picture, with the help of the Light Cone Lemma (cf. [37, Lemma 2.3]) we obtain the existence of many points $x \in G(0)$ such that $\text{Shad}_{g^{-1}(0)}(x, \sigma) \subseteq \text{Shad}_{g^{-1}(0)}(0, \sigma)$. These children can then be pushed forward via g to get children of $g(0)$.

Lemma 8.12. *Fix $0 < s < t < \delta_G$. Let $\xi_1, \xi_2 \in L(G)$ be two t -divergent points (see Lemma 8.8 for their existence). Then there exist $\sigma > 0$ and positive integers $l_1, l_2 \geq 1$ such that the following holds:*

For every $g \in G$ with $d(g(0), 0) > \sigma$, there exists a set $\Gamma(g)$ contained in one of the sets

$$\Gamma_i := \{h \in G : h(0) \in \widehat{B}(\xi_i, r_\sigma) \cap A_{l_i}(\tau)\} \quad \text{for } i = 1, 2;$$

such that the following hold:-

- (a) *The family $\{\Pi[B_h(gh(0), \sigma)] : h \in \Gamma(g)\}$ consists of mutually disjoint balls.*

(b) For every $h \in \Gamma(g)$,

$$\Pi[gh(0)] \in B_e\left(\Pi[g(0)], \frac{1}{8}\text{diam}_e\Pi[B_h(g(0), \sigma)]\right).$$

(c) There exists a constant $\beta_\sigma \in (0, 1)$ depending only on l_1, l_2 and σ (in particular independent of g) such that for every $h \in \Gamma(g)$,

$$\frac{\beta_\sigma}{4}\text{diam}\left(\Pi[B_h(g(0), \sigma)]\right) \leq \text{diam}\left(\Pi[B_h(gh(0), \sigma)]\right) \leq \frac{1}{4}\text{diam}\left(\Pi[B_h(g(0), \sigma)]\right).$$

(d) The following inequality holds

$$\sum_{h \in \Gamma(g)} \text{diam}^s\left(\Pi[B_h(gh(0), \sigma)]\right) \geq \text{diam}^s\left(\Pi[B_h(g(0), \sigma)]\right).$$

Proof. Take $\sigma > \log 2$ and so large as needed for Lemma 8.11 and so that

$$d_e(\xi_1, \xi_2) > 6\pi e^{-\alpha_\sigma \sigma}.$$

Now by the choice of r_σ , see (8.9), we have that

$$\inf\{d_e(x, y) : x \in \widehat{B}(\xi_1, r_\sigma), y \in \widehat{B}(\xi_2, r_\sigma)\} > 4\pi e^{-\alpha_\sigma \sigma}.$$

Since $\alpha_\sigma \sigma > \log 2$, it then follows from the Light Cone Lemma (Lemma 2.3 in [37]) that at least one of the balls $\widehat{B}(\xi_1, r_\sigma)$ or $\widehat{B}(\xi_2, r_\sigma)$ is contained in $g^{-1}(\Pi[B(g(0), \alpha_\sigma \sigma)])$. Assume without loss of generality that $\widehat{B}(\xi_1, r_\sigma) \subseteq g^{-1}(\Pi[B(g(0), \alpha_\sigma \sigma)])$. Consequently, if we fix two integers $l_1, l_2 \geq 1$ that we will specify later in the course of the proof, then

$$\Pi[h(0)] \in g^{-1}(\Pi[B_h(g(0), \alpha_\sigma \sigma)])$$

for every $h \in \Gamma_1$. Therefore $g(\Pi[h(0)]) \in \Pi[B(g(0), \alpha_\sigma \sigma)]$ and g maps the geodesic from 0 to $\Pi[h(0)]$ with $h(0)$ on it to the geodesic from $g(0)$ to $g\Pi[h(0)]$ with $gh(0)$ on it. Therefore $gh(0)$ lies inside of the light cone generated by 0 and $B_h(g(0), \alpha_\sigma \sigma)$ and we thus have that

$$\Pi[gh(0)] \in \Pi[B_h(g(0), \alpha_\sigma \sigma)] \subseteq B_e\left(\Pi[g(0)], \frac{1}{8}\text{diam}\Pi[B_h(g(0), \sigma)]\right)$$

where the inclusion follows from Lemma 8.11. Note that condition (b) of our Lemma has thus been established.

Assume now that $l_1 > \sigma/\tau$. Then $d(h(0), 0) > \sigma$ for all $h \in \Gamma_1$ and we can apply the Geometric Distortion Lemma (Lemma 2.2 in [37]) to get

$$(8.10) \quad d(g(0), 0) + d(h(0), 0) - 2\sigma \leq d(gh(0), 0) \leq d(g(0), 0) + d(h(0), 0)$$


for every $h \in \Gamma_1$. Note that we also needed $g(\Pi[h(0)]) \in \Pi[B(g(0), \sigma)]$ to apply the Geometric Distortion Lemma, but we already have that $g(\Pi[h(0)]) \in \Pi[B_h(g(0), \alpha_\sigma \sigma)] \subseteq \Pi[B_h(g(0), \sigma)]$. Now since $h(0) \in A_{l_1}(\tau)$, for every $h \in \Gamma_1$, we have that $l_1\tau \leq d(h(0), 0) < l_1\tau + \tau$. Hence, we get

$$(8.11) \quad d(g(0), 0) + l_1\tau - 2\sigma \leq d(gh(0), 0) \leq d(g(0), 0) + l_1\tau + \tau$$

for every $h \in \Gamma_1$. Now define $\Gamma(g)$ to be a maximal (in the sense of inclusion) subset of Γ_1 such that

$$(8.12) \quad d(h_1(0), h_2(0)) > \tau + 6\sigma$$

for all $h_1, h_2 \in \Gamma_1$ with $h_1 \neq h_2$. Let us now prove item (a). Suppose by way of contradiction that, $\Pi[B_h(gh_1(0), \sigma)] \cap \Pi[B_h(gh_2(0), \sigma)] \neq \emptyset$ for some $h_1, h_2 \in \Gamma_1$ with $h_1 \neq h_2$. This means that $B(gh_1(0), \sigma)$ intersects the light-cone viewed from 0 generated by $B_h(gh_2(0), \sigma)$. Let z_1 belong to this intersection and let $z_2 \in B_h(gh_2(0), \sigma)$ be chosen on the ray from 0 to z_1 whose endpoint we denote by ξ .



Then the following estimates are true using (8.11):

$$d(g(0), 0) + l_1\tau - 3\sigma < d(z_1, 0) < d(g(0), 0) + l_1\tau + \tau + \sigma$$

and

$$d(g(0), 0) + l_1\tau - 3\sigma < d(z_2, 0) < d(g(0), 0) + l_1\tau + \tau + \sigma.$$

Therefore

$$\begin{aligned} d(z_1, z_2) &= |d(0, z_1) - d(0, z_2)| \\ &< [d(g(0), 0) + l_1\tau + \tau + \sigma] - [d(g(0), 0) + l_1\tau - 3\sigma] \\ &= \tau + 4\sigma. \end{aligned}$$

Therefore we have that

$$\begin{aligned} d(h_1(0), h_2(0)) &= d(gh_1(0), gh_2(0)) \\ &\leq d(gh_1(0), z_1) + d(z_1, z_2) + d(z_2, gh_2(0)) \\ &< \sigma + [\tau + 4\sigma] + \sigma \\ &= \tau + 6\sigma \end{aligned}$$

which contradicts (8.12) and finishes the proof of item (a). It follows directly from the left-hand side of (8.11) and Lemma 2.1 in [37] that with $l_1 \geq 1$ chosen large enough the right-hand side of (c) holds. Similarly the left-hand side of (c) follows from the right-hand side of (8.11) and Lemma 2.1 in [37].

We have only left to show item (d) and this where the final specification of l_1 will be made. First for every $h \in \Gamma(g)$, define

$$Q_g(h) := \{f \in \Gamma(g) : d(f(0), h(0)) \leq \tau + 6\sigma\}.$$

By maximality of $\Gamma(g)$, see (8.12), we have that

$$\bigcup_{h \in \Gamma(g)} Q_g(h) = \Gamma_1.$$

Now since our group is strongly discrete, the number of elements in $Q_g(h)$ is bounded above by some constant, $C_1 = C_1(\tau, \sigma)$, depending only on $(\tau + 6\sigma)$ and of course on G . Also because of (8.11) and Lemma 2.1 in [37], there exists another constant, $C_2 = C_2(\tau, \sigma)$, such that for every $g \in G$ with $d(g(0), 0) > \sigma$ and all $h_1, h_2 \in \Gamma_1$,

$$C_2^{-1} \leq \frac{\text{diam}(\Pi[B_h(gh_2(0), \sigma)])}{\text{diam}(\Pi[B_h(gh_1(0), \sigma)])} \leq C_2.$$

Hence,

$$\begin{aligned} \sum_{h \in \Gamma_1} \text{diam}^s(\Pi[B_h(gh(0), \sigma)]) &\leq \sum_{h \in \Gamma(g)} \sum_{f \in Q_g(h)} \text{diam}^s(\Pi[B_h(gf(0), \sigma)]) \\ &\leq \sum_{h \in \Gamma(g)} C_2(\tau, \sigma)^s \sum_{f \in Q_g(h)} \text{diam}^s(\Pi[B(gh(0), \sigma)]) \\ (8.13) \quad &= C_2(\tau, \sigma)^s \sum_{h \in \Gamma(g)} [\#Q_g(h)] \text{diam}^s(\Pi[B(gh(0), \sigma)]) \\ &\leq C_2(\tau, \sigma)^s C_1(\tau, \sigma) \sum_{h \in \Gamma(g)} \text{diam}^s(\Pi[B_h(gh(0), \sigma)]). \end{aligned}$$

Now take $M = C_\sigma^{2s} C_2(\tau, \sigma)^s C_1(\tau, \sigma)$, where $C_\sigma > 0$ comes from Lemma 2.1 in [37], and choose $l_1 \geq 1$ to be one of the numbers $(n_j(\xi_1))_{j=1}^\infty$ appearing in Lemma 8.10 that is so large as required above, viz. that $l_1 > \sigma/\tau$. Now by Lemma 8.10, Lemma 2.1 in [37] and the right-hand-side of (8.10), we get that

$$\begin{aligned} \sum_{h \in \Gamma_1} \text{diam}^s(\Pi[B_h(gh(0), \sigma)]) &\geq C_\sigma^{-s} \sum_{h \in \Gamma_1} e^{-sd(gh(0), 0)} \\ &\geq C_\sigma^{-s} \sum_{h \in \Gamma_1} e^{-sd(g(0), 0)} e^{-sd(h(0), 0)} \\ &= C_\sigma^{-s} e^{-sd(g(0), 0)} \sum_{h \in \Gamma_1} e^{-sd(h(0), 0)} \\ &\geq M C_\sigma^{-s} e^{-sd(g(0), 0)} \\ &\geq M C_\sigma^{-2s} \text{diam}^s(\Pi[B(g(0), \sigma)]) \\ &\geq C_2(\tau, \sigma)^s C_1(\tau, \sigma) \text{diam}^s(\Pi[B(g(0), \sigma)]). \end{aligned}$$

Now inserting this into (8.13), we finally get

$$\sum_{h \in \Gamma(g)} \text{diam}^s(\Pi[B(gh(0), \sigma)]) \geq \text{diam}^s(\Pi[B(g(0), \sigma)])$$

which establishes (d) and finished the proof of our Lemma. \square

We are now in a position to prove the main result of this section, which is the extension of the Bishop-Jones result to the infinite-dimensional non-proper case.

Theorem 8.13. *If G is a strongly discrete group acting on \mathbb{B} , then*

$$\text{HD}(L_r(G)) = \text{HD}(L_{ur}(G)) = \delta_G.$$

Proof. As $L_{ur}(G) \subseteq L_r(G)$, we have that

$$(8.14) \quad \text{HD}(L_{ur}(G)) \leq \text{HD}(L_r(G)).$$

We shall first show that

$$\text{HD}(L_r(G)) \leq \delta_G.$$

If $\delta_G = +\infty$, then we are done and so let's assume that $\delta_G < +\infty$. Fix an arbitrary $s > \delta_G$. Write G as $(g_n)_{n=1}^\infty$. Fix $\sigma > \log 2$ and let

$$L_{r,\sigma}(G) := \bigcap_{n \geq 1} \bigcup_{k \geq n} \Pi[B_h(g_k(0), \sigma)].$$

Since $\sum_{n \geq 1} e^{-sd(g_n(0),0)} < +\infty$, we get that

$$\lim_{n \rightarrow \infty} \sum_{k \geq n} \text{diam}^s(\Pi[B_h(g_k(0), \sigma)]) \asymp_\sigma \lim_{n \rightarrow \infty} \sum_{k \geq n} e^{-sd(g_k(0),0)} = 0.$$

Thus $\text{HD}(L_{r,\sigma}(G)) \leq s$ and consequently that $\text{HD}(L_r(G)) \leq \delta_G$. since also $L_r(G) = \bigcup_{n \geq 3} L_{r,n}(G)$, using the σ -stability of Hausdorff dimension, we therefore get

$$\text{HD}(L_r(G)) = \sup_{n \geq 3} \{\text{HD}(L_{r,n}(G))\} \leq \delta_G.$$

Along with (8.14) this gives $\text{HD}(L_{ur}(G)) \leq \text{HD}(L_r(G)) \leq \delta_G$, and we are left to show that

$$(8.15) \quad \text{HD}(L_{ur}(G)) \geq \delta_G.$$

By means of Lemma 8.12 we will perform a construction to which Proposition 8.2 will apply. In the setting of Lemma 8.12, for every $n \geq 1$ let

$$E_n := E := \Gamma_1 \cup \Gamma_2.$$

We define the set $\Sigma \subseteq E^*$ and the sets $A(\omega)$ for $\omega \in \Sigma$ by induction with respect to word length in Σ . For the base of your recursion, we take $E \cap \Sigma := E := \Gamma_1 \cup \Gamma_2$ and $A(h) := \Pi[B(h(0), \sigma)]$ for all $h \in E$. For the inductive step, suppose that the set $E^n \cap \Sigma$ has been defined and that all the sets $A(f)$, $f \in E^n \cap \Sigma$ have been defined as well. To define $E^{n+1} \cap \Sigma$ consider all the elements $g = f_1, \dots, f_n \in E^n \cap \Sigma$ and declare that $f = f_1, \dots, f_{n+1} \in E^{n+1} \cap \Sigma$ if $f_{n+1} \in E$ and $f_{n+1} \in \Gamma(f_1 \circ \dots \circ f_n) = \Gamma(g)$. Then put $A(f) = \Pi[B(f_1 \circ \dots \circ f_{n+1}(0), \sigma)]$. Verifying now the hypotheses of Proposition 8.2, we see that $\widehat{\Sigma} \subseteq \Sigma$ directly by construction. Properties (c), (d), (b) and (a) of Proposition 8.2 follow respectively from property (a) of Lemma 8.12; property (d) of Lemma 8.12; the left-hand side of property (c) of Lemma 8.12 and finally from both property (b) and the right-hand-side of Property (c) of Lemma 8.12 with $\kappa = 1/4$. Therefore all the properties of Proposition 8.2 have been verified and as a result of applying it we get that

$$\text{HD}\left(\bigcap_{n \geq 1} \bigcup_{\omega \in E^n \cap \Sigma} A(\omega)\right) \geq s.$$

It now follows from (c) of Lemma 8.12, or more directly from (8.11), that

$$\bigcap_{n \geq 1} \bigcup_{\omega \in E^n \cap \Sigma} A(\omega) \subseteq L_{ur}(G),$$

and so we have that $\text{HD}(L_{ur}(G)) \geq s$. Since s was arbitrarily smaller than δ_G , we therefore get that $\text{HD}(L_{ur}(G)) \geq \delta_G$. This means that (8.15) has been established and we are done. \square

9. CONVEX-COBOUNDED GROUPS


With the proof of the main theorem behind us, if only to whet the reader's appetite, we conclude with a proof sketch of the following theorem and encourage her to look up the papers [17, 10, 11].

Theorem 9.1. *Let $G < \text{Isom}(\mathbb{H})$ be strongly discrete and convex-cobounded. Then G is finitely generated, has finite Poincaré exponent $\delta < \infty$, is of divergence type, and has a compact limit set. The δ -dimensional Hausdorff and packing measures on $L(G)$ are finite and positive and coincide up to a multiplicative constant with the δ -conformal Patterson measure which is Ahlfors δ -regular.*

Proof (sketch). The proof that the group G is finitely generated and then showing that the orbit map is a quasi-isometry follows from the Milnor-Schwarz Lemma [2, Proposition I.8.19], once we notice that $L(G)$ being compact implies that \mathcal{C} is also compact, since we are in a Hilbert space. Say G has d generators, and let $|g|$ denote the word-length of $g \in G$, i.e. the length of the shortest word that g may be expressed in terms of the d generators. Then

$$\Sigma_s(G) = \sum_{g \in G} e^{-sd(0, g(0))} \leq \sum_{g \in G} e^{-s(\varepsilon|g| - K)} = e^{sK} \sum_{g \in G} e^{-s\varepsilon|g|} \leq e^{sK} \sum_{g \in F_d} e^{-s\varepsilon|g|},$$

where ε, K are the quasi-isometry constants and F_d denotes the free group on d generators which surjects onto G . By taking $s\varepsilon$ to be sufficiently large we can force convergence of the final sum above. Thus $\delta < \infty$. Next let us prove that G is of compact type. Consider a sequence $x_n \in G(0)$ with $d(0, x_n) \rightarrow \infty$. Choose $y_n \in \partial B(0, N) \cap [0, x_n]$ as in the figure below.



Since our group is convex-cobounded there exists a sequence $z_n \in G(0)$ with $d(0, z_n) \leq N + R$. Since the action is strongly discrete there are only finitely many such z_n s for

every n and therefore we may extract a constant subsequence $(n_k)_k$ with $z_{n_k} = z$. Thus $d(y_{n_k}, y_{n_l}) \leq 2R$ and so

$$\langle x_{n_k} | x_{n_l} \rangle_0 \geq \langle y_{n_k} | y_{n_l} \rangle_0 \geq \frac{1}{2}[N + N - 2R] = N - R .$$

Here $\langle x | y \rangle_z := \frac{1}{2}[d(x, z) + d(y, z) - d(x, y)]$ denotes the Gromov product, see [41, Definition 2.7]. Now since N are arbitrary, we may extract a diagonal sequence such that for every $k, l \in \mathbb{N}$

$$\langle x_{n_k} | x_{n_l} \rangle_0 \geq \min\{k, l\} - R \xrightarrow[k, l]{} \infty .$$

Thus $(x_{n_k})_k$ is a Gromov sequence [41, Section 5] whose distances from the origin become arbitrarily large and thus we have convergence to a limit point on the boundary. Thus $L(G)$ is compact by Theorem 6.5.

Note that the compactness of the limit set allows the usual Patterson-Sullivan machinery via weak limits to go through, see [38, Theorem 1]. The Ahlfors regularity of the Patterson-Sullivan measure (and thus its equivalence with Hausdorff and packing measures) follows from a well-known argument using Sullivan's shadow lemma; see [38, Proposition 3] and [27, Section 8]. Finally, since Theorem ?? shows that the Patterson-Sullivan measure is supported on $L_r(G)$, the easy direction of the well-known Ahlfors-Thurston-Tukia argument (viz. μ is s -conformal implies that $\Sigma_s(G) = \infty$, see [33, Theorem 8.2.2 and 8.2.3]) shows that the G is of divergence type. This completes the proof. \square

10. ACKNOWLEDGEMENTS

The basic ideas behind these results were obtained by the authors during the summer of 2009 at the end of a productive conference, Dynamical Systems II, hosted at the University of North Texas in Denton. We thank our colleagues for their enthusiasm and patience in spite of the long delays in writing these notes up. TD would like to thank his undergraduate advisor BOS for introducing him to the charmed world of negative curvature, and his graduate advisor MU for suggesting that he explore the uncharted territory of Kleinian groups acting on Hilbert spaces. The proof of the Bishop-Jones relation appeared in [9]. TD thanks them both for their inspiration and guidance over the years.

REFERENCES

- [1] C. J. Bishop, P. W. Jones, Hausdorff dimension and Kleinian groups. *Acta Math.* 179 (1997), no. 1, 1-39.
- [2] M. R. Bridson, A. Haefliger, *Metric spaces of non-positive curvature*. Grundlehren der Mathematischen Wissenschaften, 319. Springer-Verlag, Berlin, 1999.
- [3] M. Burger, A. Iozzi, N. Monod, Equivariant embeddings of trees into hyperbolic spaces. *IMRN* (2005), no. 22, 1331-1369.
- [4] J. W. Cannon, W. J. Floyd, R. Kenyon, W. Parry and R. Walter, *Hyperbolic geometry*. *Flavors of geometry*, 59-115, *Math. Sci. Res. Inst. Publ.*, 31, Cambridge Univ. Press, Cambridge, 1997.
- [5] S. Cantat, S. Lamy, Normal subgroups of the Cremona group, *Acta Mathematica*, 210 (2013), 31-94.
- [6] P-A. Cherix, M. Cowling, P. Jolissaint, P. Julg, A. Valette, *Groups with the Haagerup property. Gromov's a-T-menability*. *Progress in Mathematics*, 197. Birkhuser Verlag, Basel, 2001.
- [7] K. Corlette, A. Iozzi, Limit sets of discrete groups of isometries of exotic hyperbolic spaces. *Trans. AMS* 351 (1999), 1507-1530.
- [8] S.G. Dani, Bounded orbits of flows on homogeneous spaces. *Comment. Math. Helv.* 61 (1986), 636-660.

- [9] T. Das, Kleinian groups in Hilbert spaces. Ph. D. Thesis, University of North Texas, 2012.
- [10] T. Das, D. S. Simmons, M. Urbański, Geometry and dynamics in Gromov hyperbolic metric spaces I, in preparation.
- [11] T. Das, D. S. Simmons, M. Urbański, Geometry and dynamics in Gromov hyperbolic metric spaces II: Thermodynamic formalism, in preparation.
- [12] T. Delzant, P. Py, Kähler groups, real hyperbolic spaces and the Cremona group. *Compos. Math.* 148 (2012),no.1,153-184.
- [13] M. Edelstein, On non-expansive mappings of Banach spaces, *Proc. Cambridge Philos. Soc.* 60 (1964), 439-447.
- [14] K. Falconer, *Techniques in fractal geometry*. John Wiley & Sons, Ltd., Chichester, 1997.
- [15] K. Falk and B. O. Stratmann, Constructing restricted Patterson measures for geometrically infinite Kleinian groups. *Acta Math. Sin. (Engl. Ser.)* 22 (2006), no. 2, 431-446.
- [16] K. Fan, Invariant subspaces of certain linear operators, *Bull. Amer. Math. Soc. Vol. 69, No. 6* (1963), 773-777.
- [17] L. Fishman, D. S. Simmons, M. Urbański, Diophantine approximation and the geometry of limit sets in Gromov hyperbolic metric spaces, preprint (2013). <http://arxiv.org/abs/1301.5630>
- [18] I. Garrido, J. A. Jaramillo, Y. C. Rangel, Algebras of differentiable functions on Riemannian manifolds, *Bull. Lond. Math. Soc.* 41 (2009), no. 6, 993-1001.
- [19] M. Gromov, *Asymptotic Invariants of Infinite Groups*, Geometric group theory, Vol. 2 (Sussex, 1991), 1-295, London Math. Soc. Lecture Note Ser., 182, Cambridge Univ. Press, Cambridge, 1993.
- [20] G. A. Hedlund, Fuchsian groups and transitive horocycles. *Duke Math. J.* 2 (1936), no. 3, 530-542.
- [21] S. Hersensky, F. Paulin, Hausdorff dimension of Diophantine geodesics in negatively curved manifolds. *J. Reine Angew. Math.* 539 (2001), 29-43.
- [22] C. Horbez, A short proof of Handel and Mosher's alternative for subgroups of $\text{Out}(F_N)$, preprint (2014).
- [23] M. Huff, Conformal maps on Hilbert space, *Bull. Amer. Math. Soc. Volume 82, Number 1* (1976), 147-149.
- [24] M. Kapovich, Kleinian groups in higher dimensions. *Geometry and dynamics of groups and spaces*, 487-564, *Progr. Math.*, 265, Birkhäuser, Basel, 2008.
- [25] A. Karlsson, Nonexpanding maps and Busemann functions, *Ergodic Theory Dynam. Systems* 21 (2001), 1447-1457.
- [26] A. Karlsson and G. A. Noskov, Some groups having only elementary actions on metric spaces with hyperbolic boundaries, *Geom. Dedicata* 104 (2004), 119-137.
- [27] R. D. Mauldin, T. Szarek and M. Urbański, Graph Directed Markov Systems on Hilbert Spaces, *Math. Proc. Cambridge Phil. Soc.* 147 (2009), 455-488.
- [28] R. D. Mauldin, M. Urbański, *Graph Directed Markov Systems*, Cambridge Univ. Press, 2003.
- [29] S. Lang, *Introduction to differentiable manifolds*. Interscience Publishers, New York-London 1962.
- [30] Manin, Yu. I. *Cubic forms: algebra, geometry, arithmetic*. Translated from the Russian by M. Hazewinkel. North-Holland Mathematical Library, Vol. 4. North-Holland Publishing Co., Elsevier, New York, 1974.
- [31] N. Monod, P. Py, An exotic deformation of the hyperbolic space, preprint (2012). <http://arxiv.org/abs/1201.0606>
- [32] R. Nevanlinna, *On differentiable mappings*, Analytic Functions, Princeton University Press, N. J., 1960.
- [33] P. J. Nicholls, *The ergodic theory of discrete groups*. London Mathematical Society Lecture Note Series 143. Cambridge University Press, Cambridge, 1989.
- [34] S. J. Patterson, The limit set of a Fuchsian group. *Acta Math.* 136 (1976), no. 3-4, 241-273.
- [35] F. Paulin, On the critical exponent of a discrete group of hyperbolic isometries. *Diff. Geom. Appl.* 7 (1997), 231-236.
- [36] J. G. Ratcliffe, *Foundations of hyperbolic manifolds*. 2ed. GTM 149. Springer, New York, 2006.

- [37] B. O. Stratmann, The exponent of convergence of Kleinian groups: on a theorem of Bishop and Jones, *Mathematica Gottingensis* 1 (2001) 1-16.
- [38] D. Sullivan, The density at infinity of a discrete group of hyperbolic motions, *IHÉS Publ. Math.* 50 (1979) 171-202.
- [39] D. Sullivan, Entropy, Hausdorff measures old and new, and limit sets of geometrically finite Kleinian groups, *Acta Math.* 153 (1984) 259-277.
- [40] D. Sullivan, Seminar on conformal and hyperbolic geometry, *IHÉS*, 1982.
- [41] J. Väisälä, Gromov hyperbolic spaces, *Expositiones Mathematicae* 23 (2005), no. 3, 187-231.
- [42] A. Valette, Affine isometric actions on Hilbert spaces and amenability, to appear in *Proc. Special semester on amenability*, Erwin Schrödinger Institute, Vienna, March 2007.

TUSHAR DAS, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN - LA CROSSE, LA CROSSE, WI 54601, USA.

E-mail address: `tdas@uwlax.edu`

BERND O. STRATMANN, FACHBEREICH 3 - MATHEMATIK, UNIVERSITÄT BREMEN, POSTFACH 33 04 40, BIBLIOTHEKSTRASSE 1, 28359 BREMEN, GERMANY.

E-mail address: `bos@math.uni-bremen.de`

MARIUSZ URBAŃSKI, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NORTH TEXAS, P.O. BOX 311430, DENTON, TX 76203-1430, USA.

E-mail address: `urbanski@unt.edu`